

**ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, PHENYLEPHRINE HCL-  
acetaminophen, chlorpheniramine maleate, phenylephrine hcl tablet  
OPMX LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

**Pentrexilina - Daytime  
(Acetaminophen 500mg, Chlorpheniramine Maleate 4mg, Phenylephrine HCl 10mg Tablet)**

Acetaminophen 500mg

Chlorpheniramine Maleate 4mg

Phenylephrine HCl 10mg

Pain reliever/fever reducer

Antihistamine

Nasal decongestant

magnesium stearate, povidone, silicon dioxide, sodium benzoate, sodium starch glycolate, starch, talc

- Do not take more than directed - see overdose warning
- swallow whole; do not crush, chew or dissolve
- do not exceed 6 tablets per 24 hours

---

|  |   |
|--|---|
| Adults and children 12 years of age & over | 1 tablet with water every 4-6 hours as needed |
| Children 4 to 12 years of age              | Ask a doctor                                  |
| Children under 4 years of age              | Do not use                                    |

---

Temporarily relieves these symptoms of hay fever or other respiratory allergies:

- Headache
- Nasal congestion
- Sinus congestion & pressure
- Runny nose and sneezing
- Minor aches & pain

Temporarily relieves these additional symptoms of hay fever:

- itching of the nose or throat
- itchy, watery eyes
- helps clear nasal passages
- helps decongest sinus opening and passages

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take :

- more than 6 tablets in 24 hours, which is the maximum daily amount for this product
- with other drugs contains acetaminophen
- 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult doctor promptly

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- Liver disease
- High blood pressure
- Thyroid disease
- Diabetes
- Trouble urinating due to enlarged prostate gland
- A breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if:

You get nervous, dizzy, or sleepless

Pain, nasal congestion, or cough gets worse or lasts more than 7 days

Fever gets worse or last more than 3 days

Redness or swelling is present

New symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding: ask a health professional before use.

Keep out of reach of children.

Keep out of reach of children.

**OVERDOSE WARNING:** Read directions carefully. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

store at room temperature between 20-25°C (68-77°F)

avoid excessive heat and humidity

**Uses:** temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ headache ■ sinus congestion & pressure ■ nasal congestion ■ runny nose and sneezing ■ minor aches and pains ■ temporarily relieves these additional symptoms of hay fever: ■ itching of the nose or throat ■ itchy, watery eyes ■ helps clear nasal passages ■ helps decongest sinus openings and passages

**Warnings: Liver Warnings:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 6 tablets in 24 hrs, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ Skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Directions:**

- Do not take more than directed - See Overdose warning
- Swallow whole; do not crush, chew or dissolve
- Do not exceed 6 tablets per 24 hours

|   |  |
|---|--|
| Adults and children<br>12 years of age & over | 1 tablet with water<br>every 4-6 hours as needed |
| Children 4 to 12 years of age                 | Ask a doctor                                     |
| Children under 4 years of age                 | Do not use                                       |

**Other information:** ■ Store at room temperature 20°-25°C (68°-77°F)  
■ Avoid excessive heat and humidity

**READ AND KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

Exclusively distributed by  
**OPMX**  
San Diego, CA 92164  
Phone: 619-460-4632

Made for:  
**RRX**  
San Diego, CA

REV. 09/2019

TAMPER EVIDENT: DO NOT  
USE IF POUCH IS TORN,  
BROKEN OR SHOWS ANY  
SIGN OF TAMPERING



2 Tablets

**DAYTIME**  
**PENTREXCILINA**

| ACTIVE INGREDIENTS (in each tablet) | PURPOSE                     |
|-------------------------------------|-----------------------------|
| Acetaminophen 500 mg                | Pain reliever/Fever reducer |
| Chlorpheniramine Maleate 4 mg       | Antihistamine               |
| Phenylephrine HCl 10 mg             | Nasal decongestant          |

Effective and quick relief from  
**cold & flu symptoms**

- Headache
- Nasal Congestion
- Itchy, Watery Eyes
- Runny Nose & Sneezing



**Drug Facts (continued)**

**Other information:** • Store at room temperature between 20° - 25° C (68° - 77° F) • Avoid excessive heat and humidity

**Inactive ingredients:** magnesium stearate, polydione, silicon dioxide, sodium benzoate, sodium starch glycolate, starch, talc

**Questions or comments?** Call: (619) 600-5632 (Mon-Fri 9am-5pm EST) or <https://www.opmx.us>

**TAMPER EVIDENT: DO NOT USE IF POUCH IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING**

**Effective and Quick Relief from**

- Headache
- Nasal Congestion
- Itchy, Watery Eyes
- Runny Nose & Sneezing

**PENTREXILINA®**

Acetaminophen 500 mg • Chlorpheniramine Maleate 4 mg • Phenylephrine HCl 10 mg

**DAYTIME**

**COLD & FLU**

12 Tablets (6 Pouches of 2 Tablets)

NDC 69729-223-37

Model for: **OPMX** Pharmaceuticals, Inc. San Diego, CA

8 5822 0824 00024 (08/2016)

1.349 in

**READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT INFORMATION**

**Drug Facts**

| ACTIVE INGREDIENTS (in each tablet) | PURPOSE                     |
|-------------------------------------|-----------------------------|
| Acetaminophen 500 mg                | Pain reliever/fever reducer |
| Chlorpheniramine Maleate 4 mg       | Antihistamine               |
| Phenylephrine HCl 10 mg             | Nasal decongestant          |

**Uses:**

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies: • headache • sinus congestion & pressure • nasal congestion
- runny nose and sneezing • minor aches and pains • temporarily relieves these additional symptoms of hay fever:
- itching of the nose or throat • itchy, watery eyes • helps clear nasal passages • helps decongest sinus openings and passages

**Warnings:**

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 tablets in 24 hrs, which is the maximum daily amount for this product • with other drugs containing acetaminophen
- 2 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • rash

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use:** • with any other drug containing acetaminophen (prescription or nonprescription) • if you are not sure whether a drug contains acetaminophen

**Ask a doctor or pharmacist:** • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug • if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product • if you have ever had an allergic reaction to this product or any of its ingredients • to make a child sleep

**Ask a doctor before use if you have:** • heart disease • heart disease • high blood pressure • thyroid disease • glaucoma • diabetes • trouble urinating due to an enlarged prostate gland • a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or a pharmacist before use if you are:** • taking the blood thinning drug warfarin • taking sedatives or tranquilizers

**When using this product:** • do not use more than directed • drowsiness may occur, especially in children • drowsiness may occur • alcohol, sedatives and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if:** • nervousness, dizziness, or sleeplessness occur • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur. These could be signs of a serious condition

**If pregnant or breast-feeding:** ask a health professional before use. **Keep out of reach of children.**

**Overdose warning:** Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions:**

|   |  |   |
|---|--|---|
| • Do not take more than directed - See Overdose warning | Adults and children 12 years of age & over | 1 tablet with water every 4-6 hours as needed |
| • Swallow whole; do not crush, chew or dissolve         | Children 4 to 12 years of age              | Ask a doctor                                  |
| • Do not exceed 6 tablets per 24 hours                  | Children under 4 years of age              | Do not use                                    |



## ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, PHENYLEPHRINE HCL

acetaminophen, chlorpheniramine maleate, phenylephrine hcl tablet

### Product Information

|                         |                |                    |               |
|-------------------------|----------------|--------------------|---------------|
| Product Type            | HUMAN OTC DRUG | Item Code (Source) | NDC:69729-222 |
| Route of Administration | ORAL           |                    |               |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength           | Strength |
|--|-----------------------------|----------|
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 10 mg    |
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)               | ACETAMINOPHEN               | 500 mg   |
| CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U) | CHLORPHENIRAMINE MALEATE    | 4 mg     |

### Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| SODIUM BENZOATE (UNII: OJ245FE5EU)                       |          |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) |          |
| TALC (UNII: 7SEV7J4R1U)                                  |          |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4)                       |          |
| MAGNESIUM STEARATE (UNII: 70097M6I30)                    |          |
| STARCH, CORN (UNII: O8232NY3SJ)                          |          |
| POVIDONE K30 (UNII: U725QWY32X)                          |          |

### Product Characteristics

|          |                  |              |          |
|----------|------------------|--------------|----------|
| Color    | white            | Score        | no score |
| Shape    | ROUND (BICONVEX) | Size         | 13mm     |
| Flavor   |                  | Imprint Code | S78      |
| Contains |                  |              |          |

### Packaging

| # | Item Code        | Package Description                               | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:69729-222-27 | 2 in 1 POUCH; Type 0: Not a Combination Product   | 08/26/2019           |                    |
| 2 | NDC:69729-222-37 | 12 in 1 POUCH; Type 0: Not a Combination Product  | 08/26/2019           |                    |
| 3 | NDC:69729-222-38 | 144 in 1 POUCH; Type 0: Not a Combination Product | 08/26/2019           |                    |

### Marketing Information

| Marketing Category  | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341                                  | 08/26/2019           |                    |

**Labeler** - OPMX LLC (029918743)

Revised: 11/2020

OPMX LLC