

**ALLERGY RELIEF D- cetirizine hydrochloride, pseudoephedrine hydrochloride tablet, film coated, extended release**  
**H E B**

-----  
**HEB Allergy Relief-D Drug Facts**

**Active ingredients (in each extended release tablet)**

Cetirizine HCl 5 mg

Pseudoephedrine HCl 120 mg

**Purpose**

Antihistamine

Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease

- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are**

taking tranquilizers or sedatives.

**When using this product**

- **do not use more than directed**
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

**If pregnant or breast-feeding:**

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

## Other information

- store between 20° to 25°C (68° to 77°F)
- **do not use if blister unit is broken or torn**
- see side panel for lot number and expiration date
- meets USP *Dissolution Test 2*

## Inactive ingredients

colloidal silicon dioxide, hypromellose, lactose monohydrate, low-substituted hydroxypropyl cellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

## Questions or comments?

**1-800-719-9260**

## Package/Label Principal Display Panel

Compare to Zyrtec-D<sup>®</sup> active ingredients

Allergy Relief-D

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride

Extended Release Tablets, 5 mg/120 mg

Antihistamine/Nasal Decongestant

Allergy & Congestion

12 Hour

Indoor & Outdoor Allergies

12 Hour Relief of:

Sneezing

Itchy, Watery Eyes

Runny Nose

Itchy Throat or Nose

Sinus Pressure

Nasal Congestion

actual size

12 EXTENDED RELEASE TABLETS

**H-E-B**

# Allergy Relief-D

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride  
Extended Release Tablets, 5 mg/120 mg  
Antihistamine/Nasal Decongestant

Allergy & Congestion

Compare to Zyrtec-D® active ingredients\*

NDC 37808-651-53

**H-E-B**

# Allergy Relief-D

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride  
Extended Release Tablets, 5 mg/120 mg  
Antihistamine/Nasal Decongestant

Allergy & Congestion



actual size

Indoor & Outdoor Allergies

12 Hour Relief of:

- Sneezing
- Runny Nose
- Sinus Pressure
- Itchy, Watery Eyes
- Itchy Throat or Nose
- Nasal Congestion

**12 EXTENDED RELEASE TABLETS**

### Drug Facts

**Active ingredients**  
(in each extended release tablet)  
Cetirizine HCl 5 mg ..... Antihistamine  
Pseudoephedrine HCl 120 mg ..... Nasal decongestant

### Purpose

### Uses

- temporarily relieve the severe symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- nasal congestion
- reduced swelling of nasal passages
- temporarily relieve sinus congestion and pressure
- temporarily restores free breathing through the nose

### Warnings

- Do not use
- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
  - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or molarol conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

### Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

- When using this product:
- do not use more than directed
  - drowsiness may occur
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery

### Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or dizzy
- symptoms do not improve within 7 days or are accompanied by fever

### If pregnant or breast-feeding:

- if breast-feeding: not recommended
  - if pregnant: ask a health professional before use.
- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- do not break or chew tablet; swallow tablet whole
- |                                       |  |
|---------------------------------------|--|
| adults and children 12 years and over | take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours. |
| adults 65 years and over              | ask a doctor   |
| children under 12 years of age        | ask a doctor   |
| consume with liver or kidney disease  | ask a doctor   |

14753-1F-C2

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Zyrtec-D®.

If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.

**100% GUARANTEE promise**

MADE WITH PRIDE & CARE FOR H-E-B® SAN ANTONIO, TX 78204

### Drug Facts (continued)

### Other information

- store between 20° to 25°C (68° to 77°F)
- do not use if blister unit is broken or torn
- see side panel for lot number and expiration date
- meets USP Dissolution Test 2

**Inactive ingredients** colloidal silicon dioxide, hypromellose, lactose monohydrate, low-substituted hydroxypropyl cellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

Questions or comments? 1-800-719-9260

GLUTEN FREE

12966 - 2006



0 41220 27880 0

# ALLERGY RELIEF D

cetirizine hydrochloride, pseudoephedrine hydrochloride tablet, film coated, extended release

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37808-651
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CETIRIZINE HYDROCHLORIDE</b> (UNII: 64O047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	5 mg
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 2165RE0K14)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	L147
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-651-53	12 in 1 CARTON	02/24/2021	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:37808-651-62	24 in 1 CARTON	02/18/2021	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210719	02/18/2021	

**Labeler - H E B (007924756)**

Revised: 2/2021

H E B