LEADER TUSSIN DM- dextromethorphan hydrobromide, guaifenesin solution Cardinal Health 110, LLC. dba Leader

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Cardinal Health Tussin DM Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr, USP 20 mg Guaifenesin, USP 200 mg

Purposes

Cough suppressant Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- · this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- each 20 mL contains: sodium 14 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

LEADER™

Adult | Non-Drowsy

Tussin DM

Cough + Chest Congestion DM

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 200 mg

Cough Suppressant | Expectorant

Raspberry Flavor

For Ages 12 & Over

Controls Cough

Relieves Chest Congestion

Thins & Loosens Mucus

COMPARE TO ROBITUSSIN® COUGH+CHEST CONGESTION DM

active ingredients

100% Money Back Guarantee

4 FL OZ (118 mL)

SEE NEW DOSING



LEADER2 Adult | Non-Drowsy Tussin DM Cough + Chest Congestion DM

Dextromethorphan HBr, USP 20 mg Guaifenesin, USP 200 mg Cough Suppressant | Expectorant

Drug Facts

Active ingredients Purposes (in each 20 mL)
Dextromethorphan HBr,
USP 20 mg......C.
Guaifenesin, USP 200 mg....

.Cough suppressantExpectorant

Uses

- USES

 It temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold helps loosen phiegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MACI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MACI drug. If you do not know if your prescription drug contains an MACI, ask a doctor or pharmacist before taking this product.

- Ask a doctor before use if you have
- Ask a doctor before use if you have cough that occurs with too much phlegm (mucus) cough that last or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of

overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions

- do not take more than 6 doses in any

- do not take more than 6 doses in any 24-hour period
 measure only with dosing cup provided
 keep dosing cup with product
 ml. = milliliter
 this adult product is not intended for use in children under 12 years of age

Drug Facts (continued)

	age	dose
	adults and children 12 years and over	20 mL every 4 hours
П	children under 12 years	do not use

Other information

■ each 20 mL contains: sodium 14 mg ■ store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, FD&C red no. 40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-800-719-9260

DO NOT USE IF PRINTED HECKBAND IS BROKEN OR MISSING

"This product is not manufactured or distributed by Pfizer, the distributor of Robitussin® Cough+Chest Congestion DM. 02021 Cardinal Health, All Rights Reserved, CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

DISTRIBUTED BY CARDINAL HEALTH Dublin, Ohio 43017 CardinalHealth™ www.myleader.c 1-800-200-6313





www.StopMedicineAbuse.org

LEADER

NDC 70000-0596-1

Adult | Non-Drowsy

Congestion DM

Tussin DM Cough + Chest

Dextromethorphan HBr, USP 20 mg Guaifenesin, USP 200 mg Cough Suppressant | Expectorant

Raspberry Flavor For Ages 12 & Over

Controls Cough Relieves Chest Congestion

Thins & Loosens Mucus

COMPARE TO ROBITUSSIN® COUGH+CHEST CONGESTION DM active ingredients®

100% Money Back Guarantee

4FL OZ (118 mL)

SEE NEW DOSING



63K26 E9 C1

LEADER?

NDC 70000-0596-1

Adult | Non-Drowsy

Tussin DM Cough + Chest Congestion DM

Dextromethorphan HBr, USP 20 mg Guaifenesin, USP 200 mg Cough Suppressant | Expectorant

Raspberry Flavor For Ages 12 & Over

Controls Cough Relieves Chest Congestion

Thins & Loosens Mucus



COMPARE TO ROBITUSSIN® COUGH+CHEST CONGESTION DM

4FL OZ (118 mL)

SEE NEW DOSING

LEADER TUSSIN DM

dextromethorphan hydrobromide, quaifenesin solution

Product Information

HUMAN OTC DRUG NDC:70000-0596 **Product Type** Item Code (Source) **Route of Administration** ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 20 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)		
SORBITOL (UNII: 506T60A25R)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
XANTHAN GUM (UNII: TTV12P4NEE)		

Product Characteristics			
Color	RED	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:70000- 0596-1	1 in 1 CARTON	11/24/2021		
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product			
2	NDC:70000- 0596-2	1 in 1 CARTON	11/24/2021		
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/24/2021	