

DIPHENHYDRAMINE HYDROCHLORIDE 25MG- diphenhydramine hydrochloride 25mg tablet

Strive Pharmaceuticals Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

RIGHT REMEDIES Allergy Relief

Drug Facts

Active Ingredient (in each tablet)

Diphenhydramine HCl 25 mg

Purpose

Antihistamine

Uses

- temporarily relieves these symptoms due to hay fever or upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose and throat
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing

Warnings

Do not use

- to make a child sleepy
- with other products containing diphenhydramine, even ones used on skin

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- take every 4 to 6 hours, or as directed by doctor
- do not take more than 6 times in 24 hours

adults & children 12 years of age and over	take 1 to 2 tablets
children 6 years of age to under 12 years of age	take 1 tablet
children under 6 years of age	do not use

Other information

- **each tablet contains:** calcium 18.64 mg
- store at room temperature between 20 - 25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.

Inactive ingredients

croscarmellose sodium, dicalcium phosphate, D&C red #27 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, talc, titanium dioxide

Questions or comments?

1-888-577-8033 Monday - Friday 8am - 4pm EST

RIGHT REMEDIES

Compare to the active ingredient of **Benadryl® Allergy Ultratabs®***

Distributed by: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816

Product of India

Packaged & Quality Assured in USA

REV.00-102022

READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc..

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division., owner of the registered trademark Benadryl® Allergy Ultratabs®

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING FROM BOTTLE

Packaging

Packaging: 100 tablets

RIGHT REMEDIES †compare to active ingredient in **BENADRYL® ALLERGY ULTRATAB®**
NDC 70692-148-01

Allergy Relief
sneezing, runny nose,
itchy watery eyes,
itchy throat
diphenhydramine HCl **25mg** each caplet
100 caplets

RIGHT REMEDIES †compare to active ingredient in **BENADRYL® ALLERGY ULTRATAB®**
NDC 70692-148-01

Allergy Relief
sneezing, runny nose,
itchy watery eyes,
itchy throat
diphenhydramine HCl 25mg
100 caplets

RIGHT REMEDIES †compare to active ingredient in **BENADRYL® ALLERGY ULTRATAB®**
NDC 70692-148-01

Allergy Relief
sneezing, runny nose,
itchy watery eyes,
itchy throat
diphenhydramine HCl 25mg

READ AND KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

†This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark BENADRYL® ALLERGY ULTRATAB®

Distributed by: STRIVE PHARMACEUTICALS INC.
East Brunswick, NJ 08816
CT7069214801 - REV/00-092020
3 70692 13601 2
Lot. No.:
Exp. Date:
COATING FREE AREA

Drug Facts (continued)

Keep out of reach of children.
OVERDOSE WARNING: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- take every 4 to 6 hours, or as directed by doctor
- do not take more than 6 doses in 24 hours
- adults & children 12 years and over take 1 to 2 caplets
- children 6 to under 12 years take 1 caplet
- children under 6 years do not use

Other information

- store at room temperature between 20-25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.
- TAMPER EVIDENT: Do not use if imprinted safety seal is broken or missing.**

Inactive ingredients
crosscarmellose sodium, dicalcium phosphate DT, D&C red #27 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, talc, titanium dioxide.

Questions or comments?
Call 1 (888) 577-8033

Drug Facts

Active Ingredient (in each caplet) Diphenhydramine HCl 25 mg
Purpose Antihistamine

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
- runny nose
- sneezing

Warnings

- Do not use**
- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- avoid sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts Table

Drug Facts	
Active Ingredient (in each tablet)	Purpose
Diphenhydramine HCl 25 mg.....	Antihistamine
Uses	
<ul style="list-style-type: none"> ■ temporarily relieves these symptoms due to hay fever or upper respiratory allergies: <ul style="list-style-type: none"> ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose and throat ■ temporarily relieves these symptoms due to the common cold: <ul style="list-style-type: none"> ■ runny nose ■ sneezing 	
Warnings	
Do not use	
<ul style="list-style-type: none"> ■ to make a child sleepy ■ with other products containing diphenhydramine, even ones used on skin 	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland 	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers	
When using this product	
<ul style="list-style-type: none"> ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children 	
<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.</p>	

Drug Facts (continued)	
Directions	
<ul style="list-style-type: none"> ■ take every 4 to 6 hours, or as directed by doctor ■ do not take more than 6 times in 24 hours 	
adults & children 12 years of age and over	take 1 to 2 tablets
children 6 years of age to under 12 years of age	take 1 tablet
children under 6 years of age	do not use
Other information	
<ul style="list-style-type: none"> ■ each tablet contains: calcium 18.64 mg ■ store at room temperature between 20 - 25°C (68-77°F). ■ avoid excessive heat, cold and humidity. ■ close cap tightly after use. 	
Inactive Ingredients	
croscarmellose sodium, dicalcium phosphate, D&C red #27 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, talc, titanium dioxide	
Questions or comments?	
1-888-577-8033 Monday - Friday 8am - 4pm EST	

DIPHENHYDRAMINE HYDROCHLORIDE 25MG

diphenhydramine hydrochloride 25mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-148
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40)	DIPHENHYDRAMINE	25 mg

(DIPHENHYDRAMINE - UNII:8GTS82S83M)

HYDROCHLORIDE

25 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	

Product Characteristics

Color	pink	Score	no score
Shape	CAPSULE	Size	12mm
Flavor		Imprint Code	DP
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-148-01	100 in 1 PACKAGE; Type 0: Not a Combination Product	12/28/2018	
2	NDC:70692-148-46	1 in 1 CARTON	04/06/2023	
2		72 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/20/2018	

Labeler - Strive Pharmaceuticals Inc (080028013)

Revised: 4/2023

Strive Pharmaceuticals Inc