

## **DICLOFENAC SODIUM- diclofenac sodium topical gel, 1% gel**

### **Proficient Rx LP**

-----

### **Diclofenac Sodium Topical Gel, 1% (NSAID) - arthritis pain reliever**

#### **Active ingredient**

Diclofenac sodium (NSAID\*) 1%

\*nonsteroidal anti-inflammatory drug

#### **Inactive ingredients**

Inactive ingredients:

Carbomer Homopolymer Type C, Coco-Caprylate/caprate, Isopropyl Alcohol, Mineral Oil, Polyoxyl 20 Cetostearyl Ether, Propylene Glycol, Purified Water, Strong Ammonia Solution

#### **Purpose**

Arthritis pain reliever

#### **Uses**

- for the temporary relief of arthritis pain ONLY in the following areas:
  - o hand, wrist, elbow (upper body areas)
  - o foot, ankle, knee (lower body areas)
- this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use

#### **Warnings**

##### **For external use only**

**Allergy alert:** Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

• hives • asthma (wheezing) • skin reddening • blisters • facial swelling • shock • rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains diclofenac. Liver damage may occur if you apply

- more or for a longer time than directed
- when using other drugs containing diclofenac

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
- for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.
- right before or after heart surgery
- on more than 2 body areas at the same time
- in the eyes, nose or mouth

### **Ask a doctor before use if**

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- avoid contact with eyes, nose, or mouth
- if eye contact occurs, rinse thoroughly with water

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 21 days
- redness or swelling is present in the painful area
- fever occurs
- skin irritation occurs
- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding
- feel faint
- have bloody or black stools
- vomit blood
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke
- chest pain
- trouble breathing
- leg swelling
- weakness in one part or side of body
- slurred speech

### **If pregnant or breast-feeding**

ask a health professional before use. It is especially important not to use this product during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

Use up to 21 days unless directed by your doctor

Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

| <b>Daily</b>                                                                                                                                                                       | <b>Per Dose</b>                                                                                                                                                                                                                                                                        |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>For your arthritis pain:</b> <ul style="list-style-type: none"> <li>• Use 4 times per day every day</li> <li>• Do not use on more than 2 body areas at the same time</li> </ul> | <b>Use ENCLOSED DOSING CARD to measure a dose</b> <ul style="list-style-type: none"> <li>• For each upper body area (hand, wrist, or elbow) – Squeeze out 2.25 inches (2 grams)</li> <li>• For each lower body area (foot, ankle or knee) –Squeeze out 4.5 inches (4 grams)</li> </ul> |

**Read the enclosed User Guide for complete instructions:**

- use only as directed
- do not use more than directed or for longer than directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product
- do not apply with external heat such as heating pad
- do not apply a bandage over the treated area

store ENCLOSED DOSING CARD with your Diclofenac Sodium Topical Gel, 1% Arthritis Pain. The dosing card is re-usable.

**Other information**

- Store at 20-25°C (68°F - 77°F). Keep from freezing.
- read all product information before using. Keep the dosing card, the carton and accompanying User guide for important information.

**Inactive ingredients**

Carbomer homopolymer Type C, cocoyl caprylocaprate, isopropyl alcohol, mineral oil, polyoxyl 20 cetostearyl ether, propylene glycol, purified water, strong ammonia solution.

**Question and comments 1-833-285-4151**

**PRINCIPAL DISPLAY PANEL**

**Carton Label - NDC 71205-747-00**

Diclofenac Sodium Topical Gel, 1%

(NSAID)- Arthritis pain reliever

For external use only

For daily Treatment of Arthritis Pain Anti-Inflammatory

Net Wt 3.53 OZ (100 g)



Scan Here



NDC 71205-747-00

Relabeled By: Proficient Rx LP  
Thousand Oaks, CA 91320**Diclofenac Sodium 1%****3.53 OZ (100 g) Topical Gel**

Each tube contains: Diclofenac sodium (NSAID\*) 1%  
(equivalent to 0.93% diclofenac) Arthritis pain reliever  
\*nonsteroidal anti-inflammatory drug

See package insert

Product ID: SD074700

**For external use only**

Dist. By: Dr. Reddy's Laboratories, Inc. Princeton, NJ 08540 Made in India

Store at 20°C to 25°C (68°F to 77°F)

Keep medication out of the reach of children

Diclofenac Sodium 1%  
3.53 OZ (100 g) Topical Gel  
Lot #:00000 SN# MASTER  
NDC 71205-747-00 Exp:00/00/00

Diclofenac Sodium 1%  
3.53 OZ (100 g) Topical Gel  
Lot #:00000 SN# MASTER  
NDC 71205-747-00 Exp:00/00/00

Diclofenac Sodium 1%  
3.53 OZ (100 g) Topical Gel  
Lot #:00000 SN# MASTER  
NDC 71205-747-00 Exp:00/00/00



GTIN: 00371205747008  
SN# MASTER  
Exp. 00/00/00  
Lot #:00000

**Container Label**

Diclofenac Sodium Topical Gel, 1%

(NSAID)- Arthritis pain reliever

For external use only

For daily Treatment of Arthritis Pain Anti-Inflammatory

Net Wt 3.53 OZ (100 g)

**Dosing Card****Use Enclosed Dosing Card to Measure a Dose****How to Use**

- For arthritis pain only
- Use 4 times a day
- May take up to 7 days to work for your arthritis pain
- for use on no more than 2 body areas
- Use up to 21 days unless directed by your doctor

**DICLOFENAC SODIUM**

diclofenac sodium topical gel, 1% gel

**Product Information**

|                                |                |                           |                              |
|--------------------------------|----------------|---------------------------|------------------------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:71205-747(NDC:43598-977) |
| <b>Route of Administration</b> | TOPICAL        |                           |                              |

| Active Ingredient/Active Moiety                                                           |                  |                                                    |                      |                    |
|-------------------------------------------------------------------------------------------|------------------|----------------------------------------------------|----------------------|--------------------|
| Ingredient Name                                                                           |                  |                                                    | Basis of Strength    | Strength           |
| <b>DICLOFENAC SODIUM</b> (UNII: QTG126297Q) (DICLOFENAC - UNII:144O8QL0L1)                |                  |                                                    | DICLOFENAC SODIUM    | 10 mg in 1 g       |
|                                                                                           |                  |                                                    |                      |                    |
| Inactive Ingredients                                                                      |                  |                                                    |                      |                    |
| Ingredient Name                                                                           |                  |                                                    |                      | Strength           |
| <b>CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED)</b> (UNII: 4Q93RCW27E) |                  |                                                    |                      |                    |
| <b>COCOYL CAPRYLOCAPRATE</b> (UNII: 8D9H4QU99H)                                           |                  |                                                    |                      |                    |
| <b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)                                               |                  |                                                    |                      |                    |
| <b>MINERAL OIL</b> (UNII: T5L8T28FGP)                                                     |                  |                                                    |                      |                    |
| <b>POLYOXYL 20 CETOSTEARYL ETHER</b> (UNII: YRC528SWUY)                                   |                  |                                                    |                      |                    |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)                                                |                  |                                                    |                      |                    |
| <b>WATER</b> (UNII: 059QF0KO0R)                                                           |                  |                                                    |                      |                    |
| <b>Ammonia</b> (UNII: 5138Q19F1X)                                                         |                  |                                                    |                      |                    |
|                                                                                           |                  |                                                    |                      |                    |
| Packaging                                                                                 |                  |                                                    |                      |                    |
| #                                                                                         | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
| 1                                                                                         | NDC:71205-747-00 | 1 in 1 CARTON                                      | 01/18/2023           |                    |
| 1                                                                                         |                  | 100 g in 1 TUBE; Type 0: Not a Combination Product |                      |                    |
|                                                                                           |                  |                                                    |                      |                    |
| Marketing Information                                                                     |                  |                                                    |                      |                    |
| Marketing Category                                                                        |                  | Application Number or Monograph Citation           | Marketing Start Date | Marketing End Date |
| ANDA                                                                                      |                  | ANDA210986                                         | 09/14/2020           |                    |

**Labeler** - Proficient Rx LP (079196022)

| Establishment    |         |           |                     |
|------------------|---------|-----------|---------------------|
| Name             | Address | ID/FEI    | Business Operations |
| Proficient Rx LP |         | 079196022 | RELABEL(71205-747)  |

Revised: 1/2023

Proficient Rx LP