

RITE AID ANTACID AND PAIN RELIEF- aspirin, citric acid, sodium bicarbonate tablet, effervescent
Rite Aid

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rite Aid Antacid & Pain Relief

Active ingredients

(in each effervescent tablet)

Anhydrous citric acid 1000 mg

Aspirin 325 mg **(NSAID)***

Sodium bicarbonate (heat-treated) 1916 mg

*Nonsteroidal anti-inflammatory drug

Purpose

Anhydrous citric acid 1000 mg. Antacid

Aspirin 325 mg (NSAID)*. Pain reliever

Sodium bicarbonate (heat-treated) 1916 mg. Antacid

*Nonsteroidal anti-inflammatory drug

Uses

Uses for the temporary relief of

- heartburn, acid indigestion and sour stomach when accompanied by headache or body aches and pains
- upset stomach with headache from overindulgence in food or drink
- headache, body aches, and pain alone

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

■ presently taking a prescription drug. Antacids may interact with certain prescription drugs. ■ taking a prescription drug for anticoagulation (thinning the blood), diabetes, gout or arthritis

When using this product

When using this product do not take more than 8 tablets (adults) or 4 tablets (60 years and over) in a 24-hour period or use the maximum dosage for more than 2 weeks, except under the advice and supervision of a physician.

Stop use and ask a doctor if

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- new symptoms occur

- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

Keep out of reach of children.

In case of overdose

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage
 - to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking
- adults and children 12 years and over (up to 60 years of age) - 2 tablets every 4 hours
do not exceed 8 tablets in 24 hours
- 60 years and over - 2 tablets every 4 hours do not exceed 4 tablets in 24 hours
- Children under 12 years - consult a doctor

Other information

- **each tablet contains:** sodium 524 mg
- store at room temperature (59°-86°F)

Inactive ingredients

- none

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

PN: 42124 RA C2
R: 04/21



**SATISFACTION
GUARANTEE**
If you're not satisfied, we'll
happily refund your money.

This product is not manufactured,
distributed or licensed by
the maker of Alka-Seltzer®.

© 2011 Rite Aid
DISTRIBUTED BY:
RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011
MANUFACTURED IN THE U.S.A. WITH DOMESTIC
AND IMPORTED INGREDIENTS.
www.riteaid.com
Serious side effects associated with the use of this product can be reported to the
following address, or by calling the following 1-888-228-6937 phone number.

EFFERVESCENT
TABLETS
24

ORIGINAL
FLAVOR

Antacid & PAIN RELIEF
EFFERVESCENT TABLETS
Compare to the active ingredients in Alka-Seltzer®**
NDC 11822-4212-4
Anhydrous Citric Acid/Antacid
Aspirin (NSAID)/Pain Reliever
Sodium Bicarbonate/Antacid
Fast relief of: acid indigestion,
upset stomach, heartburn with
headache, body aches & pains



DO NOT USE IF PRINTED PACKETS
ARE TORN OR PUNCTURED

Drug Facts (continued)
Keep out of reach of children. In case
of overdose, get medical help or contact
a Poison Control Center right away.
Directions
■ do not exceed recommended dosage
■ to avoid serious injury, completely
dissolve tablets in 4 oz. of water
before taking
Adults and children
2 tablets every 4
hours do not
exceed 8 tablets in
24 hours
Up to 60 years of age
2 tablets every 4
hours do not
exceed 4 tablets in
24 hours
Children under 12 years
consult a doctor
Other information
■ each tablet contains: sodium 524 mg
■ store at room temperature (59°-86°F)
Inactive ingredients
■ none

ANTACID & PAIN RELIEF
EFFERVESCENT TABLETS
Anhydrous Citric Acid/Antacid
Aspirin (NSAID)/Pain Reliever
Sodium Bicarbonate/Antacid

Drug Facts
Active ingredients
(in each effervescent tablet)
Anhydrous citric acid 1000 mg Antacid
Aspirin 325 mg (NSAID) Pain Reliever
Sodium bicarbonate (heat-treated) 1916 mg Antacid
*Nonsteroidal anti-inflammatory drug
Purpose
Uses for the temporary relief of
■ heartburn, acid indigestion and sour stomach when
accompanied by headache or body aches and pains
■ upset stomach with headache from overindulgence in
food or drink
■ headache, body aches, and pain alone
Warnings
Reye's syndrome: Children and teenagers who have or
are recovering from chicken pox or flu-like symptoms
should not use this product. When using this product, if
changes in behavior with nausea and vomiting occur,
consult a doctor because these symptoms could be an
early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction
which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an
NSAID, which may cause severe stomach bleeding. The
chance is higher if you ■ are age 60 or older ■ have had
stomach ulcers or bleeding problems ■ take a blood
thinning (anticoagulant) or steroid drug ■ take other
drugs containing prescription or nonprescription NSAIDs
(aspirin, ibuprofen, naproxen, or others) ■ have 3 or
more alcoholic drinks every day while using this product

Drug Facts (continued)
■ take more or for a longer time than directed
Do not use ■ if you are allergic to aspirin or any other pain reliever/fever
reducer ■ if you have ever had an allergic reaction to this product or any of its
ingredients.
Ask a doctor before use if ■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn ■ you have
high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are
taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet
Ask a doctor or pharmacist before use if you are
■ presently taking a prescription drug. Antacids may interact with certain
prescription drugs. ■ taking a prescription drug for anticoagulation (thinning
the blood), diabetes, gout or arthritis
When using this product do not take more than 8 tablets (adults) or 4 tablets
(60 years and over) in a 24-hour period or use the maximum dosage for more
than 2 weeks, except under the advice and supervision of a physician.
Stop use and ask a doctor if ■ you experience any of the following signs of
stomach bleeding
■ feel faint
■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ an allergic reaction occurs. Seek medical help right away.
■ pain lasts for more than 10 days or gets worse
■ new symptoms occur
■ redness or swelling is present
■ ringing in the ears or loss of hearing occurs
If pregnant or breast-feeding, ask a health professional before use. It is
especially important not to use aspirin during the last 3 months of
pregnancy unless definitely directed to do so by a doctor because it may
cause problems in the unborn child or complications during delivery.



RITE AID ANTACID AND PAIN RELIEF
aspirin, citric acid, sodium bicarbonate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-4212
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1Z RA3Q20)	SODIUM BICARBONATE	1916 mg

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	L421
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-4212-4	12 in 1 CARTON	03/16/2020	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:11822-4212-7	36 in 1 CARTON	12/09/2022	
2		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	03/16/2020	

Labeler - Rite Aid (014578892)