RAPIDOL SODIUM BICARBONATE- sodium bicarbonate powder Pharmadel LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rapidol Sodium Bicarbonate Antacid Powder (WP)

Drug Facts

Active Ingredient & Purposes

Active ingredient (in each $\frac{1}{2}$ teaspoon)	Purpose
Sodium bicarbonate 2,616	Antacid
mg	Arreaciu

Uses For the relief of an upset stomach associated with these symptoms

- heatburn
- sour stomach
- acid indigestion

Warnings

Stomach Waring: To avoid serious injury, do not take powder until it is completely dissolved. It is very important not to take this product when overly full from food or drink. Consult a doctor if severe stomach pain occurs after taking this product.

Do not use

- the maximum daily dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor
- this product except under advice and supervision of a physician if you are allergic to milk or milk products

When using this product

- may cause constipation
- may have laxative effect

Ask a doctor or pharmacist before use if you are

• presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Ask a doctor before use if you have

• a sodium-restricted diet

If pregnant or breast-feeding,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

In case of an accidental overdose, seek professional assistance or contact a Poison Control center right away.

Directions

- do not take more than your "maximum daily dosage for your age group" in a 24-hour period. See Warnings.
- take a ½ teaspoon in a 4 fl. oz. (½ a glass) of water every 2 hours up to the **maximum daily dosage** or as directed by a doctor
- dissolve completely in water before drinking

age	maximum daily dosage
adults 60 years of age and over	$\frac{1}{2}$ teaspoon, 3 times a day
adults under 60 and children 12 years of age and over	$\frac{1}{2}$ teaspoon, 6 times a day
children under 12 years of age	do not use

Other information

- each ½ teaspoonful contains: sodium 716 mg
- store in cool dry place below 77°F
- do not use if seal under cap is torn, broken or missing

None

Questions & comments?

PHARMADEL LLC.

New Castle, DE, 19720

+1-866-359-3478

www.pharmadel.com

Prinicpal Display Panel

55758-327-04

4 Oz



RAPIDOL SODIUM BICARBONATE

sodium bicarbonate powder

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
	SODIUM BICARBONATE	2.6 g in 2.6 g

	 \sim 1		
Drod	Char	3CTA	ristics
	CHAL	acte	1 13 LICS

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Pa	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
		113 g in 1 BOTTLE; Type 0: Not a Combination Product	11/23/2020		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part331	11/23/2020		

Labeler - Pharmadel LLC (030129680)

Revised: 3/2023 Pharmadel LLC