

**GERICARE SORBITOL SOLUTION- sorbitol liquid**  
**GERI-CARE PHARMACEUTICALS, CORP**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Active Ingredient**

Sorbitol 70% USP

**Purpose**

□Laxative

□**Uses**

- relieves occasional constipation and irregularity
- generally produces bowel movement in 1/4 to 1 hour when used rectally
- as a pharmaceutical aide (sweetner)
- for other uses, ask your doctor

□**Warnings**

**For rectal use only**

**Do not use**

- □when abdominal pain, nausea, or vomiting are present
- for more than one week unless directed by a doctor

□**Ask a doctor before use if**

- are taking mineral oil
- have noticed a sudden change in bowel habits that lasts over 2 weeks

□**Stop use and ask a doctor if** □you have a rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.□

□**If pregnant or breast feeding** □ask a health professional before use.

□**Keep out of reach of children.** □In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions Do not exceed recommended dose**

---

AGE	DOSE
Adults and children 12 years and over	For rectal use only. Enema dosage is 120 mL of a 25 to 30% w/v solution (1 part of this product with 2.3 parts water) in a single daily dose as needed, or as directed by your doctor.
Children under 12 years of age	Ask a doctor

---

□**Inactive ingredients** □water

□**Questions or comments? 1-800-540-3765**

**Drug Facts**

<b>Active ingredient</b> Sorbitol 70%, USP	<b>Purpose</b> Laxative
---	----------------------------

**Uses**

- relieves occasional constipation and irregularity
- generally produces bowel movement in ¼ to 1 hour when used rectally
- as a pharmaceutical aide (sweetener)
- for other uses, ask your doctor

**Warnings**

**For rectal use only**

**Do not use** • when abdominal pain, nausea, or vomiting are present

• for more than one week unless directed by a doctor.

**Ask a doctor before use if you**

- are taking mineral oil
- have noticed a sudden change in bowel habits that lasts over 2 weeks

**Stop use and ask a doctor if you have rectal bleeding or no bowel movement after using this product.** These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. →

Code#: L-126 Rev. 12/20  
 Lot.#  
 Exp. Date:



**SORBITOL SOLUTION USP**  
**70% W/W**  
**LAXATIVE**

TAMPER EVIDENT: Do not use this product if inner foil seal over the mouth of bottle is missing or broken.

**16 FL OZ (1 PT) (473 mL)**

**Drug Facts (continued)**

**Directions • do not exceed recommended dose**

Age	Dose
Adults and children 12 years and over	For rectal use only. Enema dosage is 120 mL of a 25 to 30% w/w solution (1 part of this product with 2.3 parts water) in a single daily dose as needed, or as directed by your doctor.
Children under 12 years of age	Ask a doctor.

**Other information**

- store at room temperature 15°- 30° C (59°-86°F)
- below 59°F cloudiness and thickening may occur; warming will restore clarity and fluidity without affecting product quality
- do not freeze
- Pharmacist-Preserve and dispense in tight, light-resistant container with a child resistant cap as defined in the USP

**Inactive ingredient:** water

**Questions or comments? 1-800-540-3765**

3 57896 43516 5  
 DIST. BY: GERI-CARE PHARMACEUTICALS CORP.  
 1650 63rd Street  
 Brooklyn, NY 11204

**Drug Facts**

<b>Active ingredient</b> Sorbitol 70%, USP	<b>Purpose</b> Laxative
---	----------------------------

**Uses**

- relieves occasional constipation and irregularity
- generally produces bowel movement in ¼ to 1 hour when used rectally
- as a pharmaceutical aide (sweetener)
- for other uses, ask your doctor

**Warnings**

**For rectal use only**

**Do not use** • when abdominal pain, nausea, or vomiting are present

• for more than one week unless directed by a doctor.

**Ask a doctor before use if you**

- are taking mineral oil
- have noticed a sudden change in bowel habits that lasts over 2 weeks

**Stop use and ask a doctor if you have rectal bleeding or no bowel movement after using this product.** These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. →

Code#: L-126 Rev. 12/20  
 Lot.#  
 Exp. Date:



**SORBITOL SOLUTION USP**  
**70% W/W**  
**LAXATIVE**

TAMPER EVIDENT: Do not use this product if inner foil seal over the mouth of bottle is missing or broken.

**16 FL OZ (1 PT) (473 mL)**

**Drug Facts (continued)**

**Directions • do not exceed recommended dose**

Age	Dose
Adults and children 12 years and over	For rectal use only. Enema dosage is 120 mL of a 25 to 30% w/w solution (1 part of this product with 2.3 parts water) in a single daily dose as needed, or as directed by your doctor.
Children under 12 years of age	Ask a doctor.

**Other information**

- store at room temperature 15°- 30° C (59°-86°F)
- below 59°F cloudiness and thickening may occur; warming will restore clarity and fluidity without affecting product quality
- do not freeze
- Pharmacist-Preserve and dispense in tight, light-resistant container with a child resistant cap as defined in the USP

**Inactive ingredient:** water

**Questions or comments? 1-800-540-3765**

3 57896 43516 5  
 DIST. BY: GERI-CARE PHARMACEUTICALS CORP.  
 1650 63rd Street  
 Brooklyn, NY 11204

**GERICARE SORBITOL SOLUTION**

sorbitol liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:57896-758
<b>Route of Administration</b>	RECTAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
SORBITOL (UNII: 506T60A25R) (SORBITOL - UNII:506T60A25R)	SORBITOL	70 mg in 100 mL

**Inactive Ingredients**

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-758-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	01/01/2020	

**Labeler** - GERI-CARE PHARMACEUTICALS, CORP (611196254)

Revised: 12/2020

GERI-CARE PHARMACEUTICALS, CORP