ALLERGY AND CONGESTION RELIEF- loratadine, pseudoephedrine sulfate tablet, film coated, extended release L. Perrigo Company

Perrigo Allergy and Congestion Relief Drug Facts

Active ingredients (in each tablet)

Loratadine 5 mg

Pseudoephedrine sulfate 120 mg

Purpose

Antihistamine

Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- sneezing
- itchy, watery eyes
- runny nose
- itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

heart disease

- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- each tablet contains: calcium 25 mg
- do not use if blister unit is broken or torn
- store between 20° to 25°C (68° to 77°F)
- keep in a dry place

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Claritin-D[®] 12 hour Extended Release Tablets active ingredients allergy & congestion relief

Pseudoephedrine Sulfate 120 mg/Loratadine 5 mg

Extended Release Tablets

Nasal Decongestant/Antihistamine

12 Hour

INDOOR & OUTDOOR ALLERGIES

Relief of:

Nasal and Sinus Congestion Due to Colds or Allergies

Sneezing - Runny Nose - Itchy, Watery Eyes

Itchy Throat or Nose Due to Allergies

Non-Drowsy*

*When taken as directed. See Drug Facts Panel.

Actual Size

20 EXTENDED RELEASE TABLETS

allergy & congestion relief



Compare to Claritin-D® 12 hour **Extended Release Tablets** active ingredients

NDC 0113-2007-60

allergy & congestion relief

Pseudoephedrine Sulfate 120 mg/Loratadine 5 mg **Extended Release Tablets** Nasal Decongestant/Antihistamine

INDOOR & OUTDOOR ALLERGIES

Relief of:

Nasal and Sinus Congestion Due to Colds or Allergies

Sneezing • Runny Nose • Itchy, Watery Eyes
 Itchy Throat or Nose Due to Allergies

Non-Drowsy³

*When taken as directed. See Drug Facts Panel.

Actual Size



O EXTENDED RELEASE TABLETS

7U060 55 C2

Drug Facts

Active ingredients (in each tablet)

Purpose

Loratadine 5 mg......Antihistamine Pseudoephedrine sulfate 120 mg.......Mas al decongestant

■ temporarily relieves these symptoms due to hay feveror other upper respiratory allergies:

■ sneezing ■ itchy, watery eyes
■ runny nose ■ itching of the nose or throat

■temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

■ reduces swelling of n as al pas sages

■ temporarily relieves sinus congestion and pressure ■ temporarily restores freer breathing through the nose

Warnings

Do not u se

OPEN OTHER

END

■ if you have ever had an allergic reaction to this product or any of its ingredients

■ if you are now taking a prescription monoamine oxidase inhibi tor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkins on's disease), or for 2 weeks after stopping the MAOI drug. I fyou do not know if your prescription drug contains an MAOI, ask a doctor or pharmac is thefore taking this product.

Ask a doctor before use if you have

■ heart disease ■ thyroid disease ■ high blood pressure

■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ liveror kidney disease. Your doctors hould determine if you

need a different dose

Drug Facts (continued)

When using this product do not take more than directed. Taking more than directed may cause drowsines s

Stop use and ask a doctor if

■ an allergic reaction to this product occurs. Seek medical help right away.

■ symptoms do not improve within 7 days or are accompanied by

■ nervous ness, diz ziness or slæples sness occurs

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help orcontact a Poison Control Centerright away (1 -800-222 -1 222).

Directions ■ do not divid e, crush, chew or dissolve the tablet

1 tab letevery 1 2 hours; not more than 2 tablets in 2 4 hours adults and children 12 years and over children under 12 years of age ask a doctor consumers with liver or ask a doctor kidney disease

Ot her in formation

■ each tab let contains: calcium 25 mg ■ donot use if blister unit is broken or torn

■ store between 20° to 25° C (68° to 77 °F)

■ keep in a dry place

Inactive ingredients croscamellose sodium, dibasic calcium phosphate, hypromellose, lactos e monohydrate, magnesium stearate, pharmac eut cal ink, povidone, titanium dioxide

Drug Facts (continued)

Questions or comments? 1-8 00-7 19-9260

Distributed By



Allegan, MI 49010 perrigo.com



CONVENIENT RECLOSING TAB

ALLERGY AND CONGESTION RELIEF

loratadine, pseudoephedrine sulfate tablet, film coated, extended release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-2007
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	5 mg	
PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE SULFATE	120 mg	

Inactive Ingredients		
Ingredient Name	Strength	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)		
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics			
Color	WHITE (to off-white)	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	7U0
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0113-2007- 46	10 in 1 CARTON	04/17/2018	04/17/2018	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product			
2	NDC:0113-2007- 60	20 in 1 CARTON	04/12/2018		
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076050	04/12/2018	

Labeler - L. Perrigo Company (006013346)

Revised: 2/2024 L. Perrigo Company