

MEDIQUE AT HOME ASPIRIN- aspirin tablet, film coated
Unifirst First Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Medique @ Home Aspirin

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever lasts more than 3 days
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose

Adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours

Children under 12 years: consult a doctor

Other information

- store at room temperature
- avoid excessive heat and humidity
- **Tamper Evident** child resistant packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose*, microcrystalline cellulose*, polyethylene glycol* povidone*, propylene glycol*

* may contain

Questions or comments? 1-800-634-7680

Medique® @ Home

Aspirin Tablets

Aspirin 325 mg

Pain Reliever/Fever Reducer\ (NSAID)

Pull to Open

Child-Resistant Packets

*not actual size

50 Packets

(2 Tablets Per Packet)

Compare to the active ingredient in Genuine Bayer® Tablets



Aspirin Tablets

Aspirin 325 mg
Pain Reliever/Fever Reducer
(NSAID)



50 Packets (2 Tablets Per Packet) Compare to the active ingredient in Genuine Bayer® Tablets*

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Purpose Aspirin 325 mg (NSAID)..... Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug</p> <p>Uses temporarily relieves ■ headache ■ muscle pain ■ toothache ■ menstrual pain ■ pain and fever of colds ■ minor pain of arthritis</p> <p>Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock</p> <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcohol drinks every day while using this product ■ take more or for a longer time than directed</p> <p>Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer.</p>	<p>Drug Facts (continued)</p> <p>Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma</p> <p>Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis</p> <p>Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ if ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area ■ any new symptoms appear</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p style="text-align: right;"><i>(continued on back panel)</i></p>
---	---

Drug Facts (continued)	
Directions ■ drink a full glass of water with each dose	
Adults and children 12 years and over	take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.
Children under 12 years:	consult a doctor
Other information ■ store at room temperature ■ avoid excessive heat and humidity ■ TAMPER EVIDENT child resistant packets ■ do not use any opened or torn packets	
Inactive ingredients corn starch, hypromellose*, microcrystalline cellulose*, polyethylene glycol*, povidone*, propylene glycol* *may contain	
Questions or comments? 1-800-634-7680	

†Bayer® is a registered trademark of Bayer Consumer Health and is not affiliated with Medique Products or this product.

3 47682 70533 6

Reorder 70533
Manufactured for
Medique Products
Fort Myers, FL 33967
1-800-634-7680
www.mediqueproducts.com

Retain carton for complete product information

MEDIQUE AT HOME ASPIRIN

aspirin tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-533
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients	
Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STARCH, CORN (UNII: O8232NY3S)	

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;157;ASPIRIN
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-533-33	50 in 1 BOX	02/01/2021	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	02/01/2021	

Labeler - Unifirst First Aid Corporation (832947092)

Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(47682-533)

Revised: 2/2023

Unifirst First Aid Corporation