

IBUPROFEN- ibuprofen capsule, liquid filled
Camber Consumer Care. Inc.

Active ingredient

(in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

PURPOSE

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - o headache
 - o toothache
 - o backache
 - o menstrual cramps
 - o the common cold
 - o muscular aches
 - o minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - . feel faint
 - . vomit blood
 - . have bloody or black stools
 - . have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - . chest pain
 - . trouble breathing
 - . slurred speech
 - . leg swelling
 - . weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor

because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and over:	<ul style="list-style-type: none">• take 1 capsule every 4 to 6 hours while symptoms persist• if pain or fever does not respond to 1capsule, 2 capsules may be used• do not exceed 6 capsules in 24 hours, unless directed by a doctor
children under 12 years:	<ul style="list-style-type: none">• ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- Read all warnings and directions before use. Keep carton.
- Store at 20-25°C (68-77°F)
- Avoid excessive heat above 40°C (104°F)

Inactive ingredients

FD&C blue no. 1, FD&C yellow no. 6, gelatin, lecithin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan.

Questions or comments?

call 1-888-588-1418

NDC 69230-146-52

Ibuprofen Capsules, 200mg

Pain Reliever/ Fever Reducer

(NSAID)

200mg, 160 SOFTGELS

Liquid-Filled Capsules

CAMBER
NDC 69230-146-52

Ibuprofen Capsules, 200 mg

Solubilized Ibuprofen Capsules, 200 mg
Pain Reliever/Fever Reducer (NSAID)

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

WARNING: Do not take if sealant or bottle cap imprinted with "SEAL IT FOR YOUR PROTECTION" is broken or missing.

CAMBER
NDC 69230-146-52

Ibuprofen Capsules, 200 mg

Solubilized Ibuprofen Capsules, 200 mg
Pain Reliever/Fever Reducer (NSAID)

Compare to Advil® Liqui-Gel® active ingredient*

160 CAPSULES**

*Liqui-Gel capsules

Drug Facts

Active ingredient (in each capsule)	Purpose
Solubilized ibuprofen equal to 200 mg (ibuprofen (NSAID) (present as the free acid and potassium salt))	Pain reliever / Fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- biters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (Continued)

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - sturred speech
 - weakness in one part or side of body
 - leg swelling
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
- any new symptoms appear

Drug Facts (Continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used

adults and children 12 years and over:

- take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor

children under 12 years:

- ask a doctor

Other information

- each capsule contains potassium 20 mg
- Read all warnings and directions before use. Keep carton.
- Store at 20-25°C (68-77°F)
- Avoid excessive heat above 40°C (104°F)

Inactive ingredients: FD&C blue no. 1, FD&C yellow no. 6, gelatin (soybean), medium-chain triglycerides, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan.

Questions or comments? call 1-888-588-1418

Distributed by:
Camber Consumer Care, Inc.
Piscataway, NJ 08854, USA, Made in USA Rev 04/18

♦ This product is not manufactured or distributed by Pfizer Consumer Health care, owner of the registered trademark of Advil® Liqui-Gel®.



0 6 9 2 3 0 1 4 6 5 2 0

Exp. Lot

NO VARNISH

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69230-146
Route of Administration	ORAL, ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	GREEN	Score	no score
Shape	OVAL	Size	20 mm
Flavor		Imprint Code	AT146
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69230-146-20	1 in 1 CARTON	05/01/2018	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69230-146-40	1 in 1 CARTON	05/01/2018	
2		40 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:69230-146-80	1 in 1 CARTON	05/01/2018	
3		80 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:69230-146-52	1 in 1 CARTON	05/01/2018	
4		160 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
---------------------------	---	-----------------------------	---------------------------

ANDA	ANDA206999	05/01/2018	
------	------------	------------	--

Labeler - Camber Consumer Care, Inc. (079539968)

Revised: 4/2018

Camber Consumer Care, Inc.