

**WOMENS LAXATIVE- bisacodyl tablet, delayed release  
DOLGENCORP, LLC**

-----  
**Dollar General 44-607**

***Active ingredient (in each tablet)***

Bisacodyl USP, 5 mg

***Purpose***

Stimulant laxative

***Uses***

- for relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

***Warnings***

**Do not use**

if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness and cramps

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- take with a glass of water

---

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

---

### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

### ***Inactive ingredients***

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C red #27 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

### ***Questions or comments?***

**1-888-309-9030**

### ***Principal display panel***

**DG™ | health**

Compare to  
active ingredient  
of Dulcolax Pink®\*

### **Women's Laxative**

Bisacodyl USP, 5 mg  
Stimulant Laxative  
Gentle, dependable constipation relief

**5**  
mg each

**25 Tablets**


Comfort Coated  
Actual Tablet Size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by Sanofi-Aventis Deutschland GMBH, owner of the registered trademark Dulcolax Pink®.  
50844 ORG011960756

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

**100% Satisfaction Guaranteed!**  
(888) 309-9030



Actual Tablet Size

mg each

B-0315-607-56  
OR6011960756

50844 OR6011960756  
Dulcolax Pink®  
GMBH, owner of the registered trademark  
distributed by Sanofi-Aventis Deutschland  
This product is not manufactured or

**Drug Facts**

**Active ingredient (in each tablet)**  
Bisacodyl USP, 5 mg.....Stimulant laxative

**Uses**  
■ for relief of occasional constipation and irregularly  
■ this product generally produces bowel movement in 6 to 12 hours

**Warnings**  
Do not use if you cannot swallow without chewing.  
Ask a doctor before use if you have  
■ stomach pain, nausea or vomiting  
■ a sudden change in bowel habits that lasts more than 2 weeks  
When using this product  
■ do not chew or crush tablet(s)  
■ do not use within 1 hour after taking an antacid or milk  
■ you may have stomach discomfort, faintness and cramps  
Stop use and ask a doctor if  
■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.  
■ you need to use a laxative for more than 1 week  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact

1-888-309-9030  
**Questions or comments?**

**Inactive ingredients** acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C red #27 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, polydextrose, propylene glycol, propylparaben, sodium benzoate, sodium bicarbonate, stearyl shellac glaze, simethicone, sodium alginate, acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

**Other information** ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
■ avoid excessive humidity  
■ see end flap for expiration date and lot number

**Directions** ■ take with a glass of water  
adults and children take 1 to 3 tablets in a single daily dose  
children 6 to under 12 years and over 12 years take 1 tablet in a single daily dose  
children under 6 years ask a doctor

**Drug Facts (continued)**  
Poison Control Center right away.

**Drug Facts**

**Active ingredient (in each tablet)**  
Bisacodyl USP, 5 mg.....Stimulant laxative

**Uses**  
■ for relief of occasional constipation and irregularly  
■ this product generally produces bowel movement in 6 to 12 hours

**Warnings**  
Do not use if you cannot swallow without chewing.  
Ask a doctor before use if you have  
■ stomach pain, nausea or vomiting  
■ a sudden change in bowel habits that lasts more than 2 weeks  
When using this product  
■ do not chew or crush tablet(s)  
■ do not use within 1 hour after taking an antacid or milk  
■ you may have stomach discomfort, faintness and cramps  
Stop use and ask a doctor if  
■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.  
■ you need to use a laxative for more than 1 week  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact

0.78"

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No Print / No Varnish Lot no. & Exp. date

DISTRIBUTED BY OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

100%

Satisfaction Guaranteed!

(888) 309-9030

**Dollar General 44-607**

## WOMENS LAXATIVE

bisacodyl tablet, delayed release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-168
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

### Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
AMMONIA (UNII: 5138Q19F1X)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SHELLAC (UNII: 46N107B71O)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

### Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	5
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-168-56	1 in 1 CARTON	02/23/2023	
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	02/23/2023	

**Labeler** - DOLGENCORP, LLC (068331990)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55910-168)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-168) , pack(55910-168)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55910-168)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55910-168)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55910-168)

Revised: 2/2023

DOLGENCORP, LLC