


CEPHALEXIN - cephalexin powder, for suspension
STAT RX USA LLC

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Cephalexin for Oral Suspension, USP and other antibacterial drugs, Cephalexin for Oral Suspension, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION

Cephalexin, USP is a semisynthetic cephalosporin antibiotic intended for oral administration. It is 7-(D- α -Amino- α -phenylacetamido)-3-methyl-3-cephem-4-carboxylic acid monohydrate. Cephalexin has the molecular formula $C_{16}H_{17}N_3O_4S \cdot H_2O$ and the molecular weight is 365.41.

Cephalexin has the following structural formula:



The nucleus of cephalexin is related to that of other cephalosporin antibiotics. The compound is a zwitterion; i.e., the molecule contains both a basic and an acidic group. The isoelectric point of cephalexin in water is approximately 4.5 to 5.

The crystalline form of cephalexin which is available is a monohydrate. It is a white crystalline solid having a bitter taste. Solubility in water is low at room temperature; 1 or 2 mg/mL may be dissolved readily, but higher concentrations are obtained with increasing difficulty.

The cephalosporins differ from penicillins in the structure of the bicyclic ring system. Cephalexin has a *D*-phenylglycyl group as substituent at the 7-amino position and an unsubstituted methyl group at the 3-position.

Cephalexin for Oral Suspension, USP is a white to off white granular powder with characteristic flavor, and gives orange colored viscous suspension after reconstitution with water.

After mixing, each 5 mL of Cephalexin for Oral Suspension, USP, will contain cephalexin monohydrate equivalent to 125 mg (360 micro mol) or 250 mg (720 micro mol) of anhydrous cephalexin. The suspensions also contains the following inactive ingredients: colloidal silicon dioxide, FD and C Yellow No.6, tutti-frutti flavor, orange flavor, sodium lauryl sulphate, sucrose and xanthan gum.

CLINICAL PHARMACOLOGY

Human Pharmacology:

Cephalexin is acid stable and may be given without regard to meals. It is rapidly absorbed after oral administration. Following doses of 250 mg, 500 mg, and 1 g, average peak serum levels of approximately 9, 18, and 32 mcg/mL respectively were obtained at 1 hour. Measurable levels were

present 6 hours after administration. Cephalexin is excreted in the urine by glomerular filtration and tubular secretion. Studies showed that over 90% of the drug was excreted unchanged in the urine within 8 hours. During this period, peak urine concentrations following the 250-mg, 500-mg, and 1-g doses were approximately 1000, 2200, and 5000 mcg/mL respectively.

Microbiology:

In vitro tests demonstrate that the cephalosporins are bactericidal because of their inhibition of cell-wall synthesis. Cephalexin has been shown to be active against most strains of the following microorganisms both *in vitro* and in clinical infections as described in the **INDICATIONS AND USAGE** section.

Aerobes, Gram-positive:

Staphylococcus aureus (including penicillinase-producing strains)

Streptococcus pneumoniae (penicillin-susceptible strains)

Streptococcus pyogenes

Aerobes, Gram-negative:

Escherichia coli

Haemophilus influenzae

Klebsiella pneumoniae

Moraxella (Branhamella) catarrhalis

Proteus mirabilis

Note - Methicillin-resistant staphylococci and most strains of enterococci (*Enterococcus faecalis* [formerly *Streptococcus faecalis*]) are resistant to cephalosporins, including cephalexin. It is not active against most strains of *Enterobacter* spp., *Morganella morganii*, and *Proteus vulgaris*. It has no activity against *Pseudomonas* spp. or *Acinetobacter calcoaceticus*. Penicillin - resistant *Streptococcus pneumoniae* is usually cross-resistant to beta-lactam antibiotics.

Susceptibility Tests

Dilution Techniques:

Quantitative methods are used to determine antimicrobial minimal inhibitory concentrations (MIC's). These MIC's provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MIC's should be determined using a standardized procedure. Standardized procedures are based on a dilution method¹⁻³ (broth or agar) or equivalent with standardized inoculum concentrations and standardized concentrations of cephalothin powder. The MIC values should be interpreted according to the following criteria:

| MIC (mcg/mL) Interpretation | | |
|------------------------------------|--------------|-----|
| ≤8 | Susceptible | (S) |
| 16 | Intermediate | (I) |
| ≥32 | Resistant | (R) |

A report of "Susceptible" indicates that the pathogen is likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable. A report of "Intermediate" indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies

possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where high dosage of drug can be used. This category also provides a buffer zone which prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "Resistant" indicates that the pathogen is not likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable; other therapy should be selected.

Standardized susceptibility test procedures require the use of laboratory control microorganisms to control the technical aspects of the laboratory procedures. Standard cephalothin powder should provide the following MIC values:

| Microorganism | MIC (mcg/mL) |
|-----------------------------|---------------------|
| <i>E. coli</i> ATCC 25922 | 4-16 |
| <i>S. aureus</i> ATCC 29213 | 0.12-0.5 |

Diffusion Techniques:

Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure^{2,3} requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 30-mcg cephalothin to test the susceptibility of microorganisms to cephalexin.

Reports from the laboratory providing results of the standard single-disk susceptibility test with a 30-mcg cephalothin disk should be interpreted according to the following criteria:

| Zone Diameter (mm) | Interpretation |
|---------------------------|-----------------------|
| ≥18 | Susceptible (S) |
| 15-17 | Intermediate (I) |
| ≤14 | Resistant (R) |

Interpretation should be as stated above for results using dilution techniques. Interpretation involves correlation of the diameter obtained in the disk test with the MIC for cephalexin.

As with standard dilution techniques, diffusion methods require the use of laboratory control microorganisms that are used to control the technical aspects of the laboratory procedures. For the diffusion technique, the 30-mcg cephalothin disk should provide the following zone diameters in these laboratory test quality control strains:

| Microorganism | Zone Diameter (mm) |
|-----------------------------|---------------------------|
| <i>E. coli</i> ATCC 25922 | 15-21 |
| <i>S. aureus</i> ATCC 25923 | 29-37 |

INDICATIONS AND USAGE

Cephalexin for Oral Suspension, USP is indicated for the treatment of the following infections when caused by susceptible strains of the designated microorganisms:

Respiratory tract infections caused by *Streptococcus pneumoniae* and *Streptococcus pyogenes* (Penicillin is the usual drug of choice in the treatment and prevention of streptococcal infections, including the prophylaxis of rheumatic fever. Cephalexin for Oral Suspension, USP is generally effective in the eradication of streptococci from the nasopharynx; however, substantial data establishing the efficacy of cephalexin in the subsequent prevention of rheumatic fever are not available at present.)

Otitis media due to *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Staphylococcus aureus*,

Streptococcus pyogenes, and *Moraxella catarrhalis*

Skin and skin structure infections caused by *Staphylococcus aureus* and/or *Streptococcus pyogenes*

Bone infections caused by *Staphylococcus aureus* and/or *Proteus mirabilis*

Genitourinary tract infections, including acute prostatitis, caused by *Escherichia coli*, *Proteus mirabilis*, and *Klebsiella pneumoniae*

Note - Culture and susceptibility tests should be initiated prior to and during therapy. Renal function studies should be performed when indicated.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Cephalexin for Oral Suspension, USP and other antibacterial drugs, Cephalexin for Oral Suspension, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS

Cephalexin for Oral Suspension, USP is contraindicated in patients with known allergy to the cephalosporin group of antibiotics.

WARNINGS

BEFORE THERAPY WITH CEPHALEXIN IS INSTITUTED, CAREFUL INQUIRY SHOULD BE MADE TO DETERMINE WHETHER THE PATIENT HAS HAD PREVIOUS HYPERSENSITIVITY REACTIONS TO CEPHALEXIN, CEPHALOSPORINS, PENICILLINS, OR OTHER DRUGS. IF THIS PRODUCT IS TO BE GIVEN TO PENICILLIN-SENSITIVE PATIENTS, CAUTION SHOULD BE EXERCISED BECAUSE CROSS-HYPERSENSITIVITY AMONG BETA-LACTAM ANTIBIOTICS HAS BEEN CLEARLY DOCUMENTED AND MAY OCCUR IN UP TO 10% OF PATIENTS WITH A HISTORY OF PENICILLIN ALLERGY. IF AN ALLERGIC REACTION TO CEPHALEXIN OCCURS, DISCONTINUE THE DRUG. SERIOUS ACUTE HYPERSENSITIVITY REACTIONS MAY REQUIRE TREATMENT WITH EPINEPHRINE AND OTHER EMERGENCY MEASURES, INCLUDING OXYGEN, INTRAVENOUS FLUIDS. INTRAVENOUS ANTIHISTAMINES, CORTICOSTEROIDS, PRESSOR AMINES AND AIRWAY MANAGEMENT, AS CLINICALLY INDICATED.

There is some clinical and laboratory evidence of partial cross-allergenicity of the penicillins and the cephalosporins. Patients have been reported to have had severe reactions (including anaphylaxis) to both drugs.

Any patient who has demonstrated some form of allergy, particularly to drugs, should receive antibiotics cautiously. No exception should be made with regard to cephalexin.

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including cephalexin, and may range from mild to life threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is a primary cause of "antibiotic-associated" colitis.

After the diagnosis of pseudomembranous colitis has been established, therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes,

protein supplementation, and treatment with an antibacterial drug clinically effective against *Clostridium difficile* colitis.

PRECAUTIONS

General:

Prescribing Cephalexin for Oral Suspension, USP in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Patients should be followed carefully so that any side effects or unusual manifestations of drug idiosyncrasy may be detected. If an allergic reaction to cephalexin occurs, the drug should be discontinued and the patient treated with the usual agents (e.g., epinephrine or other pressor amines, antihistamines, or corticosteroids).

Prolonged use of cephalexin may result in the overgrowth of nonsusceptible organisms. Careful observation of the patient is essential. If superinfection occurs during therapy, appropriate measures should be taken.

Positive direct Coombs' tests have been reported during treatment with the cephalosporin antibiotics. In hematologic studies or in transfusion cross-matching procedures when antiglobulin tests are performed on the minor side or in Coombs' testing of newborns whose mothers have received cephalosporin antibiotics before parturition, it should be recognized that a positive Coombs' test may be due to the drug.

Cephalexin should be administered with caution in the presence of markedly impaired renal function. Under such conditions, careful clinical observation and laboratory studies should be made because safe dosage may be lower than that usually recommended.

Indicated surgical procedures should be performed in conjunction with antibiotic therapy.

Broad-spectrum antibiotics should be prescribed with caution in individuals with a history of gastrointestinal disease, particularly colitis.

Cephalosporins may be associated with a fall in prothrombin activity. Those at risk include patients with renal or hepatic impairment, or poor nutritional state, as well as patients receiving a protracted course of antimicrobial therapy, and patients previously stabilized on anticoagulant therapy. Prothrombin time should be monitored in patients at risk and exogenous vitamin K administered as indicated.

Information for Patients :

Patients should be counseled that antibacterial drugs including Cephalexin for Oral Suspension, USP should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Cephalexin for Oral Suspension, USP is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Cephalexin for Oral Suspension, USP or other antibacterial drugs in the future.

Drug Interactions:

Metformin: In healthy subjects given single 500 mg doses of cephalexin and metformin, plasma metformin mean C_{max} and AUC increased by an average of 34% and 24%, respectively, and metformin mean renal clearance decreased by 14%. No information is available about the interaction of cephalexin and metformin following multiple doses of either drug.

Although not observed in this study, adverse effects could potentially arise from co-administration of

cephalexin and metformin by inhibition of tubular secretion via organic cationic transporter systems. Accordingly, careful patient monitoring and dose adjustment of metformin is recommended in patients concomitantly taking cephalexin and metformin.

Probenecid: As with other β -lactams, the renal excretion of cephalexin is inhibited by probenecid.

Drug/Laboratory Test Interactions:

As a result of administration of cephalexin, a false-positive reaction for glucose in the urine may occur. This has been observed with Benedict's and Fehling's solutions and also with Clinitest[®] tablets.

Carcinogenesis, Mutagenesis, Impairment of Fertility:

Lifetime studies in animals have not been performed to evaluate the carcinogenic potential of cephalexin. Tests to determine the mutagenic potential of cephalexin have not been performed. In male and female rats, fertility and reproductive performance were not affected by cephalexin oral doses up to 1.5 times the highest recommended human dose based upon mg/m².

Pregnancy:

Teratogenic effects -Pregnancy Category B- Reproduction studies have been performed on mice and rats using oral doses of cephalexin monohydrate 0.6 and 1.5 times the maximum daily human dose (66 mg/kg/day) based upon mg/m², and have revealed no harm to the fetus. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers :

The excretion of cephalexin in human milk increased up to 4 hours after a 500-mg dose; the drug reached a maximum level of 4 mcg/mL, then decreased gradually, and had disappeared 8 hours after administration. Caution should be exercised when cephalexin is administered to a nursing woman.

Pediatric Use:

The safety and effectiveness of cephalexin in pediatric patients was established in clinical trials for the dosages described in the DOSAGE AND ADMINISTRATION section. In these trials, pediatric patients may have received cephalexin capsules or cephalexin for Oral Suspension.

Geriatric Use:

Of the 701 subjects in 3 published clinical studies of cephalexin, 433 (62%) were 65 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function (see **PRECAUTIONS**, General).

ADVERSE REACTIONS

Gastrointestinal:

Onset of pseudomembranous colitis may occur during or after antibacterial treatment. (See **WARNINGS**.) Nausea and vomiting have been reported rarely. The most frequent side effect has been diarrhea. It was very rarely severe enough to warrant cessation of therapy. Dyspepsia, gastritis, and abdominal pain have also occurred. As with some penicillins and some other cephalosporins, transient

hepatitis and cholestatic jaundice have been reported rarely.

Hypersensitivity:

Allergic reactions in the form of rash, urticaria, angioedema, and, rarely, erythema multiforme, Stevens-Johnson syndrome, or toxic epidermal necrolysis have been observed. These reactions usually subsided upon discontinuation of the drug. In some of these reactions, supportive therapy may be necessary. Anaphylaxis has also been reported.

Other reactions have included genital and anal pruritus, genital moniliasis, vaginitis and vaginal discharge, dizziness, fatigue, headache, agitation, confusion, hallucinations, arthralgia, arthritis, and joint disorder. Reversible interstitial nephritis has been reported rarely. Eosinophilia, neutropenia, thrombocytopenia, hemolytic anemia, and slight elevations in AST and ALT have been reported.

In addition to the adverse reactions listed above that have been observed in patients treated with cephalexin, the following adverse reactions and altered laboratory tests have been reported for cephalosporin class antibiotics:

Adverse Reactions:

Fever, colitis, aplastic anemia, hemorrhage, renal dysfunction and toxic nephropathy.

Several cephalosporins have been implicated in triggering seizures, particularly in patients with renal impairment when the dosage was not reduced (See **INDICATIONS AND USAGE** and **PRECAUTIONS, General**). If seizures associated with drug therapy should occur, the drug should be discontinued. Anticonvulsant therapy can be given if clinically indicated.

Altered Laboratory Tests:

Prolonged prothrombin time, increased BUN, increased creatinine, elevated alkaline phosphatase, elevated bilirubin, elevated LDH, pancytopenia, leukopenia, and agranulocytosis.

OVERDOSAGE

Signs and Symptoms:

Symptoms of oral overdose may include nausea, vomiting, epigastric distress, diarrhea, and hematuria. If other symptoms are present, it is probably secondary to an underlying disease state, an allergic reaction, or toxicity due to ingestion of a second medication.

Treatment:

To obtain up-to-date information about the treatment of overdose, a good resource is your certified Regional Poison Control Center. Telephone numbers of certified poison control centers are listed in the *Physicians' Desk Reference (PDR)*. In managing overdosage, consider the possibility of multiple drug overdoses, interaction among drugs, and unusual drug kinetics in your patient.

Unless 5 to 10 times the normal dose of cephalexin has been ingested, gastrointestinal decontamination should not be necessary.

Protect the patient's airway and support ventilation and perfusion. Meticulously monitor and maintain, within acceptable limits, the patient's vital signs, blood gases, serum electrolytes, etc. Absorption of drugs from the gastrointestinal tract may be decreased by giving activated charcoal, which, in many cases, is more effective than emesis or lavage; consider charcoal instead of or in addition to gastric emptying. Repeated doses of charcoal over time may hasten elimination of some drugs that have been absorbed. Safeguard the patient's airway when employing gastric emptying or charcoal.

Forced diuresis, peritoneal dialysis, hemodialysis, or charcoal hemoperfusion have not been

established as beneficial for an overdose of cephalexin; however, it would be extremely unlikely that one of these procedures would be indicated.

The oral median lethal dose of cephalexin in rats is > 5000 mg/kg.

DOSAGE AND ADMINISTRATION

Cephalexin for Oral Suspension, USP is administered orally.

Adults:

The adult dosage ranges from 1 to 4 g daily in divided doses. The 333 mg and 750 mg strengths should be administered such that the daily dose is within 1 to 4 grams per day. The usual adult dose is 250 mg every 6 hours. For the following infections, a dosage of 500 mg may be administered every 12 hours: streptococcal pharyngitis, skin and skin structure infections, and uncomplicated cystitis in patients over 15 years of age. Cystitis therapy should be continued for 7 to 14 days. For more severe infections or those caused by less susceptible organisms, larger doses may be needed. If daily doses of Cephalexin for Oral Suspension, USP greater than 4 g are required, parenteral cephalosporins, in appropriate doses, should be considered.

Pediatric Patients :

The usual recommended daily dosage for pediatric patients is 25 to 50 mg/kg in divided doses. For streptococcal pharyngitis in patients over 1 year of age and for skin and skin structure infections, the total daily dose may be divided and administered every 12 hours.

Cephalexin Suspension

| | |
|---------------|-----------------------|
| <u>Weight</u> | <u>125 mg/5 mL</u> |
| 10 kg (22 lb) | 1/2 to 1 tsp q.i.d. |
| 20 kg (44 lb) | 1 to 2 tsp q.i.d. |
| 40 kg (88 lb) | 2 to 4 tsp q.i.d. |
| <u>Weight</u> | <u>250 mg/5 mL</u> |
| 10 kg (22 lb) | 1/4 to 1/2 tsp q.i.d. |
| 20 kg (44 lb) | 1/2 to 1 tsp q.i.d. |
| 40 kg (88 lb) | 1 to 2 tsp q.i.d. |
| or | |
| <u>Weight</u> | <u>125 mg/5 mL</u> |
| 10 kg (22 lb) | 1 to 2 tsp b.i.d. |
| 20 kg (44 lb) | 2 to 4 tsp b.i.d. |
| 40 kg (88 lb) | 4 to 8 tsp b.i.d. |
| <u>Weight</u> | <u>250 mg/5 mL</u> |
| 10 kg (22 lb) | 1/2 to 1 tsp b.i.d. |
| 20 kg (44 lb) | 1 to 2 tsp b.i.d. |
| 40 kg (88 lb) | 2 to 4 tsp b.i.d. |

In severe infections, the dosage may be doubled.

In the therapy of otitis media, clinical studies have shown that a dosage of 75 to 100 mg/kg/day in 4 divided doses is required.

In the treatment of β -hemolytic streptococcal infections, a therapeutic dosage of Cephalexin for Oral Suspension, USP should be administered for at least 10 days.

Direction for Mixing:

| Cephalexin for Oral Suspension, USP | Bottle Size | Directions for Reconstitution |
|-------------------------------------|-------------|---|
| 125 mg/5 mL | 100 mL | Add 68 mL of water in two portions to the dry mixture in the bottle. Shake well after each addition. |
| | 200 mL | Add 136 mL of water in two portions to the dry mixture in the bottle. Shake well after each addition. |
| 250 mg/5 mL | 100 mL | Add 68 mL of water in two portions to the dry mixture in the bottle. Shake well after each addition. |
| | 200 mL | Add 136 mL of water in two portions to the dry mixture in the bottle. Shake well after each addition. |

HOW SUPPLIED

Cephalexin for Oral Suspension, USP is a white to off white granular powder with characteristic flavor, and gives orange colored viscous suspension after reconstitution with water.

Cephalexin for Oral Suspension, USP, is available in:

The 125 mg/5 mL oral suspension is available as follows

100 - mL Bottles NDC 42043-142-38

200 - mL Bottles NDC 42043-142-58

The 250 mg/5 mL oral suspension is available as follows

100 - mL Bottles NDC 42043-143-38

200 - mL Bottles NDC 42043-143-58

After mixing store in a refrigerator. May be kept for 14 days without significant loss of potency. Shake well before using. Keep tightly closed.

Prior to mixing, store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

REFERENCES

1. National Committee for Clinical Laboratory Standards: Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically-Fourth Edition. Approved Standard NCCLS Document M7-A4, Vol. 17, No. 2, NCCLS, Wayne, PA, January, 1997.
2. National Committee for Clinical Laboratory Standards. Performance Standards for Antimicrobial Disk Susceptibility Tests- Sixth Edition. Approved Standard NCCLS Document M2-A6, Vol. 17, No. 1, NCCLS, Wayne, PA, January, 1997.
3. National Committee for Clinical Laboratory Standards. Performance Standards for Antimicrobial Susceptibility Testing- Eighth Informational Supplement. Approved Standard NCCLS Document M100-S8, Vol. 18, No. 1, NCCLS, Wayne, PA, January, 1998.

Manufactured for:

Orchid Healthcare

(A Division of Orchid Chemicals and Pharmaceuticals Ltd.)

At Irungattukottai - 602 105, India

Distributed by:



Karalex Pharma, LLC

470 Chestnut Ridge Road, Woodcliff Lake, NJ 07677

Revised 03/10

948025975

PACKAGE LABEL-CEPHALEXIN ORAL SUSP 125 mg/ 5mL, 100mL

Packaged and distributed by:  STAT R USA, LLC Gainesville, GA 30501

Cephalexin Oral Susp
125mg/5mL 100 mL

Generic For: **Keflex**

NDC 16590-466-32 Prod# 466-32
Lot# SAMPLE

When reconstituted each (5mL) contains: Cephalexin equivalent to 125mg anhydrous cephalexin

Mfg By: Karalex Pharma, LLC
Woodcliff Lake, NJ 07677 NDC 42043-142-38

Mfg Lot: HL06708
Discard After: 04/11 MD 4/21/2009 SAMPLE

RX ONLY-KEEP OUT OF REACH OF CHILDREN

IMPORTANT: Finish all this medication unless otherwise directed by prescriber. § Shake well before using.

BA53Y
Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed.

S 0



PACKAGE LABEL-CEPHALEXIN ORAL SUSP 250 mg/ 5mL, 200mL

Packaged and distributed by: **STAT R_x USA** Gainesville, GA 30501

Cephalexin Oral Susp
250mg/5mL **200 mL**

Generic For: **Keflex**

NDC 16590-049-36 Prod# 049-36
 Lot# SAMPLE

When reconstituted each 5mL contains: Cephalexin equivalent to 250mg of anhydrous cephalexin

Mfg By: Karalex Pharma, LLC
 Woodcliff Lake, NJ 07677 NDC 42043-143-58

Mfg Lot: C002852 MD 8/4/2010 SAMPLE
 Discard After: 12/12

RX ONLY-KEEP OUT OF REACH OF CHILDREN

AS JMF
 Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed.

IMPORTANT: Finish all this medication unless otherwise directed by prescriber. § SHAKE WELL



CEPHALEXIN

cephalexin powder, for suspension

Product Information

| | | | |
|-------------------------|-------------------------|--------------------|------------------------------|
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:16590-466(NDC:42043-142) |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------------|
| CEPHALEXIN (UNII: OBN7UDS42Y) (CEPHALEXIN ANHYDROUS - UNII:5SFF1W6677) | CEPHALEXIN | 125 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J) | |
| SUCROSE (UNII: C151H8M554) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |

Product Characteristics

| | | | |
|----------|----------------------|--------------|--|
| Color | | Score | |
| Shape | | Size | |
| Flavor | TUTTI FRUTTI, ORANGE | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:16590-466-32 | 100 mL in 1 BOTTLE | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA065326 | 12/07/2007 | |

CEPHALEXIN

cephalexin powder, for suspension

Product Information

| | | | |
|-------------------------|-------------------------|--------------------|------------------------------|
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:16590-049(NDC:42043-143) |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------------|
| CEPHALEXIN (UNII: OBN7UDS42Y) (CEPHALEXIN ANHYDROUS - UNII:5SFF1W6677) | CEPHALEXIN | 250 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| SODIUM LAURYL SULFATE (UNII: 368GB5141J) | |
| SUCROSE (UNII: C151H8M554) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |

Product Characteristics

| | | | |
|----------|----------------------|--------------|--|
| Color | | Score | |
| Shape | | Size | |
| Flavor | TUTTI FRUTTI, ORANGE | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---------------------|----------------------|--------------------|
| 1 | NDC:16590-049-36 | 200 mL in 1 BOTTLE | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA065326 | 12/04/2007 | |

Labeler - STAT RX USA LLC (786036330)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------|---------|-----------|---------------------|
| STAT RX USA LLC | | 786036330 | relabel, repack |

Revised: 5/2011

STAT RX USA LLC