DOCUSATE SODIUM - STOOL SOFTENER LAXATIVE- docusate sodium capsule, liquid filled Cardinal Health 107, LLC

Cardinal Health

Docusate Sodium - Stool Softener Laxative

Drug Facts

Active ingredient (in each softgel)

Docusate Sodium 100 mg

Purpose

Stool softener laxative

Uses

- for relief of occasional constipation (irregularity)
- This generally produces a bowel movement within 12 to 72 hours.

Warnings

Ask a doctor before use if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are presently taking mineral oil

Stop use and ask a doctor if

• you have rectal bleeding or no bowel movement after using this product.

These can be signs of a serious condition

• you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health care professional before use.

This unit dose package is not child resistant and is Intended for Institutional Use Only.

Keep out of reach of children

Overdose Warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

• take softgel with a full glass (8 oz) of water

·

adults and children 12 years of age and older	take 1 to 3 softgels daily or as directed by a doctor. This dose may be taken as a single daily dose or in divided doses.
children 6 to under 12 years of age	take 1 softgel daily or as directed by a doctor
children under 6 years of age	ask a doctor

Other information

- each softgel contains: sodium 10 mg (very low sodium)
- store at 25°C (77°F) in a dry place. Avoid excessive heat 40°C (140°F).

Inactive ingredients

edible white ink, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, isopropyl alcohol, light mineral oil, medium chain triglycerides, polyethylene glycol, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments?

(800)616-2471

Distributed by:

MAJOR® PHARMACEUTICALS

Livonia, MI 48152

Distributed By:

Cardinal Health

Dublin, OH 43017

L57466800124

Product of UAE

Packaged and Quality Assured in the USA

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

READ FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Principal Display Panel

Docusate Sodium



R122

DOCUSATE SODIUM 100 mg Stool Softener Laxative

10 SOFTGELS

Each softgel contains: sodium 10 mg (very low sodium)

Drug Facts

Active ingredient (in each softgel)

Docusate Sodium 100 mg

Purpose

Stool softener laxative

Inactive ingredients edible white ink, FD&C red no. 40, FD&C yellow no. 6, gelatin, glycerin, isopropyl alcohol, light mineral oil, medium chain triglycerides, polyethylene glycol, propylene glycol, purified water, sorbitol sorbitan solution

Uses -for relief of occasional constipation (irregularity)

-This generally produces a bowel movement within 12 to 72 hours

Directions - take softgels with a full glass (8 oz) of water

adults and children 12 years of age and older take 1 to 3 softgels daily or as directed by a doctor. This dose may be taken as a single daily dose or in divided doses.

children 6 to under 12 years of age

take 1 softgel daily or as directed by a doctor

children under 6 years of age ask a doctor

STORAGE: Store at 25° C (77° F); in a dry place.

Avoid excessive heat 40° C (140° F).

Warnings

Ask a doctor before use if you have -abdominal pain, nausea or vomiting -a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are presently taking mineral oil Stop use and ask a doctor if -you have rectal bleeding or no bowel movement after using this product .These can be signs of a serious condition.

-you need to use a laxative for more than 1 week

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. If pregnant or breast-feeding, ask a health care professional before use. Keep out of reach of children.

Overdose Warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical

attention is critical for adults as well as children even if you do not notice any signs or symptoms.

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING READ FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Questions or comments? (800) 616-2471

Distributed by MAJOR® PHARMACEUTICALS

Indianapolis, IN 46268

Product of UAE

Packaged and Quality Assured in the USA

Distributed by Cardinal Health

Dublin, OH 43017

L57466800124

Exp:

DOCUSATE SODIUM - STOOL SOFTENER LAXATIVE

docusate sodium capsule, liquid filled

Product Information

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:55154-3369(NDC:0904-7183)Route of AdministrationORAL

Active Ingredient/Active Moiety

	Ingredient Name	Basis of Strength	Strength
l	DOCUSATE SODIUM (UNII: F05Q2T2JA0) (DOCUSATE - UNII:M7P27195AG)	DOCUSATE SODIUM	100 mg

Inactive Ingredients		
Ingredient Name	Strength	
FD&C RED NO. 40 (UNII: WZB9127XOA)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)		
GLYCERIN (UNII: PDC6A3C0OX)		
ISOPROPYL ALCOHOL (UNII: ND2M416302)		
LIGHT MINERAL OIL (UNII: N6K5787QVP)		
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3MJQ0SDW1A)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SORBITOL (UNII: 506T60A25R)		
SORBITAN (UNII: 6092ICV9RU)		

Product Characteristics				
Color	red	Score	no score	
Shape	OVAL	Size	12mm	
Flavor		Imprint Code	777	
Contains				

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154- 3369-0	10 in 1 BAG	09/04/2021	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	09/04/2021	

Revised: 1/2024 Cardinal Health 107, LLC