

**PREVAGE CITY SMART- titanium dioxide, zinc oxide emulsion  
REVLON**

-----  
**ELIZABETH ARDEN PREVAGE CITY SMART BROAD SPECTRUM HYDRATING  
SHIELD**

**ACTIVE INGREDIENTS**

TITANIUM DIOXIDE 6.2% W/W .....Sunscreen  
ZINC OXIDE 5.6% W/W .....Sunscreen

**INACTIVE INGREDIENTS**

CYCLOPENTASILOXANE, WATER/AQUA/EAU, ISODODECANE, DIMETHICONE, C12-15 ALKYL BENZOATE, ISOPROPYL MYRISTATE, PENTYLENE GLYCOL, BUTYLENE GLYCOL, SILICA DIMETHYL SILYLATE, CAPRYLYL METHICONE, POLYGLYCERYL-3 POLYDIMETHYLSILOXYETHYL DIMETHICONE, ACRYLATES/POLYTRIMETHYLSILOXYMETHACRYLATE COPOLYMER, GLYCERIN, ALUMINUM HYDROXIDE, STEARIC ACID, PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, CARNOSINE, ACRYLATES/ETHYLHEXYL ACRYLATE/DIMETHICONE METHACRYLATE COPOLYMER, ARABIDOPSIS THALIANA EXTRACT, CAMELLIA SINENSIS LEAF EXTRACT, CAPRYLYL GLYCOL, DISODIUM EDTA, ERGOTHIONEINE, FERULIC ACID, HYDROGEN DIMETHICONE, HYDROXYDECYL UBIQUINOYL DIPALMITOYL GLYCERATE, LECITHIN, MICA, MICROCOCCUS LYSATE, PARFUM/FRAGRANCE, PEG/PPG-18/18 DIMETHICONE, PLANKTON EXTRACT, SODIUM PCA, TIN OXIDE, TOCOPHERYL ACETATE, TRIETHOXYCAPRYLYLSILANE, ALPHA-ISOMETHYL IONONE, LINALOOL, CHLORPHENESIN, PHENOXYETHANOL, IRON OXIDES (CI 77491, CI 77492, CI 77499), TITANIUM DIOXIDE (CI 77891).

**USES**

- Helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

**WARNINGS**

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

**DIRECTIONS FOR SUNSCREEN USE**

### Directions For Sunscreen Use:

■ Apply liberally 15 minutes before sun exposure

■ Reapply:

- at least every 2 hours
- use a water resistant sunscreen if swimming or sweating

■ Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

■ Limit time in the sun, especially from 10 a.m. – 2 p.m.

■ Wear long-sleeve shirts, pants, hats, and sunglasses

■ Children under 6 months: Ask a doctor

### **OTHER INFORMATION**

■ Protect this product from excessive heat and direct

■ Store at room temperature

As needed

- Helps prevent sunburn

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Sunscreen

### **CARTON ART**

## Drug Facts

Active Ingredient	Purpose
TITANIUM DIOXIDE 6.2% W/W	Sunscreen
ZINC OXIDE 5.6% W/W	Sunscreen

### Uses

- Helps prevent sunburn
- If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

### Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes.  
Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

### Directions For Sunscreen Use:

- Apply liberally 15 minutes before sun exposure
- Reapply:
  - at least every 2 hours
  - use a water resistant sunscreen if swimming or sweating
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
  - Limit time in the sun, especially from 10 a.m. – 2 p.m.
  - Wear long-sleeve shirts, pants, hats, and sunglasses
  - Children under 6 months: Ask a doctor

### Inactive Ingredients

CYCLOPENTASILOXANE, WATER/AQUA/AEAU, ISODODECANE, DIMETHICONE, C12-15 ALKYL BENZOATE, ISOPROPYL MYRISTATE, PENTYLENE GLYCOL, BUTYLENE GLYCOL, SILICA DIMETHYL SILYLATE, CAPRYLYL METHICONE, POLYGLYCERYL-3 POLYDIMETHYLSILOXYETHYL DIMETHICONE, ACRYLATES/POLYTRIMETHYLSILOXYMETHACRYLATE COPOLYMER, GLYCERIN, ALUMINUM HYDROXIDE, STEARIC ACID, PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE, CARNOSINE, ACRYLATES/ETHYLHEXYL ACRYLATE/ETHYLHEXYL METHACRYLATE COPOLYMER, ARABIDOPSIS THALIANA EXTRACT, CAMELLIA SINENSIS LEAF EXTRACT, CAPRYLYL GLYCOL, DISODIUM EDTA, ERGOTHIONEINE, FERULIC ACID, HYDROGEN DIMETHICONE, HYDROXYDECYL UBIQUINOYL DIPALMITOYL GLYCERATE, LECITHIN, MICA, MICROCOCCUS LYSATE, PERFUM/FRAGRANCE, PEG/PPG-18/18 DIMETHICONE, PLANKTON EXTRACT, SODIUM PCA, TIN OXIDE, TOCOPHERYL ACETATE, TRIETHOXYCAPRYLYLSILANE, ALPHA-ISOMETHYL IONONE, LINALOOL, CHLORPHENESIN, PHENOXYETHANOL, IRON OXIDES (CI 77491, CI 77492, CI 77499), TITANIUM DIOXIDE (CI 77891).

### Other information

- Protect this product from excessive heat and direct
- Store at room temperature

This sheer, oil free, multi active skin defender delivers advanced protection against today's environmental aggressors including pollution, UV rays and free radicals to keep skin looking younger and brighter. An Anti-Pollution Complex and Broad Spectrum SPF 50 physical sunscreen form an invisible protective shield to screen out pollutants and 98% of harmful UVB rays. Powerful antioxidants fight free radicals while a DNA Enzyme Complex™ supports skin's natural repair system. Essential daily care to keep skin hydrated, protected and youthful, now and over time.

Clinically, dermatologist and allergy tested. Paraben Free. Non-Comedogenic. Safe for sensitive skin. Universal Tint. For all skin types.

To use: Apply evenly to face and neck as the last step in your skincare regimen. See insert or [elizabetharden.com](http://elizabetharden.com) for complete directions prior to first use.



prevage®

### city smart

- + DNA enzyme complex™
  - + anti-pollution
  - + antioxidants
- broad spectrum SPF 50  
hydrating shield

### Drug Facts (continued)

### Questions or Comments?

Call toll free 1-844-273-3679

1.34 FL.OZ. 40 ml

©Elizabeth Arden, Inc. Dist.  
New York, NY 10003

## PREVAGE CITY SMART

titanium dioxide, zinc oxide emulsion

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:10967-670
Route of Administration	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	5.6 mg in 1 g
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	6.2 mg in 1 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>ARABIDOPSIS THALIANA WHOLE</b> (UNII: AI3L60HQ81)	
<b>PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE</b> (UNII: TYP81E471F)	
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	
<b>LINALOOL, (+/-)-</b> (UNII: D81QY6I88E)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>ALKYL (C12-15) BENZOATE</b> (UNII: A9EJ3J61HQ)	
<b>CAPRYLYL TRISILOXANE</b> (UNII: Q95M2P1KJL)	
<b>ISOMETHYL-.ALPHA.-IONONE</b> (UNII: 9XP4LC555B)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>MICA</b> (UNII: V8A1AW0880)	
<b>PEG/PPG-18/18 DIMETHICONE</b> (UNII: 9H0AO7T794)	
<b>SODIUM PYRROLIDONE CARBOXYLATE</b> (UNII: 469OTG57A2)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>HYDROGEN DIMETHICONE (20 CST)</b> (UNII: 12Z59IF64N)	
<b>HYDROXYDECYL UBIQUINOYL DIPALMITOYL GLYCERATE</b> (UNII: OV1BT2N8RC)	
<b>CYCLOMETHICONE 5</b> (UNII: 0THT5PCI0R)	
<b>PENTYLENE GLYCOL</b> (UNII: 50C1307PZG)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>POLYGLYCERYL-3 POLYDIMETHYLSILOXYETHYL DIMETHICONE (4000 MPA.S)</b> (UNII: RLA2U05Z4Q)	
<b>GREEN TEA LEAF</b> (UNII: W2ZU1RY8B0)	
<b>CAPRYLYL GLYCOL</b> (UNII: 00YIU5438U)	
<b>EDETATE DISODIUM ANHYDROUS</b> (UNII: 8NLQ36F6MM)	
<b>ERGOTHIONEINE</b> (UNII: BDZ3DQM98W)	
<b>CARNOSINE</b> (UNII: 8HO6PVN24W)	
<b>ISOPROPYL MYRISTATE</b> (UNII: 0RE8K4LNJS)	
<b>SILICA DIMETHYL SILYLATE</b> (UNII: EU2PSP0G0W)	
<b>FERULIC ACID</b> (UNII: AVM951ZWT)	
<b>STANNIC OXIDE</b> (UNII: KM7N50LOS6)	
<b>.ALPHA.-TOCOPHEROL ACETATE</b> (UNII: 9E8X80D2L0)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)	
<b>CHLORPHENESIN</b> (UNII: I670DAL4SZ)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>ISODODECANE</b> (UNII: A8289P68Y2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10967-670-62	40 g in 1 TUBE; Type 0: Not a Combination Product	03/14/2019	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	03/14/2019	

**Labeler** - REVLON (788820165)**Registrant** - REVLON (788820165)**Establishment**

Name	Address	ID/FEI	Business Operations
Revlon		809725570	manufacture(10967-670)

Revised: 1/2024

REVLON