

TYLENOL COLD AND FLU- acetaminophen tablet, film coated
R J General Corporation

TYLENOL® COLD AND FLU

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> ▪ take 2 caplets every 4 hours ▪ swallow whole; do not crush, chew or dissolve ▪ do not take more than 10 caplets in 24 hours
children under 12 years	ask a doctor

Other information

- **each caplet contains:** sodium 3 mg
- store between 20-25°C (68-77°F)
- **do not use if pouch is torn or damaged**

Inactive ingredients

carnauba wax, croscarmellose sodium, D&C yellow no. 10 aluminum lake, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, sucralose, titanium dioxide

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Made in Italy

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA

Repackaged and distributed by:

RJ General
2024 Northwest Drive
Cincinnati OH 45231

PRINCIPAL DISPLAY PANEL - 100 Caplet Pouch Box

TYLENOL®
FOR ADULTS

COLD + FLU SEVERE

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl, Guaifenesin
Pain Reliever-Fever Reducer, Cough Suppressant, Nasal Decongestant, Expectorant

- **HEAD + BODY ACHES**
- **FEVER + SORE THROAT**
- **COUGH**
- **NASAL CONGESTION**
- **MUCUS + CHEST CONGESTION**

50 Pouches of 2 Caplets each



TYLENOL[®]

FOR ADULTS

COLD + FLU SEVERE

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl, Guaifenesin
Pain Reliever-Fever Reducer, Cough Suppressant, Nasal Decongestant, Expectorant



- HEAD + BODY ACHEs
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- MUCUS + CHEST CONGESTION



TYLENOL[®]

FOR ADULTS

COLD + FLU SEVERE

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl, Guaifenesin
Pain Reliever-Fever Reducer, Cough Suppressant, Nasal Decongestant, Expectorant



50 Pouches of 2 Caplets each

How can we help?
 1-877-895-3665



Drug Facts

Active Ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

- Uses**
- For the temporary relief of the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
 - temporarily reduces fever

- Warnings**
- Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take:
 - more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product

- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin redness
 - hives
 - rash
 If a skin reaction occurs, stop use and seek medical help right away.

Some throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

 - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 - if you have ever had an allergic reaction to this product or any of its ingredients.

- Ask a doctor before use if you have**
- liver disease
 - heart disease

Drug Facts (continued)

- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with bronchial phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dose.

- Stop use and ask a doctor if:**
- nervousness, dizziness, or sleeplessness occur
 - pain, nasal congestion or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see or read warning)

adults and children 12 years and over	take 2 caplets every 4 hours
children under 12 years	ask a doctor

swallow whole; do not crush, chew or dissolve
 do not take more than 10 caplets in 24 hours

Other information

- each caplet contains: sodium 3 mg
- store between 20-25°C (68-77°F)
- do not use if patch is torn or damaged

Inactive ingredients

camelina wax, croscarmellose sodium, D&C yellow no. 10 aluminum lake, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, saccharose, titanium dioxide

Questions or comments?
 call 1-877-895-3665 (toll-free) or 315-271-8755 (collect)

TYLENOL[®]

FOR ADULTS

COLD + FLU SEVERE

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl, Guaifenesin
 Pain Reliever-Fever Reducer, Cough Suppressant, Nasal Decongestant, Expectorant


Made in Italy
 Distributed by:
**JOHNSON & JOHNSON
 CONSUMER INC.**
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 © J&JCI 2021
 Visit us at www.tylenol.com
 or call toll-free 1-877-TYLENOL
 (1-877-895-3665)



Repackaged and distributed by:
 RJ General
 2024 Northwest Drive
 Cincinnati OH 45231

PARENTS:
 Learn about teen medicine choices





TYLENOL COLD AND FLU

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70264-025(NDC:50580-402)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	TYLENOL;SEVERE
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70264-025-02	50 in 1 BOX	05/01/2021	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2021	

Labeler - R J General Corporation (122542830)

Establishment

Name	Address	ID/FEI	Business Operations
R J General Corporation		122542830	repack(70264-025)

Establishment

Name	Address	ID/FEI	Business Operations
Johnson & Johnson		878046358	manufacture(70264-025)

Revised: 10/2023

R J General Corporation