

NAPROXEN- naproxen sodium tablet, film coated
Family Dollar Services Inc

Family Wellness 44-417

Active ingredient (in each tablet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - toothache
 - the common cold
 - muscular aches
 - headache
 - menstrual cramps
 - backache
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- skin reddening
- shock
- blisters
- rash
- facial swelling
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - feel faint
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - leg swelling
 - chest pain
 - slurred speech
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- adults and children 12 years and older
 - take 1 tablet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 tablets within the first hour
 - do not exceed 2 tablets in any 8- to 12-hour period
 - do not exceed 3 tablets in a 24-hour period
- children under 12 years: ask a doctor

Other information

- **each tablet contains:** sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, talc, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal display panel

**FAMILY
WELLNESS**

†COMPARE TO THE
ACTIVE INGREDIENT IN
ALEVE® TABLETS

**NAPROXEN
Naproxen Sodium Tablets, 220 mg**

Pain Reliever/Fever Reducer (**NSAID**)

STRENGTH TO LAST 12 HOURS

24 TABLETS

ACTUAL SIZE

NDC 55319-717-08

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by
Bayer HealthCare LLC, owner of the registered
trademark Aleve® Tablets. 50844 ORG122141708

**DISTRIBUTED BY: MIDWOOD BRANDS LLC,
500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA**

NOT 100% SATISFIED?

Return within 30 days to the
store of purchase for a refund
(with receipt) or exchange.

SKU 902886

0 32251 92378 4

4

FAMILY Wellness™

NAPROXEN

Naproxen Sodium Tablets, 220 mg
Pain Reliever/Fever Reducer (NSAID)

STRENGTH TO LAST 12 HOURS

24 TABLETS

NDC 55319-717-08

COMPARE TO THE ACTIVE INGREDIENT IN ALEVE® TABLETS

12 HOUR

ACTUAL SIZE

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts
Active ingredient (in each tablet) Purpose
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - toothache
 - the common cold
 - muscular aches
- headache
- menstrual cramps
- backache
- minor pain of arthritis

■ temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- shock
- asthma (wheezing)
- skin reddening
- hives
- blisters
- rash
- facial swelling

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- adults and children 12 years and older
- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period
- children under 12 years: ask a doctor

Other information

- each tablet contains sodium 20 mg
- store at 20°-25° C (68°-77° F). Avoid high humidity and excessive heat above 40° C (104° F).
- see end flap for expiration date and lot number

Inactive ingredients croscarmellose sodium, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, talc, titanium dioxide

Questions or comments?
Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ if right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk, if stomach upset occurs
- stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:
 - vomit blood
 - feel faint
 - have bloody or black stools
 - have stomach pain that does not get better
 - you have symptoms of heart problems or stroke:
 - leg swelling
 - chest pain
 - slurred speech
 - trouble breathing
 - weakness in one part or side of body
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear

NO PRINT Glue Area

NO PRINT Glue Area

TAPE EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

No print/No varnish Lot & Exp date

NOT 100% SATISFIED?
Return within 30 days to the store of purchase for a refund (with receipt) or exchange.

NOT 100% SATISFIED?
Return within 30 days to the store of purchase for a refund (with receipt) or exchange.

This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve® Tablets. 50844 6RG122141708

DISTRIBUTED BY: MIDWOOD BRANDS LLC,
500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA

B-0500-417-08-N
ORG122141708

Family_Wellness_44-417

NAPROXEN

naproxen sodium tablet, film coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:55319-717

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)		NAPROXEN SODIUM	220 mg	
Inactive Ingredients				
Ingredient Name			Strength	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)				
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	blue	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;417	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-717-08	1 in 1 CARTON	12/13/2023	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA204872	12/13/2023		

Labeler - Family Dollar Services Inc (024472631)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55319-717)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		832867837	manufacture(55319-717)
-------------------------	--	-----------	------------------------

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55319-717)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55319-717)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55319-717)

Revised: 12/2023

Family Dollar Services Inc