

**TETROXY 25- oxytetracycline hcl powder**  
**Bimeda, Inc.**

-----  
**Tetroxy® 25**

**(oxytetracycline HCl)**

**Soluble Powder**

**CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.**

***A broad spectrum ANTOBIOTIC*** □

**FOR CONTROL AND TREATMENT OF SPECIFIC DISEASES**

**IN POULTRY, CATTLE, SHEEP, SWINE AND HONEY BEES.**

**This packet contains 10 grams of oxytetracycline hydrochloride**

**For oral use in poultry, cattle, sheep, swine and honey bees.**

**FOR USE IN DRINKING WATER ONLY**

**NOT FOR USE IN LIQUID FEED SUPPLEMENTS**

**NOT FOR USE IN HUMANS**

**KEEP OUT OF REACH OF CHILDREN**

**Restricted Drug (California) - Use Only as Directed**

**ANADA 200-146, Approved by FDA**

**INDICATIONS AND DIRECTIONS FOR USE:**

For the control of the following poultry diseases caused by organisms susceptible to oxytetracycline: Add the following amount to two gallons of stock solution when proportioner is set to meter at the rate of one ounce per gallon.

CHICKENS	DOSAGE	PACKETS/2 GALLONS STOCK SOLUTION
Infectious synovitis caused by <i>Mycoplasma synoviae</i> .	200-400 mg/gal	5 - 10
Chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> .	400-800 mg/gal	10 - 20
Fowl cholera caused by <i>Pasteurella multocida</i> .	400-800 mg/gal	10 - 20
TURKEYS		
Hexamitiasis caused by <i>Hexamita meleagridis</i> .	200-400 mg/gal	5 - 10
Infectious synovitis caused by <i>Mycoplasma synoviae</i> .	400 mg/gal	10
Growing Turkeys – Complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis).	25 mg/lb body weight daily	varies with age & water consumption (1 packet will treat 400 lb of turkeys.)

Medicate continuously at the first clinical signs of disease and continue for 7 to 14 consecutive days. If improvement is not noted within 24-48 hours, consult a poultry diagnostic laboratory or poultry pathologist to determine diagnosis and advice on dosage. See Residue Warnings.

For the control and treatment of the following diseases caused by organisms susceptible to oxytetracycline:

SWINE	DOSAGE
Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> . Bacterial pneumonia caused by <i>Pasteurella multocida</i> . <b>For Breeding Swine:</b> Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
CALVES, BEEF CATTLE AND NON-LACTATING DAIRY CATTLE	
Bacterial enteritis caused by <i>Escherichia coli</i> . Bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
SHEEP	
Bacterial enteritis caused by <i>Escherichia coli</i> . Bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline HCl per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
<b>This packet will treat 1,000 lb of swine, cattle or sheep at 10 mg/lb</b>	
HONEY BEES	
For control of American Foulbrood caused by <i>Paenibacillus larvae</i> , and European Foulbrood caused by <i>Streptococcus pluton</i> susceptible to oxytetracycline.	200 mg/colony Administer in 3 applications of sugar syrup or 3 dustings at 4 to 5 day intervals. Apply dust on the outer parts or ends of the frames. See Residue Warnings.

**CONTRAINDICATION:** Dusting of uncapped brood cells has been reported to cause death of larval honey bees. Do not dust uncapped brood cells.

**CONTRAINDICATION:** Dusting of uncapped brood cells has been reported to cause death of larval honey bees. Do not dust uncapped brood cells.

**PRECAUTIONS:** Use as sole source of oxytetracycline. Prepare fresh solutions every 24 hours.

Special Note: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate and the environmental temperature and humidity, each of which affects water consumption.

**RESIDUE WARNING:** Do not administer turkeys, cattle or sheep within 5 days of slaughter. Zero-day slaughter withdrawal in swine. Do not administer to chickens or turkeys producing eggs for human consumption. Do not administer this product with milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. For honey bees, the drug should be fed early in the spring or fall and consumed by the bees before main honey flow begins to avoid contamination of production honey. Remove at least 6 weeks prior to main honey flow.

**STORAGE:** Store at 20°C to 25°C (68°F to 77°F), excursions permitted



between 15°C and 30°C (between 59°F and 86°F).



# **Tetroxy<sup>®</sup> 25**

**(oxytetracycline hydrochloride)**

## **Soluble Powder**

**CAUTION:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

*A broad spectrum ANTIBIOTIC*

**FOR CONTROL AND TREATMENT OF SPECIFIC DISEASES  
IN POULTRY, CATTLE, SHEEP, SWINE AND HONEY BEES.**

**This packet contains 10 grams of oxytetracycline  
hydrochloride**

**For oral use in poultry, cattle, sheep, swine and honey bees.**

**FOR USE IN DRINKING WATER ONLY  
NOT FOR USE IN LIQUID FEED SUPPLEMENTS  
NOT FOR USE IN HUMANS  
KEEP OUT OF REACH OF CHILDREN**

Restricted Drug (California) - Use Only as Directed  
ANADA 200-146, Approved by FDA



**Net**

**Weight: 181.5 g (6.4 oz)**



Manufactured by:  
Bimeda, Inc.  
Le Sueur, MN 56058



## Tetroxy<sup>®</sup> 25 Soluble Powder

(oxytetracycline hydrochloride)

### INDICATIONS AND DIRECTIONS FOR USE:

For the control of the following poultry diseases caused by organisms susceptible to oxytetracycline: Add the following amount to two gallons of stock solution when proportioner is set to meter at the rate of one ounce per gallon.

CHICKENS	DOSAGE	PACKETS/2 GALLONS STOCK SOLUTION
Infectious synovitis caused by <i>Mycoplasma synoviae</i> .	200-400 mg/gal	5 - 10
Chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> .	400-800 mg/gal	10 - 20
Fowl cholera caused by <i>Pasteurella multocida</i> .	400-800 mg/gal	10 - 20
TURKEYS		
Hexamitiasis caused by <i>Hexamita meleagridis</i> .	200-400 mg/gal	5 - 10
Infectious synovitis caused by <i>Mycoplasma synoviae</i> .	400 mg/gal	10
Growing Turkeys – Complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis).	25 mg/lb body weight daily	varies with age & water consumption (1 packet will treat 400 lb of turkeys.)

Medicate continuously at the first clinical signs of disease and continue for 7 to 14 consecutive days. If improvement is not noted within 24-48 hours, consult a poultry diagnostic laboratory or poultry pathologist to determine diagnosis and advice on dosage. See Residue Warnings.

For the control and treatment of the following diseases caused by organisms susceptible to oxytetracycline:

SWINE	DOSAGE
Bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> . Bacterial pneumonia caused by <i>Pasteurella multocida</i> . <b>For Breeding Swine:</b> Leptospirosis (reducing the incidence of abortions and shedding of leptospira) caused by <i>Leptospira pomona</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline hydrochloride per pound of body weight daily. Administer up to 14 days. See Residue Warnings.

### CALVES, BEEF CATTLE AND NON-LACTATING DAIRY CATTLE

Bacterial enteritis caused by <i>Escherichia coli</i> . Bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline hydrochloride per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
--	---

### SHEEP

Bacterial enteritis caused by <i>Escherichia coli</i> . Bacterial pneumonia (shipping fever complex) caused by <i>Pasteurella multocida</i> .	Administer in the drinking water at a level of 10 mg oxytetracycline hydrochloride per pound of body weight daily. Administer up to 14 days. See Residue Warnings.
--	---

**This packet will treat 1,000 lb of swine, cattle or sheep at 10 mg/lb**

### HONEY BEES

For control of American Foulbrood caused by <i>Paenibacillus larvae</i> susceptible to oxytetracycline.	200 mg/colony Administer in 3 applications of sugar syrup or 3 dustings at 4 to 5 day intervals. Apply dust on the outer parts or ends of the frames. See Residue Warnings.
---	--

**CONTRAINDICATION:** Dusting of uncapped brood cells has been reported to cause death of larval honey bees. Do not dust uncapped brood cells.

**PRECAUTION:** Use as sole source of oxytetracycline. Prepare fresh solutions every 24 hours.

Special Note: The concentration of drug required in medicated water must be adequate to compensate for variation in the age of the animal, feed consumption rate and the environmental temperature and humidity, each of which affects water consumption.

**RESIDUE WARNINGS:** Do not administer to turkeys, cattle or sheep within 5 days of slaughter. Zero-day slaughter withdrawal in swine. Do not administer to chickens or turkeys producing eggs for human consumption. Do not administer this product with milk or milk replacers. Administer 1 hour before or 2 hours after feeding milk or milk replacers. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. For honey bees, the drug should be fed early in the spring or fall and consumed by the bees before main honey flow begins to avoid contamination of production honey. Remove at least 6 weeks prior to main honey flow.

STORAGE: Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F).



**Bimeda**

Tetroxy<sup>®</sup> is a Registered Trademark of Bimeda, Inc.  
N.A. Corp. Address: Bimeda Inc., One Tower Lane, Oakbrook Terrace, IL 60181

Product No: 1TET044  
8TET042 Rev. 09/16

## TETROXY 25

oxytetracycline hcl powder

### Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:61133-5025
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYTETRACYCLINE HYDROCHLORIDE (UNII: 4U7K4N52ZM) (OXYTETRACYCLINE ANHYDROUS - UNII:SLF0D9077S)	OXYTETRACYCLINE HYDROCHLORIDE	10 g in 181.5 g

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61133-5025-1	181.5 g in 1 POUCH		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200146	09/28/2017	

**Labeler** - Bimeda, Inc. (060492923)

**Registrant** - Bimeda, Inc. (060492923)

### Establishment

Name	Address	ID/FEI	Business Operations
Bimeda, Inc.		060492923	manufacture

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 12/2022

Bimeda, Inc.