

ACID REDUCER- ranitidine tablet
CVS Pharmacy

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 150 mg (as ranitidine hydrochloride, USP 168 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PRINCIPAL DISPLAY PANEL

CVS pharmacy™

Compare to the active ingredient in Zantac 150®*

Maximum Strength

150 mg

ACID REDUCER

RANITIDINE TABLETS, USP 150 mg

Prevents & relieves heartburn associated with acid indigestion & sour stomach

200 TABLETS

Distributed by: CVS Pharmacy, Inc.

5108604/0214



5108604

CVS pharmacy Compare to the active ingredient in Zantac 150**

Maximum Strength **150 mg**

ACID REDUCER

RANITIDINE TABLETS, USP 150 mg

Prevents & relieves heartburn associated with acid indigestion & sour stomach

value size **200 tablets**

CVS pharmacy Compare to the active ingredient in Zantac 150**

Maximum Strength **150 mg**

ACID REDUCER

RANITIDINE TABLETS, USP 150 mg

Prevents & relieves heartburn associated with acid indigestion & sour stomach

200 TABLETS  Actual Size

See actual product size on side



Drug Facts

Active ingredient (in each tablet)	Purpose
Ranitidine 150 mg (as ranitidine hydrochloride, USP 168 mg)	Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water
 - to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

Questions? call 1-800-406-7884

All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Zantac 150. Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02895 © 2014 CVSpharmacy. CVS.com 1-800-SHOP CVS Made in IN/CA V-13977

0214  PLEASE RECYCLE

#937138



0 50428 46505 9

Batch No. Expiration Date:

Non Varnish Area

t098015



ACID REDUCER			
ranitidine tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-354
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg
Inactive Ingredients			

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	OVAL (Oval Shaped)	Size	12mm
Flavor		Imprint Code	9R
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-354-65	1 in 1 CARTON		
1		65 in 1 BOTTLE		
2	NDC:59779-354-95	1 in 1 CARTON		
2		95 in 1 BOTTLE		
3	NDC:59779-354-20	1 in 1 CARTON		
3		200 in 1 BOTTLE		
4	NDC:59779-354-24	24 in 1 CARTON		
4		1 in 1 BLISTER PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200536	08/21/2012	

Labeler - CVS Pharmacy (062312574)

Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	manufacture(59779-354)