

PENNCHLOR 90G- chlortetracycline calcium granule
Pharmgate Animal Health

Pennchlor® 90 G™

Chlortetracycline
 Type A Medicated Article

Active Drug Ingredients: Chlortetracycline as Chlortetracycline Calcium Complex equivalent to 90 grams Chlortetracycline Hydrochloride/lb.

Ingredients: Chlortetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.

For Use in the Manufacture of Medicated Feeds.

CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS

Directions For Use

Indications For Use	Use Level of Chlortetracycline	lbs. of Pennchlor 90 G® per ton
<p>Chickens</p> <p>For Chickens: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline. (Feed continuously for 7 - 14 days)</p> <p>WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p>	100-200 g/ton	1.11 - 2.22
<p>For Chickens: Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. (Feed continuously for 7 - 14 days).</p> <p>WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p>	200-400 g/ton	2.22 - 4.44
<p>For Chickens: Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline (Feed for 5 days).</p> <p>WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. WITHDRAW 24 HOURS PRIOR TO SLAUGHTER.</p>	500 g/ton	5.55

Turkeys		
<p>For Turkeys: Control of infectious synovitis caused by <i>Mycoplasma synoviae</i> susceptible to chlortetracycline (Feed continuously for 7 - 14 days)</p> <p>WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p>	200 g/ton	2.22
<p>For Turkeys: Control of hexamitiasis caused by <i>Hexamita meleagrides</i> susceptible to chlortetracycline. (Feed continuously for 7 - 14 days).</p> <p>WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p>	400 g/ton	4.44
<p>For Turkeys: Turkey poults not over 4 weeks of age: Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline.</p>	400 g/ton	4.44
<p>For Turkeys: Control of complicating bacterial organisms associated with bluecomb (transmissible enteritis, coronaviral enteritis) susceptible to chlortetracycline. (Feed continuously for 7 - 14 days)</p> <p>WARNINGS: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHDRAWAL PERIOD.</p>	25 mg/lb body weight/day	
Swine		
<p>For Swine: Reducing the incidence of cervical lymphadenitis (jowl abscesses) caused by <i>Group E Streptococci</i> susceptible to chlortetracycline.</p>	50-100 g/ton	0.555 - 1.11
<p>For Breeding Swine: Control of leptospirosis (reducing the instances of abortions and shedding of leptospirae) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. (Feed continuously for not more than 14 days)</p>	400 g/ton	4.44
<p>For Swine: Treatment of bacterial enteritis caused by <i>Escherichia coli</i> and <i>Salmonella choleraesuis</i> and bacterial pneumonia caused by <i>Pasteurella multocida</i> susceptible to chlortetracycline. (Feed for not more than 14 days.)</p> <p>WARNING: ZERO-DAY WITHDRAWAL PERIOD.</p>	Feed approximately 400 g/t, varying with body weight and feed consumption to provide 10 mg/lb body weight/day.	

**CALVES, BEEF CATTLE, AND
NONLACTATING DAIRY CATTLE WARNING:
A WITHDRAWAL PERIOD HAS NOT BEEN
ESTABLISHED FOR THIS PRODUCT IN PRE-
RUMINATING CALVES. DO NOT USE IN
CALVES TO BE PROCESSED FOR VEAL.**

For Growing Cattle (over 400 lbs.): For the reduction of the incidence of liver abscesses.

70 mg/head/day

WARNING: ZERO-DAY WITHDRAWAL PERIOD.

For Beef Cattle: For the control of bacterial pneumonia associated with shipping fever complex caused by *Pasteurella* spp. susceptible to chlortetracycline

350 mg/head/day

WARNING: ZERO-DAY WITHDRAWAL PERIOD.

For Beef Cattle (under 700 lbs.): Control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline.

350 mg/head/day

WARNING: ZERO-DAY WITHDRAWAL PERIOD.

For Beef Cattle (over 700 lbs.): Control of active infection of anaplasmosis caused by *Anaplasma marginale* susceptible to chlortetracycline.

0.5 mg/ lb. body weight/day

WARNING: ZERO-DAY WITHDRAWAL PERIOD.

For Calves, Beef, and Nonlactating Dairy Cattle: For treatment of bacterial enteritis caused by *Escherichia coli* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to chlortetracycline. (Treat for not more than 5 days)

10 mg/lb. body weight/day

WARNING: ZERO-DAY WITHDRAWAL PERIOD.

Sheep

For Breeding Sheep: Reducing the incidence of (vibriotic) abortion caused by *Campylobacter fetus* infections susceptible to chlortetracycline.

80 mg/head/day

WARNING: ZERO-DAY WITHDRAWAL PERIOD.

Approved by FDA under NADA # 138-935

Pennchlor 90 G® is a registered trademark of Pharmgate Inc.

Marketed by:
Pharmgate Animal Health, LLC
14040 Industrial Road
Omaha, NE 68144



Take Time

Observe Directions

PRINCIPAL DISPLAY PANEL - 22.68 Kg Bag Label

Pennchlor[®] 90 G[™]

Chlortetracycline

Type A Medicated Article

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

See mixing directions, claims, cautions and warnings on back.

Restricted Drug (California) - Use Only as Directed.

Not For Human Use.

Net wt. 50 LB (22.68 Kg)

Pharmgate
ANIMAL HEALTH



Pharmgate
ANIMAL HEALTH

Pennchlor[®] 90 G[™]

Pennchlor[®] 90 G[™]



Chlortetracycline Type A Medicated Article

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

See mixing directions, claims, cautions and warnings on back.
Restricted Drug (California) - Use Only as Directed.
Not For Human Use.

Net wt. 50 LB (22.68 Kg)




Pharmgate
ANIMAL HEALTH

Pennchlor 90 G™
 ingate™

Pennchlor 90 G™

Chlortetracycline
Type A Medicated Article

Active Drug Ingredients		Chlortetracycline as Chlortetracycline Calcium Complex equivalent to 90 grams Chlortetracycline Hydrochloride/lb.			
Ingredients		Chlortetracycline, Calcium Carbonate, Roughage Products and Mineral Oil.			
For Use in the Manufacture of Medicated Feeds					
CAUTION: For use in Dry Feeds ONLY. NOT FOR USE IN LIQUID FEED SUPPLEMENTS					
Directions For Use					
Indications For Use	The Levels of Chlortetracycline	No. of Pounds per 100 lbs.	Indications For Use	The Levels of Chlortetracycline	No. of Pounds per 100 lbs.
Chickens For Chickens: Control of infectious epizootic caused by <i>Mycoplasma agropyri</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHERAWAL PERIOD. For Chickens: Control of chronic respiratory disease (CRD) and air sac infection caused by <i>Mycoplasma gallisepticum</i> and <i>Escherichia coli</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHERAWAL PERIOD. For Chickens: Reduction of mortality due to <i>Escherichia coli</i> infections susceptible to chlortetracycline. (Feed for 5-days) WARNING: DO NOT FEED TO CHICKENS PRODUCING EGGS FOR HUMAN CONSUMPTION. WITHERAWAL 9 HOURS PRIOR TO SLAUGHTER.	300-300g / km 200-400g / km 300g / km	1.13-2.22 2.25-4.44 5.05	Swine For Swine: Reducing the incidence of cervical lymphadenitis (owl abscesses) caused by Group E <i>Streptococci</i> susceptible to chlortetracycline. For Breeding Swine: Control of leptospirosis (precluding the instances of abortions and shedding of leptospires) caused by <i>Leptospira pomona</i> susceptible to chlortetracycline. (Feed continuously for not more than 14 days) For Swine: Treatment of bacterial enteritis caused by <i>Zootherphila coli</i> and <i>Salmonella choleraesuis</i> and bacterial proventriculitis caused by <i>Haemophilus vulgare</i> susceptible to chlortetracycline. (Feed for not more than 14 days) WARNING: ZERO-DAY WITHERAWAL PERIOD.	50-100g / km 400g / km Feed approximately 400 g/l, varying with body weight and feed consumption to provide 10mg/lb body weight/day	0.035-1.13 4.44
Turkeys For Turkeys: Control of infectious epizootic caused by <i>Mycoplasma agropyri</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHERAWAL PERIOD. For Turkeys: Control of hepatitis caused by <i>Haematele magna</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHERAWAL PERIOD. For Turkeys: Turkey production over 4 weeks of age. Reduction of mortality due to paratyphoid caused by <i>Salmonella typhimurium</i> susceptible to chlortetracycline. For Turkeys: Control of complicating bacterial organisms associated with cloacal <i>Escherichia coli</i> , <i>Escherichia coli</i> , <i>Escherichia coli</i> , <i>Escherichia coli</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO TURKEYS PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHERAWAL PERIOD.	200g / km 400g / km 400g / km 25mg / lb body weight/day	2.22 4.44 4.44	CATTLE, BEEF CATTLE, AND NON-LACTATING DAIRY CATTLE WARNING: A WITHERAWAL PERIOD HAS NOT BEEN ESTABLISHED FOR THIS PRODUCT IN THIS SPECIFIC INDICATION. DO NOT USE IN CATTLE TO BE REPRODUCED FOR VEAL. For Grazing Cattle (over 400 lbs.): For the reduction of the incidence of liver abscesses. WARNING: ZERO-DAY WITHERAWAL PERIOD. For Beef Cattle: For the control of bacterial proventriculitis associated with abomasal lesions caused by <i>Haemophilus spp.</i> susceptible to chlortetracycline. WARNING: ZERO-DAY WITHERAWAL PERIOD. For Beef Cattle (under 700 lbs.): Control of active infection of esophagitis caused by <i>Aspergillus niger</i> susceptible to chlortetracycline. WARNING: ZERO-DAY WITHERAWAL PERIOD. For Beef Cattle (over 700 lbs.): Control of active infection of esophagitis caused by <i>Aspergillus niger</i> susceptible to chlortetracycline. WARNING: ZERO-DAY WITHERAWAL PERIOD. For Calves, Beef, and Non-lactating Dairy Cattle: For treatment of bacterial enteritis caused by <i>Escherichia coli</i> and bacterial proventriculitis caused by <i>Haemophilus vulgare</i> susceptible to chlortetracycline. (Feed for not more than 5 days) WARNING: ZERO-DAY WITHERAWAL PERIOD.	70mg / head / day 100mg / head / day 200mg / head / day 0.5mg / lb body weight / day 30mg / lb body weight / day	
Sheep For Sheep: Control of bacterial enteritis caused by <i>Escherichia coli</i> susceptible to chlortetracycline. (Feed continuously for 7-14 days) WARNING: DO NOT FEED TO SHEEP PRODUCING EGGS FOR HUMAN CONSUMPTION. ZERO-DAY WITHERAWAL PERIOD.	25mg / lb body weight/day				



PENNCHLOR 90G

chlortetracycline calcium granule

Product Information

Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:51429-091
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Chlortetracycline Calcium (UNII: NR4B2SX17S) (Chlortetracycline - UNII:WCK1KIQ23Q)	Chlortetracycline Hydrochloride	198 g in 1 kg

Inactive Ingredients

Ingredient Name	Strength
Calcium Carbonate (UNII: H0G9379FGK)	
Mineral Oil (UNII: T5L8T28FGP)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51429-091-01	22.68 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA138935	04/11/2007	

Labeler - Pharmgate Animal Health (833270817)

Registrant - Pharmgate Inc. (079628671)

Revised: 2/2023

Pharmgate Animal Health