METFORMIN HYDROCHLORIDE - metformin hydrochloride tablet, film coated
MARKSANS PHARMA LIMITED

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use Metformin Hydrochloride Tablets safely and effectively. See full prescribing information for Metformin Hydrochloride Tablets.
Metformin Hydrochloride Tablets for oral use
Initial U.S. Approval: 1995

WARNING: LACTIC ACIDOSIS
See full prescribing information for complete boxed warning.

- **Postmarketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. Symptoms included malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Laboratory abnormalities included elevated blood lactate levels, anion gap acidosis, increased lactate/pyruvate ratio; and metformin plasma levels generally >5 mcg/mL. (5.1)**
- **Risk factors include renal impairment, concomitant use of certain drugs, age >65 years old, radiological studies with contrast, surgery and other procedures, hypoxic states, excessive alcohol intake, hepatic impairment and mitochondrial diseases. Steps to reduce the risk of and manage metformin-associated lactic acidosis in these high risk groups are provided in the Full Prescribing Information. (5.1)**
- **If lactic acidosis is suspected, discontinue Metformin Hydrochloride Tablets and institute general supportive measures in a hospital setting. Prompt hemodialysis is recommended. (5.1)**

RECENT MAJOR CHANGES

Boxed Warning, Warnings and Precautions (5.1)
06/2026

INDICATIONS AND USAGE

Metformin Hydrochloride Tabletis a biguanide indicated as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients 10 years of age and older with type 2 diabetes mellitus. (1)

DOSAGE AND ADMINISTRATION

Adult Dosage for Metformin Hydrochloride Tablets (2)

- Starting dose: 500 mg orally twice a day or 850 mg once a day, with meals (2.1)
- Increase the dose in increments of 500 mg weekly or 850 mg every 2 weeks, up to a maximum dose of 2550 mg per day, given in divided doses (2.1)
- Doses above 2000 mg may be better tolerated given 3 times a day with meals (2.1)

Pediatric Dosage for Metformin Hydrochloride Tablets (2)

- Starting dose: 500 mg orally twice a day, with meals (2.2)
- Increase dosage in increments of 500 mg weekly up to a maximum of 2000 mg per day, given in divided doses (2.2)

Renal Impairment: (2)

- Prior to initiation, assess renal function with estimated glomerular filtration rate (eGFR) (2.3)
- Do not use in patients with eGFR below 30 mL/minute/1.73 m^2 (2.3)
- Initiation is not recommended in patients with eGFR between 30-45 mL/minute/1.73 m^2 (2.3)
- Assess risk/benefit of continuing if eGFR falls below 45mL/minute/1.73 m^2 (2.3)
- Discontinue if eGFR falls below 30 mL/minute/1.73 m^2 (2.3)

Discontinuation for Iodinated Contrast Imaging Procedures: (2)

- Metformin Hydrochloride Tabletsmay need to be discontinued at time of, or prior to, iodinated contrast imaging procedures (2.4)

DOSAGE FORMS AND STRENGTHS

Metformin Hydrochloride Tablets: 500 mg, 850 mg and 1000 mg (3)

CONTRAINDICATIONS

- Severe renal impairment (eGFR below 30 mL/min/1.73 m^2) (4, 5.1)
- Hypersensitivity to metformin (4)
- Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma. (4)

WARNINGS AND PRECAUTIONS

- *Lactic Acidosis:* See boxed warning. (5.1)
- *Vitamin B_{12} Deficiency:* Metformin may lower vitamin B_{12} levels. Measure hematological parameters annually and vitamin B_{12} at 2 to 3 year intervals and manage any abnormalities. (5.2)
- *Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues:* Increased risk of hypoglycemia when used in combination with insulin and/or an insulin secretagogue. Lower dose of insulin or insulin secretagogue may be required (5.3)

ADVERSE REACTIONS

For Metformin Hydrochloride Tablets, the most common adverse reactions (>5.0%) are diarrhea, nausea/vomiting, flatulence, asthenia, indigestion, abdominal discomfort, and headache. (6.1)
To report SUSPECTED ADVERSE REACTIONS, contact Marksans at 1-877-376-4271 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

DRUG INTERACTIONS

- Carbonic anhydrase inhibitors may increase risk of lactic acidosis. Consider more frequent monitoring (7)
- Drugs that reduce metformin clearance (such as ranolazine, vandetanib, dolutegravir, and cimetidine) may increase the accumulation of metformin. Consider the benefits and risks of concomitant use (7)
- Alcohol can potentiate the effect of metformin on lactate metabolism. Warn patients against excessive alcohol intake (7)

USE IN SPECIFIC POPULATIONS

- Females and Males of Reproductive Potential: Advise premenopausal females of the potential for an unintended pregnancy. (8.3)
- Geriatric Use: Assess renal function more frequently. (8.5)
- Hepatic Impairment: Avoid use in patients with hepatic impairment. (8.7)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 6/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

BOXED WARNING
1 INDICATIONS & USAGE
2 DOSAGE & ADMINISTRATION
3 DOSAGE FORMS & STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
6 ADVERSE REACTIONS
7 DRUG INTERACTIONS
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.3 FEMALES & MALES OF REPRODUCTIVE POTENTIAL SECTION
8.4 Pediatric Use
8.5 Geriatric Use
8.6 RENAL IMPAIRMENT
8.7 HEPATIC IMPAIRMENT
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.3 Pharmacokinetics
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis & Mutagenesis & Impairment Of Fertility

14 CLINICAL STUDIES
16 HOW SUPPLIED/STORAGE AND HANDLING
17 PATIENT COUNSELING INFORMATION
* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

BOXED WARNING

WARNING: LACTIC ACIDOSIS

Postmarketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. The onset of metformin-associated lactic acidosis is often subtle, accompanied only by nonspecific symptoms such as malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Metformin-associated lactic acidosis was characterized by elevated blood lactate levels (>5 mmol/Liter), anion gap acidosis (without evidence of ketonuria or ketonemia), an increased lactate/pyruvate ratio; and metformin plasma levels generally >5 mcg/mL *[see Warnings and Precautions (5.1)].*

Risk factors for metformin-associated lactic acidosis include renal impairment, concomitant use of certain drugs (e.g. carbonic anhydrase inhibitors such as topiramate), age 65 years old or greater, having a radiological study with contrast, surgery and other procedures, hypoxic states (e.g., acute congestive heart failure), excessive alcohol intake, hepatic impairment and mitochondrial diseases *[see Warnings and Precautions (5.1)].*

Steps to reduce the risk of and manage metformin-associated lactic acidosis in these high risk groups are provided *[seeDosage and Administration (2.3), Contraindications (4), Warnings and Precautions (5.1)].*

If metformin-associated lactic acidosis is suspected, immediately discontinue Metformin Hydrochloride Tabletsand institute general supportive measures in a hospital setting. Prompt haemodialysis is recommended *[see Warnings and Precautions (5.1)].*

1 INDICATIONS & USAGE

Metformin Hydrochloride Tablets, USP is indicated as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients 10 years of age and older with type 2 diabetes mellitus.

2 DOSAGE & ADMINISTRATION

2.1 Adult Dosage

- The recommended starting dose of Metformin Hydrochloride Tablet is 500 mg orally twice a day or 850 mg once a day, given with meals.
- Increase the dose in increments of 500 mg weekly or 850 mg every 2 weeks on the basis of glycemic control and tolerability, up to a maximum dose of 2550 mg per day, given in divided doses.
- Doses above 2000 mg may be better tolerated given 3 times a day with meals.

2.2 Pediatric Dosage for Metformin Hydrochloride Tablet

- The recommended starting dose of Metformin Hydrochloride Tablet for pediatric patients 10 years of age and older is 500 mg orally twice a day, given with meals.
- Increase dosage in increments of 500 mg weekly on the basis of glycemic control and tolerability, up to a maximum of 2000 mg per day, given in divided doses twice daily.

2.3 Recommendations for Use in Renal Impairment

- Assess renal function prior to initiation of Metformin Hydrochloride Tablets and periodically thereafter.
- Metformin Hydrochloride Tablets are contraindicated in patients with an estimated glomerular filtration rate (eGFR) below 30 mL/minute/1.73 m^2.
- Initiation of Metformin Hydrochloride Tablets in patients with an eGFR between 30 – 45 mL/minute/1.73 m^2 is not recommended.
- In patients taking Metformin Hydrochloride Tablets whose eGFR later falls below 45 mL/min/1.73 m^2, assess the benefit risk of continuing therapy.
- Discontinue Metformin Hydrochloride Tablets if the patient's eGFR later falls below 30 mL/minute/1.73 m^2 *[see Warnings and Precautions (5.1)].*

2.4 Discontinuation for Iodinated Contrast Imaging Procedures

Discontinue Metformin Hydrochloride Tablets at the time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR between 30 and 60 mL/min/1.73 m2; in patients with a history of liver disease, alcoholism, or heart failure; or in patients who will be administered intraarterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure; restart Metformin Hydrochloride Tablets if renal function is stable.

3 DOSAGE FORMS & STRENGTHS

Metformin Hydrochloride Tablets are available as:

- *500 mg:* White to off-white colored, film coated, round shaped tablet with beveled edges on both side and debossed with '134' on one side and plain on other side.
- *850 mg:* White to off-white colored, film coated, round shaped tablet with beveled edges on both side and debossed with '131' on one side and plain on other side.
- *1000 mg:* White to off-white colored, film coated, oval shaped tablet with '132' debossed on one side and breakline on both sides.

4 CONTRAINDICATIONS

Metformin Hydrochloride Tablets are contraindicated in patients with:

- Severe renal impairment (eGFR below 30 mL/min/1.73 m^2) [see Warnings and Precautions (5.1)].
- Hypersensitivity to metformin
- Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma.

5 WARNINGS AND PRECAUTIONS

5.1 Lactic Acidosis

There have been postmarketing cases of metformin-associated lactic acidosis, including fatal cases. These cases had a subtle onset and were accompanied by nonspecific symptoms such as malaise, myalgias, abdominal pain, respiratory distress, or increased somnolence; however, hypotension and resistant bradyarrhythmias have occurred with

severe acidosis. Metformin-associated lactic acidosis was characterized by elevated blood lactate concentrations (>5 mmol/L), anion gap acidosis (without evidence of ketonuria or ketonemia), and an increased lactate: pyruvate ratio; metformin plasma levels were generally >5 mcg/mL. Metformin decreases liver uptake of lactate increasing lactate blood levels which may increase the risk of lactic acidosis, especially in patients at risk.
If metformin-associated lactic acidosis is suspected, general supportive measures should be instituted promptly in a hospital setting, along with immediate discontinuation of metformin hydrochloride tablets. In metformin hydrochloride tablets, treated patients with a diagnosis or strong suspicion of lactic acidosis, prompt hemodialysis is recommended to correct the acidosis and remove accumulated metformin (metformin hydrochloride is dialyzable with a clearance of up to 170 mL/min under good hemodynamic conditions). Hemodialysis has often resulted in reversal of symptoms and recovery.

Educate patients and their families about the symptoms of lactic acidosis and, if these symptoms occur, instruct them to discontinue metformin hydrochloride tablets and report these symptoms to their healthcare provider.

For each of the known and possible risk factors for metformin-associated lactic acidosis, recommendations to reduce the risk of and manage metformin-associated lactic acidosis are provided below:

- *Renal impairment*—The postmarketing metformin-associated lactic acidosis cases primarily occurred in patients with significant renal impairment.

The risk of metformin accumulation and metformin-associated lactic acidosis increases with the severity of renal impairment because metformin is substantially excreted by the kidney. Clinical recommendations based upon the patient's renal function include [*see Dosage and Administration (2.1), Clinical Pharmacology (12.3)]:*

- Before initiating metformin hydrochloride tablets, obtain an estimated glomerular filtration rate (eGFR)
- Metformin hydrochloride tablets are contraindicated in patients with an eGFR less than 30 mL/min/1.73 m^2 *[see Contraindications (4)]* .
- Initiation of metformin hydrochloride tablets are not recommended in patients with eGFR between 30-45 mL/min/1.73 m^2.
- Obtain an eGFR at least annually in all patients taking metformin hydrochloride tablets.In patients at risk for the development of renal impairment (e.g., the elderly), renal function should be assessed more frequently.
- In patients taking metformin hydrochloride tablets whose eGFR falls below 45 mL/min/1.73 m^2, assess the benefit and risk of continuing therapy.

- *Drug interactions*—The concomitant use of metformin hydrochloride tablets with specific drugs may increase the risk of metformin-associated lactic acidosis: those that impair renal function, result in significant hemodynamic change, interfere with acid-base balance, or increase metformin accumulation. Consider more frequent monitoring of patients.

- *Age 65 or greater*—The risk of metformin-associated lactic acidosis increases with the patient's age because elderly patients have a greater likelihood of having hepatic, renal, or cardiac impairment than younger patients. Assess renal function more frequently in elderly patients.

- *Radiologic studies with contrast*— Administration of intravascular iodinated contrast agents in metformin-treated patients has led to an acute decrease in renal function and the occurrence of lactic acidosis. Stop metformin hydrochloride tablets at the

time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR between 30 and 60 mL/min/1.73 m^2; in patients with a history of hepatic impairment, alcoholism or heart failure; or in patients who will be administered intra-arterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure, and restart metformin hydrochloride tablets if renal function is stable.

- *Surgery and other procedures*—Withholding of food and fluids during surgical or other procedures may increase the risk for volume depletion, hypotension, and renal impairment. Metformin hydrochloride tablets should be temporarily discontinued while patients have restricted food and fluid intake.
- *Hypoxic states*—Several of the postmarketing cases of metformin-associated lactic acidosis occurred in the setting of acute congestive heart failure (particularly when accompanied by hypoperfusion and hypoxemia). Cardiovascular collapse (shock), acute myocardial infarction, sepsis, and other conditions associated with hypoxemia have been associated with lactic acidosis and may cause prerenal azotemia. When such an event occurs, discontinue metformin hydrochloride tablets.
- *Excessive alcohol intake*—Alcohol potentiates the effect of metformin on lactate metabolism. Patients should be warned against excessive alcohol intake while receiving metformin hydrochloride tablets.
- *Hepatic impairment*—Patients with hepatic impairment have developed cases of metformin associated lactic acidosis. This may be due to impaired lactate clearance resulting in higher lactate blood levels. Therefore, avoid use of metformin hydrochloride tablets in patients with clinical or laboratory evidence of hepatic disease.
- *Mitochondrial Diseases*—Metformin may increase the risk of lactic acidosis in patients with mitochondrial diseases such as mitochondrial encephalomyopathy with lactic acidosis, and stroke-like episodes (MELAS) syndrome and maternally inherited diabetes and deafness (MIDD).Metformin Hydrochloride tablet is not recommended in patients with mitochondrial diseases such as MELAS syndrome and MIDD.
- If a patient taking Metformin Hydrochloride Tablet develops elevated lactic acid levels (with or without accompanying neurologic signs and symptoms), discontinue Metformin Hydrochloride Tablet. Depending on the clinical context, consider evaluating the patient for MELAS, MIDD, or other mitochondrial disease.

5.2 Vitamin B_{12} Deficiency

In metformin hydrochloride tablets clinical trials of 29 weeks duration, a decrease to subnormal levels of previously normal serum vitamin B_{12} levels was observed in approximately 7% of patients. Such decrease, possibly due to interference with B_{12} absorption from the B_{12}-intrinsic factor complex, may be associated with anemia but appears to be rapidly reversible with discontinuation of metformin hydrochloride tablets or vitamin B_{12} supplementation. Certain individuals (those with inadequate vitamin B_{12} or calcium intake or absorption) appear to be predisposed to developing subnormal vitamin B_{12} levels. Measure hematologic parameters on an annual basis vitamin B_{12} at 2 to 3 year intervals in patients on metformin hydrochloride tablets and manage any abnormalities *[see Adverse Reactions (6.1)].*

5.3 Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues

Insulin and insulin secretagogues (e.g., sulfonylurea) are known to cause hypoglycemia. Metformin hydrochloride tablets may increase the risk of hypoglycemia when combined with insulin and/or an insulin secretagogue. Therefore, a lower dose of insulin or insulin secretagogue may be required to minimize the risk of hypoglycemia when used in combination with metformin hydrochloride tablets *[see Drug Interactions (7)].*

5.4 Macrovascular outcomes

There have been no clinical studies establishing conclusive evidence of macrovascular risk reduction with metformin hydrochloride tablets.

6 ADVERSE REACTIONS

The following adverse reactions are also discussed elsewhere in the labeling:

- Lactic Acidosis *[see Boxed Warning and Warnings and Precautions (5.1)]*
- Vitamin B_{12} Deficiency *[see Warnings and Precautions (5.2)]*
- Hypoglycemia *[see Warnings and Precautions (5.3)]*

6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Metformin Hydrochloride Tablets

In a U.S. clinical trial of metformin hydrochloride tablets in patients with type 2 diabetes mellitus, a total of 141 patients received metformin hydrochloride tablets up to 2550 mg per day. Adverse reactions reported in greater than 5% of metformin hydrochloride tablets treated patients and that were more common than in placebo-treated patients, are listed in Table 1.

Table 1: Adverse Reactions from Clinical Trial of Metformin Hydrochloride Tablets Occurring >5% and More Common than Placebo in Patients with Type 2 Diabetes Mellitus

	Metformin Hydrochloride Tablet (n=141)	**Placebo (n=145)**
Diarrhea	53%	12%
Nausea/Vomiting	26%	8%
Flatulence	12%	6%
Asthenia	9%	6%
Indigestion	7%	4%
Abdominal Discomfort	6%	5%
Headache	6%	5%

Diarrhea led to discontinuation of metformin hydrochloride tablets

in 6% of patients. Additionally, the following adverse reactions were reported in ≥1% to ≤5.0% of metformin hydrochloride tablets

treated patients and were more commonly reported with metformin hydrochloride tablets than placebo: abnormal stools, hypoglycemia, myalgia, lightheaded, dyspnea, nail disorder, rash, sweating increased, taste disorder, chest discomfort, chills, flu syndrome, flushing, palpitation.

In metformin hydrochloride tabletsclinical trials of 29-week duration, a decrease to subnormal levels of previously normal serum vitamin B_{12} levels was observed in approximately 7% of patients.

Pediatric Patients

In clinical trials with metformin hydrochloride tabletsin pediatric patients with type 2 diabetes mellitus, the profile of adverse reactions was similar to that observed in adults.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of metformin. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Cholestatic, hepatocellular, and mixed hepatocellular liver injury have been reported with postmarketing use of metformin.

7 DRUG INTERACTIONS

Table 2 presents clinically significant drug interactions with metformin hydrochloride tablets.

Table 2: Clinically Significant Drug Interactions with Metformin Hydrochloride Tablets

Carbonic Anhydrase Inhibitors	
Clinical Impact:	Carbonic anhydrase inhibitors frequently cause a decrease in serum bicarbonate and induce non-anion gap, hyperchloremic metabolic acidosis. Concomitant use of these drugs with metformin hydrochloride tablets may increase the risk for lactic acidosis.
Intervention:	Consider more frequent monitoring of these patients.
Examples:	Topiramate, zonisamide, acetazolamide or dichlorphenamide.
Drugs that Reduce Metformin Hydrochloride TabletsClearance	
Clinical Impact:	Concomitant use of drugs that interfere with common renal tubular transport systems involved in the renal elimination of metformin (e.g., organic cationic transporter-2 [OCT2] / multidrug and toxin extrusion [MATE] inhibitors) could increase systemic exposure to metformin and may increase the risk for lactic acidosis *[see Clinical Pharmacology (12.3)].*
Intervention:	Consider the benefits and risks of concomitant use with metformin hydrochloride tablets.
Examples:	Ranolazine, vandetanib, dolutegravir, and cimetidine.
Alcohol	
Clinical Impact:	Alcohol is known to potentiate the effect of metformin on lactate metabolism.
Intervention:	Warn patients against excessive alcohol intake while receiving metformin hydrochloride tablets.
Insulin Secretagogues or Insulin	
Clinical Impact:	Coadministration of metformin hydrochloride tablets with an insulin secretagogue (e.g., sulfonylurea) or insulin may increase the risk of hypoglycemia.
Intervention:	Patients receiving an insulin secretagogue or insulin may require lower doses of the insulin secretagogue or insulin.
Drugs Affecting Glycemic Control	
Clinical Impact:	Certain drugs tend to produce hyperglycemia and may lead to loss of glycemic control.
Intervention:	When such drugs are administered to a patient metformin hydrochloride tablets, observe the patient closely for loss of blood glucose control. When such drugs are withdrawn from a patient receiving metformin hydrochloride tablets, observe the patient closely for hypoglycemia.
Examples:	Thiazides and other diuretics, corticosteroids, phenothiazines, thyroid products, estrogens, oral contraceptives, phenytoin, nicotinic acid, sympathomimetics, calcium channel blockers, and isoniazid.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk summary

Limited data with metformin hydrochloride tabletsin pregnant women are not sufficient to determine a drug-associated risk for major birth defects or miscarriage. Published studies with metformin use during pregnancy have not reported a clear association with metformin and major birth defect or miscarriage risk *[see Data].* There are risks to the mother and fetus associated with poorly controlled diabetes mellitus in pregnancy *[see Clinical Considerations].*

No adverse developmental effects were observed when metformin was administered to pregnant Sprague Dawley rats and rabbits during the period of organogenesis at doses up to 2- and 5 times, respectively, a 2550 mg clinical dose, based on body surface area *[see Data].*

The estimated background risk of major birth defects is 6–10% in women with pre-gestational diabetes mellitus with an HbA1C >7 and has been reported to be as high as 20–25% in women with a HbA1C >10. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2–4% and 15–20%, respectively.

Clinical Considerations

Disease-associated maternal and/or embryo/fetal risk

Poorly-controlled diabetes mellitus in pregnancy increases the maternal risk for diabetic ketoacidosis, pre-eclampsia, spontaneous abortions, preterm delivery, stillbirth and delivery complications. Poorly controlled diabetes mellitus increases the fetal risk for major birth defects, stillbirth, and macrosomia related morbidity.

Data

Human Data

Published data from post-marketing studies have not reported a clear association with metformin and major birth defects, miscarriage, or adverse maternal or fetal outcomes when metformin was used during pregnancy. However, these studies cannot definitely establish the absence of any metformin-associated risk because of methodological limitations, including small sample size and inconsistent comparator groups.

Animal Data

Metformin hydrochloride did not adversely affect development outcomes when administered to pregnant rats and rabbits at doses upto 600 mg/kg/day. This represents an exposure of about 2 and 5 times a 2550 mg clinical dose based on body surface area comparisons for rats and rabbits,respectively. Determination of fetal concentrations demonstrated a partial placental barrier to metformin.

8.2 Lactation

Risk Summary

Limited published studies report that metformin is present in human milk *[see Data]*.However, there is insufficient information to determine the effects of metformin on the breastfed infant and no available information on the effects of metformin on milk

production. Therefore, the developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for metformin hydrochloride tablets and any potential adverse effects on the breastfed child from metformin hydrochloride tablets or from the underlying maternal condition.

Data

Published clinical lactation studies report that metformin is present in human milk which resulted in infant doses approximately 0.11% to 1% of the maternal weight-adjusted dosage and a milk/plasma ratio ranging between 0.13 and 1. However, the studies were not designed to
definitely establish the risk of use of metformin during lactation because of small sample size and limited adverse event data collected in infants.

8.3 FEMALES & MALES OF REPRODUCTIVE POTENTIAL SECTION

Discuss the potential for unintended pregnancy with premenopausal women as therapy with metformin hydrochloride tablets may result in ovulation in some anovulatory women.

8.4 Pediatric Use

The safety and effectiveness of metformin hydrochloride tabletsfor the treatment of type 2 diabetes mellitus have been established in pediatric patients 10 to 16 years old. Safety and effectiveness of metformin hydrochloride tabletshave not been established in pediatric patients less than 10 years old.
Use of metformin hydrochloride tablets in pediatric patients 10 to 16 years old for the treatment of type 2 diabetes mellitus is supported by evidence from adequate and well-controlled studies of metformin hydrochloride tablets in adults with additional data from a controlled clinical study in pediatric patients 10 to 16 years old with type 2 diabetes mellitus, which demonstrated a similar response in glycemic control to that seen in adults *[see Clinical Studies (14.1)]* . In this study, adverse reactions were similar to those described in adults. A maximum daily dose of 2000 mg of metformin hydrochloride tablets is recommended. *[See Dosage and Administration (2.2).]*

8.5 Geriatric Use

Controlled clinical studies of metformin hydrochloride tablets did not include sufficient numbers of elderly patients to determine whether they respond differently from younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy and the higher risk of lactic acidosis.
Assess renal function more frequently in elderly patients *[see Warnings and Precautions (5.1)]* .

8.6 RENAL IMPAIRMENT

Metformin is substantially excreted by the kidney, and the risk of metformin accumulation and lactic acidosis increases with the degree of renal impairment. Metformin hydrochloride tablets are contraindicated in severe renal impairment, patients with an estimated glomerular filtration rate (eGFR) below 30 mL/min/1.73 m^2 *[see Dosage and Administration (2.3) , Contraindications (4) , Warnings and Precautions (5.1) , and Clinical Pharmacology (12.3)*].

8.7 HEPATIC IMPAIRMENT

Use of metformin in patients with hepatic impairment has been associated with some

cases of lactic acidosis. Metformin hydrochloride tablets are not recommended in patients with hepatic impairment. *[see Warnings and Precautions (5.1)].*

10 OVERDOSAGE

Overdose of metformin hydrochloride has occurred, including ingestion of amounts greater than 50 grams. Hypoglycemia was reported in approximately 10% of cases, but no causal association with metformin has been established. Lactic acidosis has been reported in approximately 32% of metformin overdose cases *[see Warnings and Precautions (5.1)]* . Metformin is dialyzable with a clearance of up to 170 mL/min under good hemodynamic conditions. Therefore, hemodialysis may be useful for removal of accumulated drug from patients in whom metformin overdosage is suspected.

11 DESCRIPTION

Metformin Hydrochloride Tablets, USP contain the antihyperglycemic agent metformin, which is a biguanide, in the form of monohydrochloride. The chemical name of metformin hydrochloride is *N, N*-dimethylimidodicarbonimidic diamide hydrochloride. The structural formula is as shown below:



Metformin hydrochloride is a white to off-white crystalline compound with a molecular formula of $C_4H_{11}N_5$ • HCl and a molecular weight of 165.63. It is freely soluble in water and is practically insoluble in acetone, ether, and chloroform. The pK_a of metformin is 12.4. The pH of a 1% aqueous solution of metformin hydrochloride is 6.68.

Metformin Hydrochloride Tablets, USP contain 500 mg, 850 mg or 1000 mg of metformin hydrochloride, which is equivalent to 389.93 mg, 662.88 mg 779.86 mg metformin base, respectively.

Each tablet contains the inactive ingredients magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc and titanium dioxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Metformin is an antihyperglycemic agent which improves glucose tolerance in patients with type 2 diabetes mellitus, lowering both basal and postprandial plasma glucose. Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing peripheral glucose uptake and utilization. With metformin therapy, insulin secretion remains unchanged while fasting insulin levels and day-long plasma insulin response may decrease.

12.3 Pharmacokinetics

Absorption

The absolute bioavailability of a metformin hydrochloride 500 mg tablet given under fasting conditions is approximately 50% to 60%. Studies using single oral doses of metformin hydrochloride tablets 500 to 1500 mg, and 850 to 2550 mg, indicate that there is a lack of dose proportionality with increasing doses, which is due to decreased absorption rather than an alteration in elimination. At usual clinical doses and dosing schedules of metformin hydrochloride tablets, steady state plasma concentrations of metformin are reached within 24 to 48 hours and are generally <1 µg/mL.

Following a single oral dose of metformin hydrochloride extended-release tablets, C_{max} is achieved with a median value of 7 hours and a range of 4 to 8 hours. Peak plasma levels are approximately 20% lower compared to the same dose of metformin hydrochloride tablets, however, the extent of absorption (as measured by AUC) is comparable to metformin hydrochloride tablets.

At steady state, the AUC and C_{max} are less than dose proportional for metformin hydrochloride extended-release tablets within the range of 500 mg to 2000 mg administered once daily. Peak plasma levels are approximately 0.6, 1.1, 1.4, and 1.8 mcg/mL for 500, 1000, 1500, and 2000 mg once-daily doses, respectively. The extent of metformin absorption (as measured by AUC) from metformin hydrochloride extended-release tablets at a 2000 mg once-daily dose is similar to the same total daily dose administered as metformin hydrochloride tablets 1000 mg twice daily. After repeated administration of metformin hydrochloride extended-release tablets, metformin did not accumulate in plasma.

Effect of food: Food decreases the extent of absorption and slightly delays the absorption of metformin, as shown by approximately a 40% lower mean peak plasma concentration (C_{max}), a 25% lower area under the plasma concentration versus time curve (AUC), and a 35-minute prolongation of time to peak plasma concentration (T_{max}) following administration of a single 850 mg tablet of metformin hydrochloride with food, compared to the same tablet strength administered fasting.

Distribution

The apparent volume of distribution (V/F) of metformin following single oral doses of metformin hydrochloride tablets 850 mg averaged 654 ± 358 L. Metformin is negligibly bound to plasma proteins, Metformin partitions into erythrocytes, most likely as a function of time.

Metabolism

Intravenous single-dose studies in normal subjects demonstrate that metformin is excreted unchanged in the urine and does not undergo hepatic metabolism (no metabolites have been identified in humans) nor biliary excretion.

Elimination

Renal clearance (see Table 3) is approximately 3.5 times greater than creatinine clearance, which indicates that tubular secretion is the major route of metformin elimination. Following oral administration, approximately 90% of the absorbed drug is eliminated via the renal route within the first 24 hours, with a plasma elimination half-life of approximately 6.2 hours. In blood, the elimination half-life is approximately 17.6 hours, suggesting that the erythrocyte mass may be a compartment of distribution.

Special Populations

Renal Impairment

In patients with decreased renal function, the plasma and blood half-life of metformin is prolonged, and the renal clearance is decreased (see Table 3) [See *Dosage and Administration (2.3) , Contraindications (4) , Warnings and Precautions (5.1) and Use in Specific Populations (8.6)*].

Hepatic Impairment

No pharmacokinetic studies of metformin have been conducted in patients with hepatic impairment *[See Warnings and Precautions (5.1) and Use in Specific Populations (8.7)*].

Geriatrics

Limited data from controlled pharmacokinetic studies of metformin hydrochloride tablets in healthy elderly subjects suggest that total plasma clearance of metformin is decreased, the half-life is prolonged, and C_{max} is increased, compared to healthy young subjects. it appears that the change in metformin pharmacokinetics with aging is primarily accounted for by a change in renal function (see Table 3). [*See Warnings and Precautions (5.1)*and *Use in Specific Populations (8.5)*].

Table 3: Select Mean (±S.D.) Metformin Pharmacokinetic Parameters Following Single or Multiple Oral Doses of Metformin Hydrochloride Tablets

Subject Groups: Metformin Hydrochloride Tablets dose[a] (number of subjects)	**C_{max}[b] (mcg/mL)**	**T_{max}[c] (hrs)**	**Renal Clearance (mL/min)**
Healthy, nondiabetic adults:			
500 mg single dose (24)	1.03 (±0.33)	2.75 (±0.81)	600 (±132)
850 mg single dose (74)[d]	1.60 (±0.38)	2.64 (±0.82)	552 (±139)
850 mg three times daily for 19 doses[e] (9)	2.01 (±0.42)	1.79 (±0.94)	642 (±173)
Adults with type 2 diabetes mellitus:			
850 mg single dose (23)	1.48 (±0.5)	3.32 (±1.08)	491 (±138)
850 mg three times daily for 19 doses[e] (9)	1.90 (±0.62)	2.01 (±1.22)	550 (±160)
Elderly[f], healthy nondiabetic adults:			

850 mg single dose (12)	2.45 (±0.70)	2.71 (±1.05)	412 (±98)
Renal-impaired adults: **850 mg single dose**			
Mild (CL_{cr}[g] 61-90 mL/min) (5)	1.86 (±0.52)	3.20 (±0.45)	384 (±122)
Moderate (CL_{cr} 31-60 mL/min) (4)	4.12 (±1.83)	3.75 (±0.50)	108(±57)
Severe (CL_{cr} 10-30 mL/min) (6)	3.93 (±0.92)	4.01 (±1.10)	130 (±90)

[a]All doses given fasting except the first 18 doses of the multiple dose studies

[b]Peak plasma concentration

[c]Time to peak plasma concentration

[d]Combined results (average means) of five studies: mean age 32 years (range 23-59 years)

[e]Kinetic study done following dose 19, given fasting

[f]Elderly subjects, mean age 71 years (range 65-81 years)

[g]CL_{cr}= creatinine clearance normalized to body surface area of 1.73 m^2.

Pediatrics

After administration of a single oral metformin hydrochloride 500 mg tablet with food, geometric mean metformin C_{max} and AUC differed less than 5% between pediatric type 2 diabetic patients (12 to 16 years of age) and gender- and weight-matched healthy adults (20 to 45 years of age), all with normal renal function.

Gender

Metformin pharmacokinetic parameters did not differ significantly between normal subjects and patients with type 2 diabetes mellitus when analyzed according to gender (males = 19, females = 16).

Race

No studies of metformin pharmacokinetic parameters according to race have been performed.

Drug Interactions

In Vivo *Assessment of Drug Interactions*

Table 4: Effect of Coadministered Drug on Plasma Metformin Systemic

Exposure

Coadministered Drug	Dose of Coadministered Drug*	Dose of Metformin*	Geometric Mean Ratio (ratio with/without coadministered drug) No Effect = 1.00		
				AUC†	Cmax
No dosing adjustments required for the following:					
Glyburide	5 mg	850 mg	metformin	0.91‡	0.93‡
Furosemide	40 mg	850 mg	metformin	1.09‡	1.22‡
Nifedipine	10 mg	850 mg	metformin	1.16	1.21
Propranolol	40 mg	850 mg	metformin	0.90	0.94
Ibuprofen	400 mg	850 mg	metformin	1.05‡	1.07‡
Cationic drugs eliminated by renal tubular secretion may reduce metformin elimination [See *Warnings and Precautions (5)* and *Drug Interactions (7).*]					
Cimetidine	400 mg	850 mg	metformin	1.40	1.61
Carbonic anhydrase inhibitors may cause metabolic acidosis [See *Warnings and Precautions (5.1)* and *Drug Interactions (7).*]					
Topiramate	100 mg§	500 mg§	metformin	1.25§	1.17

* All metformin and coadministered drugs were given as single doses

† AUC = AUC(INF)

‡ Ratio of arithmetic means

§ At steady state with topiramate 100 mg every 12 hours and metformin 500 mg every 12 hours; AUC = AUC_{0-12h}

Table 5: Effect of Metformin on Coadministered Drug Systemic Exposure

Coadministered Drug	Dose of Coadministered Drug*	Dose of Metformin*	Geometric Mean Ratio (ratio with/without metformin) No Effect = 1.00		
				AUC†	Cmax
No dosing adjustments required for the following:					
Glyburide	5 mg	850 mg	glyburide	0.78‡	0.63‡
Furosemide	40 mg	850 mg	furosemide	0.87‡	0.69‡
Nifedipine	10 mg	850 mg	nifedipine	1.10§	1.08
Propranolol	40 mg	850 mg	propranolol	1.01§	1.02
Ibuprofen	400 mg	850 mg	ibuprofen	0.97¶	1.01¶
Cimetidine	400 mg	850 mg	cimetidine	0.95§	1.01

* All metformin and coadministered drugs were given as single doses
† AUC = AUC(INF) unless otherwise noted
‡ Ratio of arithmetic means, p-value of difference <0.05
§ AUC (0-24 hr) reported
¶ Ratio of arithmetic means

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis & Mutagenesis & Impairment Of Fertility

Long-term carcinogenicity studies have been performed in rats (dosing duration of 104 weeks) and mice (dosing duration of 91 weeks) at doses up to and including 900 mg/kg/day and 1500 mg/kg/day, respectively. These doses are both approximately 3 times the maximum recommended human daily dose of 2550 mg based on body surface area comparisons. No evidence of carcinogenicity with metformin was found in either male or female mice. Similarly, there was no tumorigenic potential observed with metformin in male rats. There was, however, an increased incidence of benign stromal uterine polyps in female rats treated with 900 mg/kg/day.

There was no evidence of a mutagenic potential of metformin in the following *in vitro*

tests: Ames test (*S. typhimurium),* gene mutation test (mouse lymphoma cells), or chromosomal aberrations test (human lymphocytes). Results in the *in vivo* mouse micronucleus test were also negative.

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately 2 times the maximum recommended human daily dose of 2550 mg based on body surface area comparisons.

14 CLINICAL STUDIES

14.1 Metformin Hydrochloride Tablets

Adult Clinical Studies

A double-blind, placebo-controlled, multicenter US clinical trial involving obese patients with type 2 diabetes mellitus whose hyperglycemia was not adequately controlled with dietary management alone (baseline fasting plasma glucose [FPG] of approximately 240 mg/dL) was conducted. Patients were treated with metformin hydrochloride tablets (up to 2550 mg/day) or placebo for 29 weeks. The results are presented in Table 6.

Table 6: Mean Change in Fasting Plasma Glucose and HbA1c at Week 29 Comparing Metformin Hydrochloride Tablets vs Placebo in Patients with Type 2 Diabetes Mellitus

	Metformin Hydrochloride Tablets(n=141)	**Placebo (n=145)**	**p-Value**
FPG (mg/dL)			
Baseline	241.5	237.7	NS*
Change at FINAL VISIT	–53.0	6.3	0.001
Hemoglobin A1c (%)			
Baseline	8.4	8.2	NS*
Change at FINAL VISIT	–1.4	0.4	0.001

*Not statistically significant

Mean baseline body weight was 201 lbs and 206 lbs in the metformin hydrochloride tablets and placebo arms, respectively. Mean change in body weight from baseline to week 29 was -1.4 lbs and -2.4 lbs in the metformin hydrochloride tablets and placebo arms, respectively. A 29-week, double-blind, placebo-controlled study of metformin hydrochloride tablets and glyburide, alone and in combination, was conducted in obese patients with type 2 diabetes mellitus who had failed to achieve adequate glycemic control while on maximum doses of glyburide (baseline FPG of approximately 250 mg/dL). Patients randomized to the combination arm started therapy with metformin hydrochloride tablets 500 mg and glyburide 20 mg. At the end of each week of the first 4 weeks of the trial, these patients had their dosages of metformin hydrochloride tablets increased by 500 mg if they had failed to reach target fasting plasma glucose. After week 4, such dosage adjustments were made monthly, although no patient was allowed to exceed metformin hydrochloride tablets 2500 mg. Patients in the metformin hydrochloride tablets only arm (metformin plus placebo) discontinued glyburide and followed the same titration schedule. Patients in the glyburide arm continued the same dose of glyburide. At the end of the trial, approximately 70% of the patients in the combination group were taking metformin hydrochloride tablets 2000 mg/glyburide 20 mg or metformin hydrochloride tablets 2500 mg/glyburide 20 mg. The results are displayed in Table 7.

Table 7: Mean Change in Fasting Plasma Glucose and HbA1c at Week 29 Comparing Metformin Hydrochloride Tablets/Glyburide (Comb) vs Glyburide (Glyb) vs Metformin Hydrochloride Tablets(GLU): in Patients with Type 2 Diabetes Mellitus with Inadequate Glycemic Control on Glyburide

	Comb (n=213)	Glyb (n=209)	GLU (n=210)	p-Values		
				Glyb vs Comb	GLU vs Comb	GLU vs Glyb
Fasting Plasma Glucose (mg/dL) Baseline	250.5	247.5	253.9	NS*	NS*	NS*
Change at FINAL VISIT	–63.5	13.7	–0.9	0.001	0.001	0.025
Hemoglobin A1c (%) Baseline	8.8	8.5	8.9	NS*	NS*	0.007
Change at FINAL VISIT	–1.7	0.2	–0.4	0.001	0.001	0.001

* Not statistically significant

Mean baseline body weight was 202 lbs, 203 lbs, and 204 lbs in the metformin hydrochloride tablets /glyburide, glyburide, and metformin hydrochloride tablets arms, respectively. Mean change in body weight from baseline to week 29 was 0.9 lbs, -0.7 lbs, and -8.4 lbs in the metformin hydrochloride tablets /glyburide, glyburide, and metformin hydrochloride tabletsarms, respectively.

Pediatric Clinical Studies

A double-blind, placebo-controlled study in pediatric patients aged 10 to 16 years with type 2 diabetes mellitus (mean FPG 182.2 mg/dL), treatment with metformin hydrochloride tablets (up to 2000 mg/day) for up to 16 weeks (mean duration of treatment 11 weeks) was conducted. The results are displayed in Table 8.

Table 8: Mean Change in Fasting Plasma Glucose at Week 16 Comparing Metformin Hydrochloride Tablets vs Placebo in Pediatric Patients[a] with Type 2 Diabetes Mellitus

	Metformin Hydrochloride Tablets	Placebo	p-Value
FPG (mg/dL)	**(n=37)**	**(n=36)**	
Baseline	162.4	192.3	
Change at FINAL VISIT	–42.9	21.4	<0.001

[a] Pediatric patient mean age 13.8 years (range 10-16 years)

Mean baseline body weight was 205 lbs and 189 lbs in the metformin hydrochloride tablets and placebo arms, respectively. Mean change in body weight from baseline to week 16 was -3.3 lbs and -2.0 lbs in the metformin hydrochloride tablets and placebo arms, respectively.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Metformin Hydrochloride Tablets, USP:

500 mg	Bottles of 30	NDC 25000-134-03
500 mg	Bottles of 1000	NDC 25000-134-14
500 mg	Bulk (16455 tablets)	NDC 25000-134-98
850 mg	Bottles of 30	NDC 25000-131-03
850 mg	Bottles of 500	NDC 25000-131-12
850 mg	Bulk (9680 tablets)	NDC 25000-131-98
1000 mg	Bottles of 30	NDC 25000-132-03
1000 mg	Bottles of 500	NDC 25000-132-12
1000 mg	Bulk (8225 tablets)	NDC 25000-132-98

Metformin Hydrochloride Tablets, USP 500 mg are white to off-white colored, film coated, round shaped tablet with beveled edges on both side and debossed with ‘134’ on one side and plain on other side.

Metformin Hydrochloride Tablets, USP 850 mg are white to off-white colored, film coated, round shaped tablet with beveled edges on both side and debossed with ‘131’ on one side and plain on other side.

Metformin Hydrochloride Tablets, USP 1000 mg are white to off-white colored, film coated, oval shaped tablet with ‘132’ debossed on one side and breakline on both sides.

16.2 Storage

Store at 20°–25° C (68°–77° F); excursions permitted to 15°–30° C (59°–86° F). [See USP Controlled Room Temperature.]

Dispense in light-resistant containers.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).
Lactic Acidosis:
Explain the risks of lactic acidosis, its symptoms, and conditions that predispose to its

development. Advise patients to discontinue metformin hydrochloride tablets immediately and to promptly notify their healthcare provider if unexplained hyperventilation, myalgias, malaise, unusual somnolence or other nonspecific symptoms occur. Counsel patients against excessive alcohol intake and inform patients about importance of regular testing of renal function while receiving metformin hydrochloride tablets Instruct patients to inform their doctor that they are taking metformin hydrochloride tablets prior to any surgical or radiological procedure, as temporary discontinuation may be required *[see Warnings and Precautions (5.1)]*

.

Hypoglycemia

Inform patients that hypoglycemia may occur when metformin hydrochloride tablets are coadministered with oral sulfonylureas and insulin. Explain to patients receiving concomitant therapy the risks of hypoglycemia, its symptoms and treatment, and conditions that predispose to its development *[see Warnings and Precautions (5.3)]*

.

Vitamin B_{12} Deficiency:

Inform patients about importance of regular haematological parameters while receiving metformin hydrochloride tablets

[see Warnings and Precautions (5.2)]

.

Females of Reproductive Age:

Inform females that treatment with metformin hydrochloride tablets

may result in ovulation in some premenopausal anovulatory women which may lead to unintended pregnancy *[see Use in Specific Populations (8.3)] .*

Rx only

Manufactured for:
Time-Cap Labs, Inc.
7 Michael Avenue, Farmingdale,
NY 11735, USA

Manufactured by:
Marksans Pharma Ltd.
Plot No.L-82, L-83, Verna Indl. Estate,
Verna, GOA - 403 722, India

Rev.06/2026

PATIENT MEDICATION INFORMATION SECTION

Metformin Hydrochloride Tablets

(met for' min hye '' droe klor' ide)

Rx only

Read the Patient Information that comes with Metformin Hydrochloride Tablets before you start taking it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your healthcare provider about your medical condition or treatment.

What is the most important information I should know about Metformin Hydrochloride Tablets?

Serious side effects can happen in people taking Metformin Hydrochloride Tablets including:

Lactic Acidosis. Metformin hydrochloride, the medicine in Metformin hydrochloride tablets, can cause a rare, but serious, side effect called lactic acidosis (a build-up of lactic acid in the blood) that can cause death. Lactic acidosis is a medical emergency and must be treated in a hospital.

Stop taking Metformin Hydrochloride Tablets and call your healthcare provider right away if you get any of the following symptoms of lactic acidosis:

- feel very weak and tired
- have unusual (not normal) muscle pain
- have trouble breathing
- have unusual sleepiness or sleep longer than usual
- have unexplained stomach or intestinal problems with nausea and vomiting, or diarrhea
- feel cold, especially in your arms and legs
- feel dizzy or lightheaded
- have a slow or irregular heartbeat

You have a higher chance of getting lactic acidosis if you:

- have kidney problems. People whose kidneys are not working properly should not take metformin hydrochloride tablets,
- have liver problems.
- have congestive heart failure that requires treatment with medicines.
- drink a lot of alcohol (very often or short-term “binge” drinking).
- get dehydrated (lose a large amount of body fluids). This can happen if you are sick with a fever, vomiting, or diarrhea. Dehydration can also happen when you sweat a lot with activity or exercise and do not drink enough fluids.
- have certain x-ray tests with injectable dyes or contrast agents.
- have surgery
- have a heart attack, severe infection, or stroke.
- are 80 years of age or older and have not had your kidney function tested
- have a genetically inherited disease affecting mitochondria (the energy-producing parts within cells) such as mitochondrial encephalomyopathy with lactic acidosis, and stroke-like episodes (MELAS) syndrome and maternally inherited diabetes and deafness (MIDD).

What are Metformin Hydrochloride Tablets?

- Metformin Hydrochloride Tablets are prescription medicines that contain metformin hydrochloride. Metformin Hydrochloride Tablets are used with diet and exercise to help control high blood sugar (hyperglycemia) in adults with type 2 diabetes.
- Metformin Hydrochloride Tablets are not for people with type 1 diabetes.
- Metformin Hydrochloride Tablets are not for people with diabetic ketoacidosis (increased ketones in your blood or urine).

Metformin Hydrochloride Tablets and Metformin Hydrochloride Extended-Release Tablets have the same active ingredient. However, Metformin Hydrochloride Tablets work longer in your body. Both of these medicines help control your blood sugar in a number of ways.

These include helping your body respond better to the insulin it makes naturally, decreasing the amount of sugar your liver makes, and decreasing the amount of sugar your intestines absorb. Metformin hydrochloride tablets do not cause your body to make more insulin.

Who should not take Metformin Hydrochloride Tablets?

Some conditions increase your chance of getting lactic acidosis or cause other problems if you take these medicines. Most of the conditions listed below can increase your chance of getting lactic acidosis.

Do not take Metformin Hydrochloride Tablets if you:

- have kidney problems
- are allergic to the metformin hydrochloride in metformin hydrochloride tabletsor any of the ingredients in metformin hydrochloride tablets. See the end of this leaflet for a complete list of ingredients in metformin hydrochloride tablets.
- are going to get an injection of dye or contrast agents for an x-ray procedure or if you are going to have surgery and not able to eat or drink much. In these situations, metformin hydrochloride tablets will need to be stopped for a short time. Talk to your healthcare provider about when you should stop metformin hydrochloride tabletsand when you should start metformin hydrochloride tabletsagain. See "**What is the most important information I should know about Metformin Hydrochloride Tablets?"**

- **What should I tell my healthcare provider before taking Metformin Hydrochloride Tablets?**

Before taking metformin hydrochloride tablets, tell your healthcare provider if you:

- have type 1 diabetes. Metformin hydrochloride tabletsshould not be used to treat people with type 1 diabetes.
- have a history or risk for diabetic ketoacidosis (high levels of certain acids, known as ketones, in the blood or urine). Metformin hydrochloride tabletsshould not be used for the treatment of diabetic ketoacidosis.
- have kidney problems.
- have liver problems.
- have heart problems, including congestive heart failure.
- are older than 80 years. If you are over 80 years old you should not take metformin hydrochloride tabletsunless your kidneys have been checked and they are normal.
- drink alcohol very often or drink a lot of alcohol in the short-term ("binge" drinking).
- have a genetically inherited disease affecting mitochondria (the energy-producing parts within cells) such as MELAS syndrome and MIDD.
- are taking insulin.
- have any other medical conditions.
- are pregnant or plan to become pregnant. It is not known if metformin hydrochloride tabletswill harm your unborn baby. If you are pregnant, talk with your healthcare provider about the best way to control your blood sugar while you are pregnant.
- are breast-feeding or plan to breast-feed. It is not known if metformin hydrochloride tabletspasses into your breast milk. Talk with your healthcare provider about the best way to feed your baby while you take metformin hydrochloride tablets.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

- Metformin hydrochloride tablets may affect the way other medicines work, and other medicines may affect how metformin hydrochloride tablets works.

Can Metformin Hydrochloride Tablets be used in children?

Metformin hydrochloride tabletshas been shown to effectively lower glucose levels in children (ages 10-16 years) with type 2 diabetes. Metformin hydrochloride tabletshas not been studied in children younger than 10 years old. Metformin hydrochloride tabletshas not been studied in combination with other oral glucose-control medicines or insulin in children. If you have any questions about the use of metformin hydrochloride tabletsin children, talk with your doctor or other healthcare provider.

How should I take Metformin Hydrochloride Tablets?

- Take metformin hydrochloride tabletsexactly as your healthcare provider tells you.
- Metformin hydrochloride tablets should be taken with meals to help lessen an upset stomach side effect.
- Swallow metformin hydrochloride tablets whole.
- You may sometimes pass a soft mass in your stools (bowel movement) that looks like metformin hydrochloride tablets.
- When your body is under some types of stress, such as fever, trauma (such as a car accident), infection, or surgery, the amount of diabetes medicine that you need may change. Tell your healthcare provider right away if you have any of these problems.
- Your healthcare provider should do blood tests to check how well your kidneys are working before and during your treatment with metformin hydrochloride tablets.
- Your healthcare provider will check your diabetes with regular blood tests, including your blood sugar levels and your haemoglobin A1C.
- Follow your healthcare provider's instructions for treating blood sugar that is too low (hypoglycemia). Talk to your healthcare provider if low blood sugar is a problem for you. See **"What are the possible side effects of Metformin Hydrochloride Tablets?"**
- Check your blood sugar as your healthcare provider tells you to.
- Stay on your prescribed diet and exercise program while taking metformin hydrochloride tablets.
- If you miss a dose of metformin hydrochloride tablets, take your next dose as prescribed unless your healthcare provider tells you differently. Do not take an extra dose the next day.
- If you take too much metformin hydrochloride tablets, call your healthcare provider, local Poison Control Center, or go to the nearest hospital emergency room right away.

What should I avoid while taking Metformin Hydrochloride Tablets?

Do not drink a lot of alcoholic drinks while taking metformin hydrochloride tablets. This means you should not binge drink for short periods, and you should not drink a lot of alcohol on a regular basis. Alcohol can increase the chance of getting lactic acidosis.

What are the side effects of Metformin Hydrochloride Tablets?

- **Lactic Acidosis. Metformin, the active ingredient in metformin hydrochloride tablets, can cause a rare but serious condition called lactic acidosis (a buildup of an acid in the blood) that can cause death. Lactic acidosis is a medical emergency and must be treated in the hospital.**

Call your doctor right away if you have any of the following symptoms, which could be signs of lactic acidosis:

- you feel cold in your hands or feet
- you feel dizzy or lightheaded
- you have a slow or irregular heartbeat
- you feel very weak or tired
- you have trouble breathing
- you feel sleepy or drowsy
- you have stomach pains, nausea or vomiting

Most people who have had lactic acidosis with metformin have other things that, combined with the metformin, led to the lactic acidosis. Tell your doctor if you have any of the following, because you have a higher chance for getting lactic acidosis with metformin hydrochloride tabletsif you:

- have severe kidney problems, or your kidneys are affected by certain x-ray tests that use injectable dye
- have liver problems
- drink alcohol very often, or drink a lot of alcohol in short-term "binge" drinking
- get dehydrated (lose a large amount of body fluids). This can happen if you are sick with a fever, vomiting, or diarrhea. Dehydration can also happen when you sweat a lot with activity or exercise and do not drink enough fluids
- have surgery
- have a heart attack, severe infection, or stroke

Common side effects of metformin hydrochloride tabletsinclude diarrhea, nausea, and upset stomach. These side effects generally go away after you take the medicine for a while. Taking your medicine with meals can help reduce these side effects. Tell your doctor if the side effects bother you a lot, last for more than a few weeks, come back after they’ve gone away, or start later in therapy. You may need a lower dose or need to stop taking the medicine for a short period or for good.

About 3 out of every 100 people who take metformin hydrochloride tabletshave an unpleasant metallic taste when they start taking the medicine. It lasts for a short time.

Metformin hydrochloride tabletsrarely cause hypoglycemia (low blood sugar) by themselves. However, hypoglycemia can happen if you do not eat enough, if you drink alcohol, or if you take other medicines to lower blood sugar.

How should I store Metformin Hydrochloride Tablets?

Store Metformin Hydrochloride Tablets at 68°F to 77°F (20°C to 25°C).

Keep Metformin Hydrochloride Tabletsand all medicines out of the reach of children.

General information about the use of Metformin Hydrochloride Tablets

If you have questions or problems, talk with your doctor or other healthcare provider. You can ask your doctor or pharmacist for the information about metformin hydrochloride tabletsthat is written for health care professionals. Medicines are sometimes prescribed for purposes other than those listed in a patient information leaflet. Do not use metformin hydrochloride tabletsfor a condition for which it was not prescribed. Do not share your medicine with other people.

What are the ingredients of Metformin Hydrochloride Tablets?

Active ingredients of metformin hydrochloride tablets: metformin hydrochloride

Metformin hydrochloride tablets, USP contain 500 mg, 850 mg, or 1000 mg of metformin hydrochloride. Each tablet contains the inactive ingredients magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc and titanium dioxide.

What is type 2 diabetes?

Type 2 diabetes is a condition in which your body does not make enough insulin, and the insulin that your body produces does not work as well as it should. Your body can also make too much sugar. When this happens, sugar (glucose) builds up in the blood. This can lead to serious medical problems.

The main goal of treating diabetes is to lower your blood sugar to a normal level.

High blood sugar can be lowered by diet and exercise, and by certain medicines when necessary.

Talk to your healthcare provider about how to prevent, recognize, and take care of low blood sugar (hypoglycemia), high blood sugar (hyperglycemia), and problems you have because of your diabetes.

Manufactured for:

Time-Cap Labs, Inc.

7 Michael Avenue, Farmingdale, NY 11735, USA

Manufactured by:

Marksans Pharma Ltd.

Plot No.L-82, L-83, Verna Indl. Estate,

Verna, GOA - 403 722, India

Rev. 06/26

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Metformin Hydrochloride Tablets USP, 500 mg
30's Container pack

Size : 90 x 35 mm

PANTONE 165 C PANTONE 300 C PANTONE 273 C

PANTONE 327 C Black

Metformin Hydrochloride Tablets USP, 500 mg
1000's Container pack

Size : 140 x 80 mm

PANTONE 165 C PANTONE 300 C PANTONE 273 C

PANTONE 327 C Black

Metformin Hydrochloride Tablets USP, 850 mg

30's Container pack



Metformin Hydrochloride Tablets USP, 850 mg

500's Container pack



Metformin Hydrochloride Tablets USP, 1000 mg
30's Container pack



Metformin Hydrochloride Tablets USP, 1000 mg
500's Container pack



METFORMIN HYDROCHLORIDE

metformin hydrochloride tablet, film coated

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:25000-134
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
METFORMIN HYDROCHLORIDE (UNII: 786Z46389E) (METFORMIN - UNII:9100L32L2N)	METFORMIN HYDROCHLORIDE	500 mg

Inactive Ingredients	
Ingredient Name	**Strength**
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	

Product Characteristics

Color	WHITE (White to off white)	**Score**	no score
Shape	ROUND	**Size**	12mm
Flavor		**Imprint Code**	134
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-134-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2012	
2	NDC:25000-134-14	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2012	
3	NDC:25000-134-51	2600 in 1 BAG; Type 0: Not a Combination Product	06/01/2012	
4	NDC:25000-134-98	1 in 1 BOX	06/01/2012	
4		16455 in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090888	06/01/2012	

METFORMIN HYDROCHLORIDE

metformin hydrochloride tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:25000-131
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METFORMIN HYDROCHLORIDE (UNII: 786Z46389E) (METFORMIN - UNII:9100L32L2N)	METFORMIN HYDROCHLORIDE	850 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	WHITE (white to off-white)	**Score**	no score
Shape	ROUND	**Size**	13mm
Flavor		**Imprint Code**	131
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-131-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2012	
2	NDC:25000-131-12	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2012	
3	NDC:25000-131-52	1800 in 1 BAG; Type 0: Not a Combination Product	06/01/2012	
4	NDC:25000-131-98	1 in 1 BOX	06/01/2012	
4		9680 in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090888	06/01/2012	

METFORMIN HYDROCHLORIDE

metformin hydrochloride tablet, film coated

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:25000-132
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METFORMIN HYDROCHLORIDE (UNII: 786Z46389E) (METFORMIN - UNII:9100L32L2N)	METFORMIN HYDROCHLORIDE	1000 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE (white to off-white)	**Score**	2 pieces
Shape	OVAL	**Size**	19mm
Flavor		**Imprint Code**	132
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:25000-132-03	30 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2012	
2	NDC:25000-132-12	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2012	
3	NDC:25000-132-53	1500 in 1 BAG; Type 0: Not a Combination Product	06/01/2012	
4	NDC:25000-132-98	1 in 1 BOX	06/01/2012	
4		8225 in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090888	06/01/2012	

Labeler - MARKSANS PHARMA LIMITED (925822975)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	MANUFACTURE(25000-131, 25000-132, 25000-134)

Revised: 6/2026 MARKSANS PHARMA LIMITED