

**PAIN AND FEVER CHILDRENS- acetaminophen suspension**  
**P & L Development, LLC**

-----  
**Drug Facts**

**Active ingredient (in each 5 mL)**

**Acetaminophen 160 mg**

**Purpose**

**Pain reliever/fever reducer**

**Uses**

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

**If a skin reaction occurs, stop use and seek medical help right away.**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

**Ask a doctor before use if your child has**

liver disease.

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin.

**When using this product,**

**do not exceed recommended dosage (see overdose warning)**

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not take more than directed (see overdose warning)**
- **shake well before using**
- mL= milliliter
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by a doctor
- find the right dose on chart below. If possible, use weight to dose, use age.

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2	ask a doctor
24-35	2-3	5 mL
36-47	4-5	7.5 ml
48-59	6-8	10 mL
60-71	9-10	12.5 mL
72-95	11	15 mL

\*or as directed by a doctor

### **Other information**

- **each 5 mL contains:** sodium 2 mg
- store between 20-25°C (68-77°F)
- do not refrigerate

### **Inactive ingredients**

acesulfame potassium, citric acid, butylparaben, carboxymethylcellulose sodium, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

### **Questions or Comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

\*\*Compare to active ingredients in Children's Tylenol® Oral Suspension

#### **children's pain relief**

Acetaminophen 160 mg per 5 mL

oral suspension

Pain Reliever/Fever Reducer

alcohol free

aspirin free

ibuprofen free

for ages 2-11 years

cherry flavor

FL OZ (mL)

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

Manufactured by:

PL Developments

11865 S. Alameda St  
Lynwood, CA 90262

## Product Label

Compare to the active ingredient in Children's Tylenol® Oral Suspension\*\*  
NDC 49580-0282-2

**ready in case**

**\$1.25**


**children's pain relief**

**Acetaminophen 160 mg per 5 mL Oral Suspension**

pain reliever/fever reducer

for ages 2-11 years  
alcohol free  
aspirin free  
ibuprofen free

4 fl oz (118 mL)



cherry flavor

**Drug Facts**

**Active ingredient (in each 5 mL)**  
Acetaminophen 160 mg.....Pain reliever/fever reducer

**Purpose**

**Uses** ■ temporarily reduces fever  
■ temporarily relieves minor aches and pains due to ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.


**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

Manufactured by:  
**PL Developments**  
11865 S. Alameda St  
Lynwood, CA 90262

PLD-8231D LB009649



3 49580 28240 7

**Drug Facts (continued)**

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see Overdose warning).**

**Drug Facts (continued under label) PEEL HERE** →

**Drug Facts (continued)**

**Stop use and ask a doctor if** ■ new symptoms occur  
■ redness or swelling is present ■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions** ■ this product does not contain directions or complete warnings for adult use ■ do not take more than directed (see Overdose warning) ■ mL = milliliter ■ shake well before using ■ use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device. ■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give more than 5 days unless directed by a doctor ■ find the right dose on chart. If possible, use weight to dose; otherwise, use age.

**Drug Facts (continued)**

Weight (lb)	Age (year)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

\*or as directed by a doctor

**Other information**  
■ each 5 mL contains: sodium 2 mg ■ store between 20-25°C (68-77°F) ■ do not refrigerate

**Inactive ingredients** acesulfame potassium, butylparaben, carboxymethylcellulose sodium, citric acid, FD&C red #40, flavors, glycerin, high fructose corn syrup, microcrystalline cellulose, propylene glycol, purified water, sodium benzoate, sorbitol, xanthan gum

**Questions or comments?**  
▶ Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

### READYinCASE Children's Pain Relief Cherry Flavor

## PAIN AND FEVER CHILDRENS

acetaminophen suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49580-0282
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	

### Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49580-0282-4	1 in 1 BOX	05/31/2016	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:49580-0282-2	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/31/2016	
3	NDC:49580-0282-8	1 in 1 BOX	05/31/2016	
3		237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/31/2016	

**Labeler** - P & L Development, LLC (101896231)

Revised: 2/2024

P & L Development, LLC