METFORMIN ER 500 MG- metformin er 500 mg tablet
METFORMIN ER 750 MG- metformin er 750 mg tablet
Granules India Ltd

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use METFORMIN HYDROCHLORIDE EXTENDED-RELEASE TABLETS safely and effectively. See full prescribing information for METFORMIN HYDROCHLORIDE EXTENDED-RELEASE TABLETS.

METFORMIN HYDROCHLORIDE extended-release tablets, for oral use
Initial U.S. Approval: 1995

WARNING: LACTIC ACIDOSIS
See full prescribing information for complete boxed warning.

• Postmarketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. Symptoms included malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Laboratory abnormalities included elevated blood lactate levels, anion gap acidosis, increased lactate/pyruvate ratio; and metformin plasma levels generally >5 mcg/mL. (5.1)

• Risk factors include renal impairment, concomitant use of certain drugs, age >65 years old, radiological studies with contrast, surgery and other procedures, hypoxic states, excessive alcohol intake, hepatic impairment, and mitochondrial diseases. Steps to reduce the risk of and manage metformin associated lactic acidosis in these high risk groups are provided in the Full Prescribing Information. (5.1)

• If lactic acidosis is suspected, discontinue Metformin hydrochloride extended-release tablets and institute general supportive measures in a hospital setting. Prompt hemodialysis is recommended. (5.1)

RECENT MAJOR CHANGES

Indications and Usage (1) 02/2026
Boxed Warnings 04/2026

INDICATIONS AND USAGE

Metformin hydrochloride extended-release tablets are biguanide indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus. (1)

DOSAGE AND ADMINISTRATION

Recommended Dosage in Adults:

• Swallow Metformin hydrochloride extended-release tablets whole and never crush, cut or chew (2.2)

• Starting dosage: 500 mg orally once daily with the evening meal (2.2)

• Increase the dosage in increments of 500 mg weekly, up to a maximum dosage of 2,000 mg once daily with the evening meal (2.2)

• Patients receiving Metformin hydrochloride immediate-release tablets may be switched to Metformin hydrochloride extended-release tablets once daily at the same total daily dosage, up to 2,000 mg once daily (2.2)

Recommended Dosage in Patients with Renal Impairment

•Prior to initiation, assess renal function with estimated glomerular filtration rate (eGFR) (2.4)

o Do not use in patients with eGFR below 30 mL/minute/1.73 m^2 (2.4)

o Initiation is not recommended in patients with eGFR between 30 to 45 mL/minute/1.73 m^2 (2.4)

o Assess risk/benefit of continuing if eGFR falls below 45 mL/minute/1.73 m^2 (2.4)

o Discontinue if eGFR falls below 30 mL/minute/1.73 m^2 (2.4)

Discontinuation for Iodinated Contrast Imaging Procedures:

• Metformin hydrochloride extended-release tablets may need to be discontinued at time of, or prior to, iodinated contrast imaging procedures (2.5)

DOSAGE FORMS AND STRENGTHS

- Metformin Hydrochloride Extended-Release Tablets: 500 mg and 750 mg (3)

CONTRAINDICATIONS

• Severe renal impairment (eGFR below 30 mL/min/1.73 m^2) (4, 5.1)

• Hypersensitivity to metformin (4)

• Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma.

WARNINGS AND PRECAUTIONS

- *Lactic Acidosis:* See boxed warning. (5.1)
- *Vitamin B_{12} Deficiency:* Metformin may lower vitamin B_{12} levels. Measure hematological parameters annually and vitamin B_{12} at 2 to 3 year intervals and manage any abnormalities. (5.2)
- *Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues:* Increased risk of hypoglycemia when used in combination with insulin and/or an insulin secretagogue. Lower dose of insulin or insulin secretagogue may be required (5.3)

ADVERSE REACTIONS

For Metformin hydrochloride extended-release tablets, the most common adverse reactions (>5%) are diarrhea, nausea/vomiting, flatulence, asthenia, indigestion, abdominal discomfort, and headache. (6.1) (6)

To report SUSPECTED ADVERSE REACTIONS, contact Granules Pharmaceuticals Inc. at 1-877-770-3183 or FDA at 1-800-FDA-1088 or *www.fda.gov/medwatch* (6)

DRUG INTERACTIONS

• Carbonic anhydrase inhibitors may increase risk of lactic acidosis. Consider more frequent monitoring (7)

• Drugs that reduce metformin clearance (such as ranolazine, vandetanib, dolutegravir, and cimetidine) may increase the accumulation of metformin. Consider the benefits and risks of concomitant use (7)

• Alcohol can potentiate the effect of metformin on lactate metabolism. Warn patients against excessive alcohol intake

USE IN SPECIFIC POPULATIONS

• Females and Males of Reproductive Potential: Advise premenopausal females of the potential for an unintended pregnancy. (8.3)

• Geriatric Use: Assess renal function more frequently. (8.5)

• Hepatic Impairment: Avoid use in patients with hepatic impairment. (8.7)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 7/2026

FULL PRESCRIBING INFORMATION: CONTENTS*

BOXED WARNING
RECENT MAJOR CHANGES
1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.2 Recommended Dosage for Metformin Hydrochloride Extended-Release Tablets in Adults With Type 2 Diabetes Mellitus
2.3 Switching Between Metformin Hydrochloride Immediate-Release Tablets and Metformin Hydrochloride Extended-Release Tablets
2.4 Recommended Dosage in Patients With Renal Impairment
2.5 Discontinuation for Iodinated Contrast Imaging Procedures
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Lactic Acidosis
5.2 Vitamin B_{12} Deficiency
5.3 Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues
5.4 Macrovascular Outcomes
6 ADVERSE REACTIONS
6.1 Clinical Studies Experience
6.2 Postmarketing Experience
7 DRUG INTERACTIONS
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy

8.2 Lactation
8.3 Females and Males of Reproductive Potential
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Renal Impairment
8.7 Hepatic Impairment
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.3 Pharmacokinetics
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
14 CLINICAL STUDIES
14.2 Metformin Hydrochloride Extended-Release Tablets
16 HOW SUPPLIED/STORAGE AND HANDLING
16.1 How Supplied
16.2 Storage
17 PATIENT COUNSELING INFORMATION
* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

BOXED WARNING

WARNING: LACTIC ACIDOSIS

Postmarketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. The onset of metformin associated lactic acidosis is often subtle, accompanied only by nonspecific symptoms such as malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Metformin associated lactic acidosis was characterized by elevated blood lactate levels (>5 mmol/Liter), anion gap acidosis (without evidence of ketonuria or ketonemia), an increased lactate/pyruvate ratio; and metformin plasma levels generally >5 mcg/mL [see*Warnings and Precautions (5.1)*].

Risk factors for metformin-associated lactic acidosis include renal impairment, concomitant use of certain drugs (e.g. carbonic anhydrase inhibitors such as topiramate), age 65 years old or greater, having a radiological study with contrast, surgery and other procedures, hypoxic states (e.g., acute congestive heart failure), excessive alcohol intake, hepatic impairment, and mitochondrial diseases *[see Warnings and Precautions (5.1)]*.

Steps to reduce the risk of and manage metformin-associated lactic acidosis in these high risk groups are provided [see *Dosage and Administration (2), Contraindications (4), Warnings and Precautions (5.1)*].

If metformin-associated lactic acidosis is suspected, immediately discontinue metformin hydrochloride extended-release tablets and institute general supportive measures in a hospital setting. Prompt hemodialysis is recommended [see *Warnings and Precautions (5.1)*].

1 INDICATIONS AND USAGE

Metformin hydrochloride extended-release tablets are indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus.

2 DOSAGE AND ADMINISTRATION

NA

2.2 Recommended Dosage for Metformin Hydrochloride Extended-Release Tablets in Adults With Type 2 Diabetes Mellitus

• Swallow metformin hydrochloride extended-release tablets whole and never crush, cut or chew.
• The recommended starting dosage of metformin hydrochloride extended-release is 500 mg orally once daily with the evening meal.
• Increase the dosage in increments of 500 mg weekly on the basis of glycemic control and tolerability, up to a maximum of 2,000 mg once daily with the evening meal.
• If glycemic control is not achieved with metformin hydrochloride extended-release tablets 2,000 mg once daily, consider a trial of metformin hydrochloride extended-release tablets 1,000 mg twice daily.
• If a dose of metformin hydrochloride extended-release tablets is missed, instruct patients not to take two doses the same day and to resume their usual dose of metformin hydrochloride extended-release tablets with the next schedule dose.

2.3 Switching Between Metformin Hydrochloride Immediate-Release Tablets and Metformin Hydrochloride Extended-Release Tablets

• Patients may switch between metformin hydrochloride immediate-release tablets and metformin hydrochloride extended-release tablets at the same total daily dosage.
• Patients receiving metformin hydrochloride extended-release tablets who require dosages higher than 2,000 mg daily may switch to metformin hydrochloride immediate-release tablets and titrate to the maximum total daily dosage of 2,550 mg.

2.4 Recommended Dosage in Patients With Renal Impairment

• Assess renal function prior to initiation of metformin hydrochloride extended-release tablets and periodically thereafter.
• Metformin hydrochloride extended-release tablets are contraindicated in patients with an estimated glomerular filtration rate (eGFR) below 30 mL/minute/1.73 m^2.
• Initiation of metformin hydrochloride extended-release tablets in patients with an eGFR between 30 to 45 mL/minute/1.73 m^2 is not recommended.
• In patients taking metformin hydrochloride extended-release tablets whose eGFR later falls below 45 mL/min/1.73 m2, assess the benefit risk of continuing therapy.
• Discontinue metformin hydrochloride extended-release tablets if the patient's eGFR later falls below 30 mL/minute/1.73 m^2 [see *Warnings and Precautions (5.1)*].

2.5 Discontinuation for Iodinated Contrast Imaging Procedures

Discontinue metformin hydrochloride extended-release tablets at the time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR between 30 and 60 mL/min/1.73 m^2; in patients with a history of liver disease, alcoholism, or heart failure; or in patients who will be administered intra-arterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure; restart metformin hydrochloride extended-release tablets if renal function is stable.

3 DOSAGE FORMS AND STRENGTHS

Metformin hydrochloride extended-release tablets, USP are available as:
• *Extended-release tablets: 500 mg* white to off-white uncoated, modified capsule

shaped tablets debossed with "G7" on one side and plain on other side.

• *Extended-release tablets: 750 mg* white to off white uncoated, modified capsule shaped tablets debossed with "G8" on one side and plain on other side.

4 CONTRAINDICATIONS

Metformin hydrochloride extended-release tablets, USP are contraindicated in patients with:

- Severe renal impairment (eGFR below 30 mL/min/1.73 m ^{2}) [see *Warnings and Precautions (5.1)].*
- Hypersensitivity to metformin.
- Acute or chronic metabolic acidosis, including diabetic ketoacidosis, with or without coma.

5 WARNINGS AND PRECAUTIONS

NA

5.1 Lactic Acidosis

There have been postmarketing cases of metformin-associated lactic acidosis, including fatal cases. These cases had a subtle onset and were accompanied by nonspecific symptoms such as malaise, myalgias, abdominal pain, respiratory distress, or increased somnolence; however, hypotension and resistant bradyarrhythmias have occurred with severe acidosis. Metformin-associated lactic acidosis was characterized by elevated blood lactate concentrations (>5 mmol/L), anion gap acidosis (without evidence of ketonuria or ketonemia), and an increased lactate: pyruvate ratio; metformin plasma levels were generally >5 mcg/mL. Metformin decreases liver uptake of lactate increasing lactate blood levels which may increase the risk of lactic acidosis, especially in patients at risk.

If metformin-associated lactic acidosis is suspected, general supportive measures should be instituted promptly in a hospital setting, along with immediate discontinuation of metformin hydrochloride extended-release tablets. In metformin hydrochloride extended-release tablets treated patients with a diagnosis or strong suspicion of lactic acidosis, prompt hemodialysis is recommended to correct the acidosis and remove accumulated metformin (metformin hydrochloride is dialyzable with a clearance of up to 170 mL/min under good hemodynamic conditions). Hemodialysis has often resulted in reversal of symptoms and recovery.

Educate patients and their families about the symptoms of lactic acidosis and, if these symptoms occur, instruct them to discontinue metformin hydrochloride extended-release tablets and report these symptoms to their healthcare provider.

For each of the known and possible risk factors for metformin-associated lactic acidosis, recommendations to reduce the risk of and manage metformin-associated lactic acidosis are provided below:
• *Renal impairment* — The postmarketing metformin-associated lactic acidosis cases primarily occurred in patients with significant renal impairment.
The risk of metformin accumulation and metformin-associated lactic acidosis increases with the severity of renal impairment because metformin is substantially excreted by the kidney. Clinical recommendations based upon the patient's renal function include [see *Dosage and Administration (2.2), Clinical Pharmacology (12.3)*]:
o Before initiating metformin hydrochloride extended-release tablets, obtain an estimated glomerular filtration rate (eGFR).
o Metformin hydrochloride extended-release tablets are contraindicated in patients with

an eGFR less than 30 mL/min/1.73 m ^{2} [see *Contraindications (4)*].
o Initiation of metformin hydrochloride extended-release tablets are not recommended in patients with eGFR between 30 to 45 mL/min/1.73 m ^{2}.
o Obtain an eGFR at least annually in all patients taking metformin hydrochloride extended-release tablets. In patients at risk for the development of renal impairment (e.g., the elderly), renal function should be assessed more frequently.
o In patients taking metformin hydrochloride extended-release tablets whose eGFR falls below 45 mL/min/1.73 m ^{2}, assess the benefit and risk of continuing therapy.
• *Drug interactions* — The concomitant use of metformin hydrochloride extended-release tablets with specific drugs may increase the risk of metformin-associated lactic acidosis: those that impair renal function, result in significant hemodynamic change, interfere with acid-base balance, or increase metformin accumulation. Consider more frequent monitoring of patients.
• *Age 65 or greater* — The risk of metformin-associated lactic acidosis increases with the patient's age because elderly patients have a greater likelihood of having hepatic, renal, or cardiac impairment than younger patients. Assess renal function more frequently in elderly patients.
• *Radiologic studies with contrast* — Administration of intravascular iodinated contrast agents in metformin-treated patients has led to an acute decrease in renal function and the occurrence of lactic acidosis. Stop metformin hydrochloride extended-release tablets at the time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR between 30 and 60 mL/min/1.73 m ^{2}; in patients with a history of hepatic impairment, alcoholism or heart failure; or in patients who will be administered intra-arterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure, and restart metformin hydrochloride extended-release tablets if renal function is stable.
• *Surgery and other procedures* — Withholding of food and fluids during surgical or other procedures may increase the risk for volume depletion, hypotension, and renal impairment. Metformin hydrochloride extended-release tablets should be temporarily discontinued while patients have restricted food and fluid intake.
• *Hypoxic states* — Several of the postmarketing cases of metformin-associated lactic acidosis occurred in the setting of acute congestive heart failure (particularly when accompanied by hypoperfusion and hypoxemia). Cardiovascular collapse (shock), acute myocardial infarction, sepsis, and other conditions associated with hypoxemia have been associated with lactic acidosis and may cause prerenal azotemia. When such an event occurs, discontinue metformin hydrochloride extended-release tablets.
• *Excessive alcohol intake* — Alcohol potentiates the effect of metformin on lactate metabolism. Patients should be warned against excessive alcohol intake while receiving metformin hydrochloride extended-release tablets.
• *Hepatic impairment* — Patients with hepatic impairment have developed cases of metformin-associated lactic acidosis. This may be due to impaired lactate clearance resulting in higher lactate blood levels. Therefore, avoid use of metformin hydrochloride extended-released tablets in patients with clinical or laboratory evidence of hepatic disease.

• *Mitochondrial Diseases* - Metformin may increase the risk of lactic acidosis in patients with mitochondrial diseases such as mitochondrial encephalomyopathy with lactic acidosis, and stroke-like episodes (MELAS) syndrome and maternally inherited diabetes and deafness (MIDD). Metformin hydrochloride extended-released tablets are not recommended in
patients with mitochondrial diseases such as MELAS syndrome and MIDD.

If a patient taking metformin hydrochloride extended-released tablets develops elevated lactic acid levels (with or without accompanying neurologic signs and symptoms), discontinue metformin hydrochloride extended-released tablets. Depending on the clinical context, consider evaluating the patient for MELAS, MIDD, or other mitochondrial disease.

5.2 Vitamin B $_{12}$ Deficiency

In metformin hydrochloride tablets clinical trials of 29-week duration, a decrease to subnormal levels of previously normal serum vitamin B $_{12}$ levels was observed in

approximately 7% of patients. Such decrease, possibly due to interference with B_{12} absorption from the B_{12}-intrinsic factor complex, may be associated with anemia but appears to be rapidly reversible with discontinuation of metformin hydrochloride tablets or vitamin B_{12} supplementation. Certain individuals (those with inadequate vitamin B_{12} or calcium intake or absorption) appear to be predisposed to developing subnormal vitamin B_{12} levels. Measure hematologic parameters on an annual basis and vitamin B_{12} at 2 to 3 year intervals in patients on metformin hydrochloride extended-release tablets and manage any abnormalities [see *Adverse Reactions (6.1)*].

5.3 Hypoglycemia with Concomitant Use with Insulin and Insulin Secretagogues

Insulin and insulin secretagogues (e.g., sulfonylurea) are known to cause hypoglycemia. Metformin hydrochloride extended-release tablets may increase the risk of hypoglycemia when combined with insulin and/or an insulin secretagogue. Therefore, a lower dose of insulin or insulin secretagogue may be required to minimize the risk of hypoglycemia when used in combination with metformin hydrochloride extended-release tablets [see *Drug Interactions (7)*].

5.4 Macrovascular Outcomes

There have been no clinical studies establishing conclusive evidence of macrovascular risk reduction with metformin hydrochloride extended-release tablets.

6 ADVERSE REACTIONS

The following adverse reactions are also discussed elsewhere in the labeling:

- Lactic Acidosis [see *Boxed Warning and Warnings and Precautions (5.1)*]
- Vitamin B_{12} Deficiency [see *Warnings and Precautions (5.2)*]
- Hypoglycemia [see *Warnings and Precautions (5.3)*]

6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Metformin Hydrochloride Extended-Release Tablets
In placebo-controlled trials, in adults with type 2 diabetes mellitus, 781 patients were administered metformin hydrochloride extended-release tablets. Adverse reactions reported in greater than 5% of the metformin hydrochloride extended-release tablets patients, and that were more common in metformin hydrochloride extended-release tablets - than placebo-treated patients, are listed in Table 1.

Table 1: Adverse Reactions from Clinical Trials of Metformin Hydrochloride Extended-Release Tablets Occurring >5% and More Common than Placebo in Adults with Type 2 Diabetes Mellitus

	Metformin Hydrochloride Extended- Released Tablets (n=781)	**Placebo (n=195)**
Diarrhea	10%	3%
Nausea/Vomiting	7%	2%

Diarrhea led to discontinuation of metformin hydrochloride extended-release tablets in 0.6% of patients. Additionally, the following adverse reactions were reported in ≥1% to ≤5% of metformin hydrochloride extended-release tablets patients and were more commonly reported with metformin hydrochloride extended-release tablets than placebo: abdominal pain, constipation, distention abdomen, dyspepsia/heartburn, flatulence, dizziness, headache, upper respiratory infection, taste disturbance.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of metformin. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Cholestatic, hepatocellular, and mixed hepatocellular liver injury have been reported with postmarketing use of metformin.

7 DRUG INTERACTIONS

Table 2 presents clinically significant drug interactions with metformin hydrochloride extended-release tablets.

Table 2: Clinically Significant Drug Interactions with Metformin Hydrochloride Extended-Release Tablets

Carbonic Anhydrase Inhibitors	
Clinical Impact:	Carbonic anhydrase inhibitors frequently cause a decrease in serum bicarbonate and induce non-anion gap, hyperchloremic metabolic acidosis. Concomitant use of these drugs with metformin hydrochloride extended-release tablets may increase the risk for lactic acidosis.
Intervention:	Consider more frequent monitoring of these patients.
Examples:	Topiramate, zonisamide, acetazolamide or dichlorphenamide.
Drugs that Reduce Metformin Hydrochloride Extended-release Tablets Clearance	
Clinical Impact:	Concomitant use of drugs that interfere with common renal tubular transport systems involved in the renal elimination of metformin (e.g., organic cationic transporter-2 [OCT2] / multidrug and toxin extrusion [MATE] inhibitors) could increase systemic exposure to metformin and may increase the risk for lactic acidosis *[see Clinical Pharmacology (12.3)]*.
Intervention:	Consider the benefits and risks of concomitant use with metformin hydrochloride extended-release tablets.
Examples:	Ranolazine, vandetanib, dolutegravir, and cimetidine.
Alcohol	
Clinical Impact:	Alcohol is known to potentiate the effect of metformin on lactate metabolism.
Intervention:	Warn patients against excessive alcohol intake while receiving metformin hydrochloride extended-release tablets.
Insulin Secretagogues or Insulin	
Clinical Impact:	Coadministration of metformin hydrochloride extended-release tablets with an insulin secretagogue (e.g., sulfonylurea) or insulin may increase the risk of hypoglycemia.
Intervention:	Patients receiving an insulin secretagogue or insulin may require lower doses of the insulin secretagogue or insulin.
Drugs Affecting Glycemic Control	
Clinical Impact:	Certain drugs tend to produce hyperglycemia and may lead to loss of glycemic control.
Intervention:	When such drugs are administered to a patient receiving metformin hydrochloride extended-release tablets, observe the patient closely for loss of blood glucose control. When such drugs are withdrawn from a patient receiving metformin hydrochloride extended-release tablets, observe the patient closely for hypoglycemia.
Examples:	Thiazides and other diuretics, corticosteroids, phenothiazines, thyroid products, estrogens, oral contraceptives, phenytoin, nicotinic acid, sympathomimetics, calcium channel blockers, and isoniazid.

8 USE IN SPECIFIC POPULATIONS

NA

8.1 Pregnancy

Risk Summary

Limited data with metformin hydrochloride extended-release tablets in pregnant women are not sufficient to determine a drug-associated risk for major birth defects or miscarriage. Published studies with metformin use during pregnancy have not reported a clear association with metformin and major birth defect or miscarriage risk [see *Data*]. There are risks to the mother and fetus associated with poorly controlled diabetes mellitus in pregnancy [see *Clinical Considerations*].

No adverse developmental effects were observed when metformin was administered to pregnant Sprague Dawley rats and rabbits during the period of organogenesis at doses up to 2-and 5-times, respectively, a 2,550 mg clinical dose, based on body surface area [see *Data*].

The estimated background risk of major birth defects is 6 to 10% in women with pre-gestational diabetes mellitus with an HbA1C >7 and has been reported to be as high as 20 to 25% in women with a HbA1C >10. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Clinical Considerations

Disease-associated maternal and/or embryo/fetal risk

Poorly-controlled diabetes mellitus in pregnancy increases the maternal risk for diabetic ketoacidosis, pre-eclampsia, spontaneous abortions, preterm delivery, stillbirth and delivery complications. Poorly controlled diabetes mellitus increases the fetal risk for major birth defects, stillbirth, and macrosomia related morbidity.

Data

Human Data

Published data from post-marketing studies have not reported a clear association with metformin and major birth defects, miscarriage, or adverse maternal or fetal outcomes when metformin was used during pregnancy. However, these studies cannot definitely establish the absence of any metformin-associated risk because of methodological limitations, including small sample size and inconsistent comparator groups.

Animal Data

Metformin hydrochloride did not adversely affect development outcomes when administered to pregnant rats and rabbits at doses up to 600 mg/kg/day. This represents an exposure of about 2 and 5 times a 2,550 mg clinical dose based on body surface area comparisons for rats and rabbits, respectively. Determination of fetal concentrations demonstrated a partial placental barrier to metformin.

8.2 Lactation

Risk Summary

Limited published studies report that metformin is present in human milk [see *Data*]. However, there is insufficient information to determine the effects of metformin on the breastfed infant and no available information on the effects of metformin on milk production. Therefore, the developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for metformin hydrochloride extended-release tablets and any potential adverse effects on the breastfed child from metformin hydrochloride extended-release tablets or from the underlying maternal condition.

Data

Published clinical lactation studies report that metformin is present in human milk which resulted in infant doses approximately 0.11% to 1% of the maternal weight-adjusted dosage and a milk/plasma ratio ranging between 0.13 and 1. However, the studies were not designed to definitely establish the risk of use of metformin during lactation because

of small sample size and limited adverse event data collected in infants.

8.3 Females and Males of Reproductive Potential

Discuss the potential for unintended pregnancy with premenopausal women as therapy with metformin hydrochloride extended-release tablets may result in ovulation in some anovulatory women.

8.4 Pediatric Use

Use of metformin hydrochloride extended-release tablets for this indication is supported by evidence from adequate and well-controlled trials of metformin hydrochloride immediate-release tablets in adults with type 2 diabetes mellitus.

8.5 Geriatric Use

Controlled clinical studies of metformin hydrochloride extended-release tablets did not include sufficient numbers of elderly patients to determine whether they respond differently from younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy and the higher risk of lactic acidosis. Assess renal function more frequently in elderly patients [see *Warnings and Precautions (5.1)*].

8.6 Renal Impairment

Metformin is substantially excreted by the kidney, and the risk of metformin accumulation and lactic acidosis increases with the degree of renal impairment. Metformin hydrochloride extended-release tablets are contraindicated in severe renal impairment, patients with an estimated glomerular filtration rate (eGFR) below 30 mL/min/1.73 m^2 [see *Dosage and Administration (2.4), Contraindications (4), Warnings and Precautions (5.1), and Clinical Pharmacology (12.3)*].

8.7 Hepatic Impairment

Use of metformin in patients with hepatic impairment has been associated with some cases of lactic acidosis. Metformin hydrochloride extended-release tablets are not recommended in patients with hepatic impairment. [see *Warnings and Precautions (5.1)*].

10 OVERDOSAGE

In the event of an overdose with metformin hydrochloride extended-release tablets, consider contacting the Poison Help line (1-800-222-1222) or a medical toxicologist for additional overdosage management recommendations.

Overdose of metformin hydrochloride has occurred, including ingestion of amounts greater than 50 grams. Hypoglycemia was reported in approximately 10% of cases, but no causal association with metformin has been established. Lactic acidosis has been reported in approximately 32% of metformin overdose cases [see *Warnings and Precautions (5.1)]*. Metformin is dialyzable with a clearance of up to 170 mL/min under good hemodynamic conditions. Therefore, hemodialysis may be useful for removal of accumulated drug from patients in whom metformin overdosage is suspected.

11 DESCRIPTION

Metformin hydrochloride extended-release tablets, USP contain the antihyperglycemic agent metformin, which is a biguanide, in the form of monohydrochloride. The chemical name of metformin hydrochloride is N,N-dimethylimidodicarbonimidic diamide hydrochloride . The structural formula is as shown below:



Metformin hydrochloride, USP is a white to off-white crystalline compound with a molecular formula of $C_4H_{11}N_5 \cdot HCl$ and a molecular weight of 165.62. Metformin hydrochloride is freely soluble in water, slightly soluble in ethanol, practically insoluble in acetone and in methylene chloride. The pK_a of metformin is 12.4. The pH of a 1% aqueous solution of metformin hydrochloride is 6.35.

Metformin hydrochloride extended-release tablets USP, contains 500 mg or 750 mg of metformin hydrochloride, which is equivalent to 389.93 mg, 584.9 mg metformin base, respectively.

Metformin hydrochloride extended-release tablets USP, 500 mg tablets contain the inactive ingredients hypromellose, magnesium stearate, and polyvinyl pyrrolidone

Metformin hydrochloride extended-release tablets USP, 750 mg tablets contain the inactive ingredients hypromellose, magnesium stearate, and polyvinyl pyrrolidone

Meets USP Dissolution Test 10

12 CLINICAL PHARMACOLOGY

NA

12.1 Mechanism of Action

Metformin is an antihyperglycemic agent which improves glucose tolerance in patients with type 2 diabetes mellitus, lowering both basal and postprandial plasma glucose. Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing peripheral glucose uptake and utilization. With metformin therapy, insulin secretion remains unchanged while fasting insulin levels and day-long plasma insulin response may decrease.

12.3 Pharmacokinetics

Absorption

The absolute bioavailability of a metformin hydrochloride tablets, 500 mg tablet given under fasting conditions is approximately 50% to 60%. Studies using single oral doses of metformin hydrochloride tablets 500 to 1,500 mg and 850 to 2,550 mg, indicate that there is a lack of dose proportionality with increasing doses, which is due to decreased absorption rather than an alteration in elimination. At usual clinical doses and dosing schedules of metformin hydrochloride tablets, steady state plasma concentrations of metformin are reached within 24 to 48 hours and are generally <1 µg/mL.

Following a single oral dose of metformin hydrochloride extended-release tablets, C_{max} is achieved with a median value of 7 hours and a range of 4 to 8 hours. Peak plasma levels are approximately 20% lower compared to the same dose of metformin hydrochloride tablets, however, the extent of absorption (as measured by AUC) is comparable to metformin hydrochloride tablets.

At steady state, the AUC and C_{max} are less than dose proportional for metformin hydrochloride extended-release tablets within the range of 500 to 2,000 mg administered once daily. Peak plasma levels are approximately 0.6, 1.1, 1.4 and 1.8 mcg/mL for 500, 1,000, 1,500, and 2,000 mg once-daily doses, respectively. The extent

of metformin absorption (as measured by AUC) from metformin hydrochloride extended-release tablets at a 2,000 mg once-daily dose is similar to the same total daily dose administered as metformin hydrochloride tablets 1,000 mg twice daily. After repeated administration of metformin hydrochloride extended-release tablets, metformin did not accumulate in plasma.

Effect of food: Food decreases the extent of absorption and slightly delays the absorption of metformin, as shown by approximately a 40% lower mean peak plasma concentration (C_{max}), a 25% lower area under the plasma concentration versus time curve (AUC), and a 35-minute prolongation of time to peak plasma concentration (T_{max}) following administration of a single 850 mg tablet of metformin hydrochloride tablets with food, compared to the same tablet strength administered fasting.

Although the extent of metformin absorption (as measured by AUC) from the metformin hydrochloride extended-release tablet increased by approximately 50% when given with food, there was no effect of food on C_{max} and T_{max} of metformin. Both high and low fat meals had the same effect on the pharmacokinetics of metformin hydrochloride extended-release tablets.

Distribution
The apparent volume of distribution (V/F) of metformin following single oral doses of metformin hydrochloride tablets 850 mg averaged 654 ± 358 L. Metformin is negligibly bound to plasma proteins. Metformin partitions into erythrocytes, most likely as a function of time.

Metabolism
Intravenous single-dose studies in normal subjects demonstrate that metformin is excreted unchanged in the urine and does not undergo hepatic metabolism (no metabolites have been identified in humans) nor biliary excretion.

Elimination
Renal clearance (see Table 3) is approximately 3.5 times greater than creatinine clearance, which indicates that tubular secretion is the major route of metformin elimination. Following oral administration, approximately 90% of the absorbed drug is eliminated via the renal route within the first 24 hours, with a plasma elimination half-life of approximately 6.2 hours. In blood, the elimination half-life is approximately 17.6 hours, suggesting that the erythrocyte mass may be a compartment of distribution.

Specific Populations

Patients With Renal Impairment
In patients with decreased renal function the plasma and blood half-life of metformin is prolonged and the renal clearance is decreased (see Table 3) [see *Dosage and Administration (2.3),Contraindications (4), Warnings and Precautions (5.1) and Use in Specific Populations (8.6)].*

Patients With Hepatic Impairment
No pharmacokinetic studies of metformin have been conducted in patients with hepatic impairment [see *Warnings and Precautions (5.1) and Use in Specific Populations (8.7)]*.

Geriatric Patients
Limited data from controlled pharmacokinetic studies of metformin hydrochloride tablets in healthy elderly subjects suggest that total plasma clearance of metformin is decreased, the half-life is prolonged, and C_{max} is increased, compared to healthy young subjects. It appears that the change in metformin pharmacokinetics with aging is primarily accounted for by a change in renal function (see Table 3). [see *Warnings and Precautions (5.1)and Use in Specific Populations (8.5)].*

Table 3: Select Mean (±S.D.) Metformin Pharmacokinetic Parameters Following Single or Multiple Oral Doses of Metformin Hydrochloride Tablets

Subject Groups: Metformin Hydrochloride	**C [b]**	**T [c]**	**Renal**

Tablets dose [a] (number of subjects)	C_{max} [b] (mcg/mL)	T_{max} [c] (hrs)	Clearance (mL/min)
Healthy, nondiabetic adults:			
500 mg single dose (24)	1.03 (±0.33)	2.75 (±0.81)	600 (±132)
850 mg single dose (74) [d]	1.6 (±0.38)	2.64 (±0.82)	552 (±139)
850 mg three times daily for 19 doses [e] (9)	2.01 (±0.42)	1.79 (±0.94)	642 (±173)
Adults with type 2 diabetes mellitus:			
850 mg single dose (23)	1.48 (±0.5)	3.32 (±1.08)	491 (±138)
850 mg three times daily for 19 doses [e] (9)	1.9 (±0.62)	2.01 (±1.22)	550 (±160)
Elderly[f], healthy nondiabetic adults: 850 mg single dose (12)	2.45 (±0.7)	2.71 (±1.05)	412 (±98)
Renal-impaired adults: **850 mg single dose**	1.86 (±0.52)	3.2 (±0.45)	384 (±122)
Mild (CL_{cr}[g] 61 to 90 mL/min) (5)	4.12 (±1.83)	3.75 (±0.5)	108 (±57)
Moderate (CL_{cr} 31 to 60 mL/min) (4) **Severe** (CL_{cr} 10 to 30 mL/min) (6)	3.93 (±0.92)	4.01 (±1.1)	130 (±90)

[a] All doses given fasting except the first 18 doses of the multiple dose studies
[b] Peak plasma concentration
[c] Time to peak plasma concentration
[d] Combined results (average means) of five studies: mean age 32 years (range 23 to 59 years)
[e] Kinetic study done following dose 19, given fasting
[f] Elderly subjects, mean age 71 years (range 65 to 81 years)
[g] CL_{cr} = creatinine clearance normalized to body surface area of 1.73 m^2

Sex
Metformin pharmacokinetic parameters did not differ significantly between normal subjects and patients with type 2 diabetes mellitus when analyzed according to sex (males=19, females=16).

Race
No studies of metformin pharmacokinetic parameters according to race have been performed.

Drug Interaction Studies
In Vivo Assessment of Drug Interactions
Table 4: Effect of Coadministered Drug on Plasma Metformin Systemic Exposure

Coadministered Drug	Dose of Coadministered Drug*	Dose of Metformin*	Geometric Mean Ratio (ratio with/without coadministered drug) No Effect = 1		
				AUC †	C_{max}
No dosing adjustments required for the following:					
Glyburide	5 mg	850 mg	metformin	0.91 ‡	0.93 ‡
Furosemide	40 mg	850 mg	metformin	1.09 ‡	1.22 ‡
Nifedipine	10 mg	850 mg	metformin	1.16	1.21
Propranolol	40 mg	850 mg	metformin	0.9	0.94
Ibuprofen	400 mg	850 mg	metformin	1.05 ‡	1.07 ‡
Cationic drugs eliminated by renal tubular secretion may reduce metformin elimination [see *Warnings and Precautions (5.9)* and *Drug*					

Interactions (7.2).]					
Cimetidine	400 mg	850 mg	metformin	1.4	1.61
Carbonic anhydrase inhibitors may cause metabolic acidosis [see *Warnings and Precautions (5.1)* and *Drug Interactions (7.1).*]					
Topiramate	100 mg §	500 mg §	metformin	1.25 §	1.17

* All metformin and coadministered drugs were given as single doses

† AUC = AUC (INF)
‡ Ratio of arithmetic means
§At steady state with topiramate 100 mg every 12 hours and metformin 500 mg every 12 hours;
AUC = AUC0-12h

Table 5: Effect of Metformin on Coadministered Drug Systemic Exposure

Coadministered Drug	**Dose of Coadministered Drug***	**Dose of Metformin***	**Geometric Mean Ratio (ratio with/without metformin) No Effect = 1**		
				AUC‡	**C_{max}**
No dosing adjustments required for the following:					
Glyburide	5 mg	850 mg	glyburide	0.78 ‡	0.63 ‡
Furosemide	40 mg	850 mg	furosemide	0.87 ‡	0.69 ‡
Nifedipine	10 mg	850 mg	nifedipine	1.1 §	1.08
Propranolol	40 mg	850 mg	propranolol	1.01 §	1.02
Ibuprofen	400 mg	850 mg	ibuprofen	0.97 ¶	1.01 ¶
Cimetidine	400 mg	850 mg	cimetidine	0.95 §	1.01

* All metformin and coadministered drugs were given as single doses
† AUC = AUC (INF) unless otherwise noted
‡ Ratio of arithmetic means, p-value of difference <0.05
§AUC (0-24 hr) reported
¶ Ratio of arithmetic means

13 NONCLINICAL TOXICOLOGY

NA

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term carcinogenicity studies have been performed in rats (dosing duration of 104 weeks) and mice (dosing duration of 91 weeks) at doses up to and including 900 mg/kg/day and 1,500 mg/kg/day, respectively. These doses are both approximately 3 times the maximum recommended human daily dose of 2,550 mg based on body surface area comparisons. No evidence of carcinogenicity with metformin was found in either male or female mice. Similarly, there was no tumorigenic potential observed with metformin in male rats. There was, however, an increased incidence of benign stromal uterine polyps in female rats treated with 900 mg/kg/day.

There was no evidence of a mutagenic potential of metformin in the following *in vitro* tests: Ames test (*S. typhimurium*), gene mutation test (mouse lymphoma cells), or chromosomal aberrations test (human lymphocytes). Results in the *in vivo* mouse micronucleus test were also negative.

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately 2 times the maximum recommended human daily dose of 2,550 mg based on body surface area comparisons.

14 CLINICAL STUDIES

NA

14.2 Metformin Hydrochloride Extended-Release Tablets

A 24-week, double-blind, placebo-controlled trial of metformin hydrochloride extended-release tablets, taken once daily with the evening meal, was conducted in adults with type 2 diabetes mellitus who had failed to achieve glycemic control with diet and exercise. Patients entering the trial had a mean baseline HbA_{1c} of 8% and a mean baseline FPG of 176 mg/dL. The treatment dose was increased to 1,500 mg once daily if at Week 12 HbA_{1c} was ≥7% but <8% (patients with HbA_{1c} ≥8% were discontinued from the trial). At the final visit (24-week), mean HbA1c had increased 0.2% from baseline in placebo patients and decreased 0.6% with metformin hydrochloride extended-release tablets.

A 16-week, double-blind, placebo-controlled, dose-response trial of metformin hydrochloride extended-release tablets, taken once daily with the evening meal or twice daily with meals, was conducted in adults with type 2 diabetes mellitus who had failed to achieve glycemic control with diet and exercise. The results are shown in Table 6.

Table 6: Mean Changes from Baseline* in HbA1c and Fasting Plasma Glucose at Week 16 Comparing Metformin Extended-Release Tablets vs Placebo in Adults with Type 2 Diabetes Mellitus

	Metformin Hydrochloride Extended-Release Tablets					
	500 mg Once Daily	**1,000 mg Once Daily**	**1,500 mg Once Daily**	**2,000 mg Once Daily**	**1,000 mg Twice Daily**	**Placebo**
Hemoglobin A_{1c} (%)	**(n=115)**	**(n=115)**	**(n=111)**	**(n=125)**	**(n=112)**	**(n=111)**
Baseline	8.2	8.4	8.3	8.4	8.4	8.4
Change at FINAL VISIT	–0.4	–0.6	–0.9	–0.8	–1.1	0.1
p-value [a]	<0.001	<0.001	<0.001	<0.001	<0.001	–
FPG (mg/dL)	**(n=126)**	**(n=118)**	**(n=120)**	**(n=132)**	**(n=122)**	**(n=113)**
Baseline	182.7	183.7	178.9	181	181.6	179.6
Change at FINAL VISIT	–15.2	–19.3	–28.5	–29.9	–33.6	7.6
p-value [a]	<0.001	<0.001	<0.001	<0.001	<0.001	–

[a] All comparisons versus Placebo

Mean baseline body weight was 193 lbs, 192 lbs, 188 lbs, 196 lbs, 193 lbs and 194 lbs in the metformin hydrochloride extended-release tablets 500 mg, 1,000 mg, 1,500 mg, and 2,000 mg once daily, 1,000 mg twice daily and placebo arms, respectively. Mean change in body weight from baseline to week 16 was -1.3 lbs, 1.3 lbs, -0.7 lbs, -1.5 lbs, -2.2 lbs and -1.8 lbs, respectively.

A 24-week, double-blind, randomized trial of metformin hydrochloride extended-release tablets, taken once daily with the evening meal, and metformin hydrochloride tablets, taken twice daily (with breakfast and evening meal), was conducted in adults with type 2 diabetes mellitus who had been treated with metformin hydrochloride tablets 500 mg twice daily for at least 8 weeks prior to study entry. The results are shown in Table 7.

Table 7: Mean Changes from Baseline* in HbA1c and Fasting Plasma Glucose at Week 24 Comparing Metformin Hydrochloride Extended-Release Tablets vs Metformin Hydrochloride Tablets in Adults with Type 2 Diabetes Mellitus

	Metformin	**Metformin Hydrochloride Extended release tablets**

	Hydrochloride Tablets 500 mg Twice Daily	- Release tablets 1,000 mg Once Daily	1,500 mg Once Daily
Hemoglobin A1c (%)	**(n=67)**	**(n=72)**	**(n=66)**
Baseline	7.06	6.99	7.02
Change at FINAL VISIT	0.14 [a]	0.27	0.13
(95% CI)	(–0.04, 0.31)	(0.11, 0.43)	(–0.02, 0.28)
FPG (mg/dL)	**(n=69)**	**(n=72)**	**(n=70)**
Baseline	127.2	131	131.4
Change at FINAL VISIT	14	11.5	7.6
(95% CI)	(7, 21)	(4.4, 18.6)	(1, 14.2)

†a n=68

Mean baseline body weight was 210 lbs, 203 lbs and 193 lbs in the metformin hydrochloride tablets 500 mg twice daily, and metformin hydrochloride extended-release tablets 1,000 mg and 1,500 mg once daily arms, respectively. Mean change in body weight from baseline to week 24 was 0.9 lbs, 1.1 lbs and 0.9 lbs, respectively.

16 HOW SUPPLIED/STORAGE AND HANDLING

NA

16.1 How Supplied

Table 8: Metformin hydrochloride extended-release tablets, USP Available Strengths, Units, and Appearance

Metformin hydrochloride extended-release tablets, USP			
500 mg	Bottles of 30	NDC 62207-491-54	White to off-white uncoated, modified capsule shaped tablets debossed with “G7” on one side and plain on other side.
	Bottles of 90	NDC 62207-491-57	
	Bottles of 100	NDC 62207-491-43	
	Bottles of 500	NDC 62207-491-47	
	Bottles of 1,000	NDC 62207-491-49	
750 mg	Bottles of 30	NDC 62207-492-54	White to off-white uncoated modified capsule shaped tablets debossed with “G8” on one side and plain on other side.
	Bottles of 90	NDC 62207-492-57	
	Bottles of 100	NDC 62207-492-43	
	Bottles of 500	NDC 62207-492-47	
	Bottles	NDC	

of 1,000	62207-492-49

16.2 Storage

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Dispense in light-resistant container.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Lactic Acidosis:
Explain the risks of lactic acidosis, its symptoms, and conditions that predispose to its development. Advise patients to discontinue metformin hydrochloride extended-release tablets immediately and to promptly notify their healthcare provider if unexplained hyperventilation, myalgias, malaise, unusual somnolence or other nonspecific symptoms occur. Counsel patients against excessive alcohol intake and inform patients about importance of regular testing of renal function while receiving metformin hydrochloride extended-release tablets. Instruct patients to inform their doctor that they are taking metformin hydrochloride extended-release tablets prior to any surgical or radiological procedure, as temporary discontinuation may be required [see *Warnings and Precautions (5.1)].*

Hypoglycemia
Inform patients that hypoglycemia may occur when metformin hydrochloride extended-release tablets are coadministered with oral sulfonylureas and insulin. Explain to patients receiving concomitant therapy the risks of hypoglycemia, its symptoms and treatment, and conditions that predispose to its development [see *Warnings and Precautions (5.3)].*

Vitamin B_{12} Deficiency:
Inform patients about importance of regular hematological parameters while receiving metformin hydrochloride extended-release tablets [see *Warnings and Precautions (5.2)].*

Females of Reproductive Age:
Inform females that treatment with metformin hydrochloride extended-release tablets may result in ovulation in some premenopausal anovulatory women which may lead to unintended pregnancy [see Use in Specific Populations (8.3)*].*

Metformin Hydrochloride Extended-Release Tablets Administration Information:
Inform patients that metformin hydrochloride extended-release tablets must be swallowed whole and not crushed, cut, or chewed, and that the inactive ingredients may occasionally be eliminated in the feces as a soft mass that may resemble the original tablet.

Manufactured by:
Granules India Limited
Hyderabad-500084, India

Manufactured for:
Granules Pharmaceuticals Inc.
Chantilly, VA 20151

Revised: 07/2026

SPL PATIENT PACKAGE INSERT SECTION

PATIENT INFORMATION
Metformin Hydrochloride Extended-Release Tablets
(met-FOR-min HYE-droe-KLOR-ide)

Read the Patient Information that comes with metformin hydrochloride extended-release tablets before you start taking it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your healthcare provider about your medical condition or treatment.

What is the most important information I should know about metformin hydrochloride extended-release tablets?

Serious side effects can happen in people taking metformin hydrochloride extended-release tablets, including:

Lactic Acidosis. Metformin hydrochloride, the medicine in metformin hydrochloride extended-release tablets, can cause a rare, but serious, side effect called lactic acidosis (a build-up of lactic acid in the blood) that can cause death. Lactic acidosis is a medical emergency and must be treated in a hospital.

Stop taking in metformin hydrochloride extended-release tablets and call your healthcare provider right away if you get any of the following symptoms of lactic acidosis:
• feel very weak and tired
• have unusual (not normal) muscle pain
• have trouble breathing
• have unusual sleepiness or sleep longer than usual
• have unexplained stomach or intestinal problems with nausea and vomiting, or diarrhea
• feel cold, especially in your arms and legs
• feel dizzy or lightheaded
• have a slow or irregular heartbeat

You have a higher chance of getting lactic acidosis if you:
• have kidney problems. People whose kidneys are not working properly should not take metformin
hydrochloride extended-release tablets.
• have liver problems.
• have congestive heart failure that requires treatment with medicines.
• drink a lot of alcohol (very often or short-term “binge” drinking).
• get dehydrated (lose a large amount of body fluids). This can happen if you are sick with a fever, vomiting, or diarrhea. Dehydration can also happen when you sweat a lot with activity or exercise and do not drink enough fluids.
• have certain x-ray tests with injectable dyes or contrast agents.
• have surgery.
• have a heart attack, severe infection, or stroke.
• are 80 years of age or older and have not had your kidney function tested.

• have a genetically inherited disease affecting mitochondria (the energy-producing parts within cells) such as mitochondrial encephalomyopathy with lactic acidosis, and stroke-like episodes (MELAS) syndrome and maternally inherited diabetes and deafness (MIDD).

What are metformin hydrochloride extended-release tablets?
• Metformin hydrochloride extended-release tablets are prescription medicines that contain metformin hydrochloride. Metformin hydrochloride extended-release tablets are used with diet and exercise to help control high blood sugar (hyperglycemia) in adults with type 2 diabetes.
• Metformin hydrochloride extended-release tablets are not for people with type 1 diabetes.
• Metformin hydrochloride extended-release tablets are not for people with diabetic ketoacidosis (increased ketones in your blood or urine).

Metformin hydrochloride extended-release tablets works longer in your body. This medicine help control your blood sugar in a number of ways. These include helping your body respond better to the insulin it makes naturally, decreasing the amount of sugar your liver makes, and decreasing the amount of sugar your intestines absorb. Metformin hydrochloride extended-release tablets do not cause your body to make

more insulin.

Who should not take metformin hydrochloride extended-release tablets?
Some conditions increase your chance of getting lactic acidosis, or cause other problems if you take either of these medicines. Most of the conditions listed below can increase your chance of getting lactic acidosis.

Do not take metformin hydrochloride extended-release tablets if you:
• have kidney problems
• are allergic to the metformin hydrochloride in metformin hydrochloride extended-release tablets or any of the ingredients in metformin hydrochloride extended-release tablets. See the end of this leaflet for a complete list of ingredients in metformin hydrochloride extended-release tablets.
• are going to get an injection of dye or contrast agents for an x-ray procedure or if you are going to have surgery and not able to eat or drink much. In these situations, metformin hydrochloride extended-release tablets will need to be stopped for a short time. Talk to your healthcare provider about when you should stop metformin hydrochloride extended-release tablets and when you should start metformin hydrochloride extended-release tablets again. See **“What is the most important information I should know about metformin hydrochloride extended-release tablets?"**

What should I tell my healthcare provider before taking metformin hydrochloride extended-release tablet?
Before taking metformin hydrochloride extended-release tablets, tell your healthcare provider if you:
• have type 1 diabetes. Metformin hydrochloride extended-release tablets should not be used to treat people with type 1 diabetes.

• have a history or risk for diabetic ketoacidosis (high levels of certain acids, known as ketones, in the blood or urine). Metformin hydrochloride extended-release tablets should not be used for the treatment of diabetic ketoacidosis.
• have kidney problems.
• have liver problems.
• have heart problems, including congestive heart failure.
• are older than 80 years. If you are over 80 years old you should not take metformin hydrochloride extended-release tablets unless your kidneys have been checked and they are normal.
• drink alcohol very often, or drink a lot of alcohol in short-term “binge” drinking.
• are taking insulin.
• have any other medical conditions.
• are pregnant or plan to become pregnant. It is not known if metformin hydrochloride extended-release tablets will harm your unborn baby. If you are pregnant, talk with your healthcare provider about the best way to control your blood sugar while you are pregnant.
• are breast-feeding or plan to breast-feed. It is not known if metformin hydrochloride passes into your breast milk. Talk with your healthcare provider about the best way to feed your baby while you take metformin hydrochloride extended-release tablets.

• have a genetically inherited disease affecting mitochondria (the energy-producing parts within cells) such as MELAS syndrome and MIDD.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.
• Metformin hydrochloride extended-release tablets may affect the way other medicines work, and other medicines may affect how metformin hydrochloride extended-release tablets works.

Can metformin hydrochloride extended-release tablets be used in children?
Metformin hydrochloride extended-release tablets has not been studied in children.

How should I take metformin hydrochloride extended-release tablets?

• Take metformin hydrochloride extended-release tablets exactly as your healthcare provider tells you.
• Metformin hydrochloride extended-release tablets should be taken with meals to help lessen an upset stomach side effect.
• Swallow metformin hydrochloride extended-release tablets whole. Do not crush, cut, or chew metformin hydrochloride extended-release tablets.
• You may sometimes pass a soft mass in your stools (bowel movement) that looks like metformin hydrochloride extended-release tablets. This is not harmful and will not affect the way metformin hydrochloride extended-release tablets works to control your diabetes.
• When your body is under some types of stress, such as fever, trauma (such as a car accident), infection, or surgery, the amount of diabetes medicine that you need may change. Tell your healthcare provider right away if you have any of these problems.
• Your healthcare provider should do blood tests to check how well your kidneys are working before and during your treatment with metformin hydrochloride extended-release tablets.
• Your healthcare provider will check your diabetes with regular blood tests, including your blood sugar levels and your hemoglobin A1C.
• Follow your healthcare provider's instructions for treating blood sugar that is too low (hypoglycemia). Talk to your healthcare provider if low blood sugar is a problem for you. See **"What are the possible side effects of metformin hydrochloride extended-release tablets?"**
• Check your blood sugar as your healthcare provider tells you to.
• Stay on your prescribed diet and exercise program while taking metformin hydrochloride extended-release tablets.
• If you miss a dose of metformin hydrochloride extended-release tablets, take your next dose as
prescribed unless your healthcare provider tells you differently. Do not take an extra dose the next day.
• If you take too much metformin hydrochloride extended-release tablets, call your healthcare provider,
local Poison Control Center, or go to the nearest hospital emergency room right away.

What should I avoid while taking metformin hydrochloride extended-release tablets?
Do not drink a lot of alcoholic drinks while taking metformin hydrochloride extended-release tablets. This means you should not binge drink for short periods, and you should not drink a lot of alcohol on a regular basis. Alcohol can increase the chance of getting lactic acidosis.

What are the side effects of metformin hydrochloride extended-release tablets?
• Lactic acidosis. Metformin, the active ingredient in metformin hydrochloride extended-release tablets, can cause a rare but serious condition called lactic acidosis (a buildup of an acid in the blood) that can cause death. Lactic acidosis is a medical emergency and must be treated in the hospital.

Call your doctor right away if you have any of the following symptoms, which could be signs of lactic
acidosis:
• you feel cold in your hands or feet
• you feel dizzy or lightheaded
• you have a slow or irregular heartbeat
• you feel very weak or tired
• you have trouble breathing
• you feel sleepy or drowsy
• you have stomach pains, nausea or vomiting

Most people who have had lactic acidosis with metformin have other things that, combined with the metformin, led to the lactic acidosis. Tell your doctor if you have any of the following, because you have a higher chance for getting lactic acidosis with metformin hydrochloride extended-release tablets, if you:

• have severe kidney problems, or your kidneys are affected by certain x–ray tests that use injectable dye
• have liver problems
• drink alcohol very often, or drink a lot of alcohol in short-term “binge” drinking
• get dehydrated (lose a large amount of body fluids). This can happen if you are sick with a fever, vomiting, or diarrhea. Dehydration can also happen when you sweat a lot with activity or exercise and do not drink enough fluids
• have surgery
• have a heart attack, severe infection, or stroke

Common side effects of metformin hydrochloride extended-release tablets, include diarrhea, nausea, and upset stomach. These side effects generally go away after you take the medicine for a while. Taking your medicine with meals can help reduce these side effects. Tell your doctor if the side effects bother you a lot, last for more than a few weeks, come back after they've gone away, or start later in therapy. You may need a lower dose or need to stop taking the medicine for a short period or for good.

About 3 out of every 100 people who take metformin hydrochloride extended-release tablets have an unpleasant metallic taste when they start taking the medicine. It lasts for a short time.

Metformin hydrochloride extended-release tablets rarely cause hypoglycemia (low blood sugar) by themselves. However, hypoglycemia can happen if you do not eat enough, if you drink alcohol, or if you take other medicines to lower blood sugar.

How should I store metformin hydrochloride extended-release tablets?
Store metformin hydrochloride extended-release tablet at 68°F to 77°F (20°C to 25°C).

Keep metformin hydrochloride extended-release tablets and all medicines out of the reach of children.

General information about the use of metformin hydrochloride extended-release tablet
If you have questions or problems, talk with your doctor or other healthcare provider. You can ask your doctor or pharmacist for the information about metformin hydrochloride extended-release tablets that is written for healthcare professionals. Medicines are sometimes prescribed for purposes other than those listed in a patient information leaflet. Do not use metformin hydrochloride extended-release tablets for a condition for which it was not prescribed. Do not share your medicine with other people.

What are the ingredients of metformin hydrochloride extended-release tablets?
Active ingredients of metformin hydrochloride extended-release tablets: metformin hydrochloride.

Inactive ingredients in each tablet of metformin hydrochloride extended-release tablet 500 mg: hypromellose, magnesium stearate, and polyvinyl pyrrolidone.

Inactive ingredients in each tablet of metformin hydrochloride extended-release tablet 750 mg: hypromellose, magnesium stearate, and polyvinyl pyrrolidone.

What is type 2 diabetes?
Type 2 diabetes is a condition in which your body does not make enough insulin, and the insulin that your body produces does not work as well as it should. Your body can also make too much sugar. When this happens, sugar (glucose) builds up in the blood. This can lead to serious medical problems.
The main goal of treating diabetes is to lower your blood sugar to a normal level. High blood sugar can be lowered by diet and exercise, and by certain medicines when necessary.

Talk to your healthcare provider about how to prevent, recognize, and take care of low blood sugar (hypoglycemia), high blood sugar (hyperglycemia), and problems you have because of your diabetes.

Manufactured by:

Granules India Limited
Hyderabad-500084, India

Manufactured for:
Granules Pharmaceuticals Inc.
Chantilly, VA 20151

Revised: 07/2026

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL



Metformin Hydrochloride Extended-Release Tablets USP, 500 mg

Batch No. : XXXXXXX

No. of Units : 5,000 Tablets.

Mfg.Dt. : MM/YYYY

Exp.Dt. : MM/YYYY

Mfg.Lic.No. : 37/RR/AP/2003/F/R

NDC No. : 62207-491-31

Storage : Store at 20°–25° C (68°–77° F); excursions permitted to 15°–30° C (59°–86° F). [See USP Controlled Room Temperature.] Dispense in light-resistant containers.

Caution : For repacking only.

Country of Origin: INDIA

Product code : XXXXXXX

Manufactured by:
Granules India Limited

Manufactured for:
Granules Pharmaceuticals Inc.
Chantilly, VA 20151

Metformin Hydrochloride Extended-Release Tablets USP, 750 mg

Batch No. : XXXXXXX

No. of Units : 3500 Tablets.

Mfg.Dt. : MM/YYYY

Exp.Dt. : MM/YYYY

Mfg.Lic.No. : 37/RR/AP/2003/F/R

NDC No. : 62207-492-40

Storage : Store at 20°–25° C (68°–77° F); excursions permitted to 15°–30° C (59°–86° F). [See USP Controlled Room Temperature.] Dispense in light-resistant containers.

Caution : For repacking only.

Country of Origin: INDIA

Product code : XXXXXXXX

Manufactured by:
Granules India Limited

Manufactured for:
Granules Pharmaceuticals Inc.
Chantilly, VA 20151

NDC 62207-492-54

Metformin Hydrochloride Extended-Release Tablets, USP

750 mg

Rx only

30 Tablets

Each tablet contains 750 mg of metformin hydrochloride, USP.
See enclosed package insert for dosage information.

Warning: Keep out of the reach of children.

Store at 20° to 25° C (68° to 77° F); excursions permitted within 15° to 30° C (59° to 86° F). [See USP Controlled Room Temperature.]

Dispense in light-resistant container.

Dispense with Patient Information available at:
https://granulesindia.com/product/metformin-er-tab-2/

Manufactured by:
Granules India Limited
Hyderabad – 500 081, India.

MADE IN INDIA

R 10/22 20000000XXXX

37/RR/AP/2003/F/R

N 3 62207 49254 6

Un Varnish Zone

Metformin Hydrochloride Extended-Release Tablets USP, 750 mg

Batch No. : XXXXXXX
Mfg.Dt. : MM/YYYY
Exp.Dt. : MM/YYYY
Mfg.Lic.No. : 37/RR/AP/2003/F/R
NDC No. : 62207-492-74

No. of Units : 5 x 3,500
Gross Wt. : 00.000 kg
Tare Wt. : 00.000 kg
Net Wt. : 00.000 kg
Case No. : 000

Storage : Store at 20°–25° C (68°–77° F); excursions permitted to 15°–30° C (59°–86° F). [See USP Controlled Room Temperature.] Dispense in light-resistant containers.

Caution : For repacking only.

(01)50362207492749(17)000000(10)XXXXXXXX

Country of Origin: INDIA

8903832000000005

Product code : XXXXXXXX

Manufactured by:
Granules India Limited

Manufactured for:
Granules Pharmaceuticals Inc.
Chantilly, VA 20151

Metformin Hydrochloride Extended-Release Tablets USP, 500 mg

Batch No. : XXXXXXX
Mfg.Dt. : MM/YYYY
Exp.Dt. : MM/YYYY
Mfg.Lic.No. : 37/RR/AP/2003/F/R
NDC No. : 62207-491-35

No. of Units : 5 x 5,000
Gross Wt. : 00.000 kg
Tare Wt. : 00.000 kg
Net Wt. : 00.000 kg
Case No. : 000

Storage : Store at 20°–25° C (68°–77° F); excursions permitted to 15°–30° C (59°–86° F). [See USP Controlled Room Temperature.] Dispense in light-resistant containers.

Caution : For repacking only.

(01)50362207491353(17)000000(10)XXXXXXXX

Country of Origin: INDIA

8903832000000004

Product code : XXXXXXXX

Manufactured by:
Granules India Limited

Manufactured for:
Granules Pharmaceuticals Inc.
Chantilly, VA 20151

METFORMIN ER 500 MG

metformin er 500 mg tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:62207-491

Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METFORMIN HYDROCHLORIDE (UNII: 786Z46389E) (METFORMIN - UNII:9100L32L2N)	METFORMIN HYDROCHLORIDE	500 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white (white to off-white)	**Score**	no score
Shape	CAPSULE	**Size**	18mm
Flavor		**Imprint Code**	G7
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62207-491-54	30 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	
2	NDC:62207-491-49	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	
3	NDC:62207-491-43	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	
4	NDC:62207-491-47	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	
5	NDC:62207-491-31	5000 in 1 POUCH; Type 0: Not a Combination Product	06/28/2018	
6	NDC:62207-491-35	25000 in 1 CARTON; Type 0: Not a Combination Product	06/28/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209313	06/28/2018	

METFORMIN ER 750 MG

metformin er 750 mg tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:62207-492
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METFORMIN HYDROCHLORIDE (UNII: 786Z46389E) (METFORMIN - UNII:9100L32L2N)	METFORMIN HYDROCHLORIDE	750 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white (white to off-white)	**Score**	no score
Shape	CAPSULE	**Size**	19mm
Flavor		**Imprint Code**	G8
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62207-492-54	30 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	
2	NDC:62207-492-49	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	
3	NDC:62207-492-43	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	
4	NDC:62207-492-47	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2019	
5	NDC:62207-492-40	3500 in 1 POUCH; Type 0: Not a Combination Product	06/28/2018	
6	NDC:62207-492-74	25000 in 1 CARTON; Type 0: Not a Combination Product	06/28/2018	10/15/2020

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209313	06/28/2018	

Labeler - Granules India Ltd (915000087)

Establishment

Name	Address	ID/FEI	Business Operations
Granules India Ltd		918609236	analysis(62207-491, 62207-492) , label(62207-491, 62207-492) , manufacture(62207-491, 62207-492) , pack(62207-491, 62207-492)

Revised: 7/2026 Granules India Ltd