DEXOLYTE SOLUTION- devolyte injection, solution Clipper

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Dexolyte Solution Injection

INDICATIONS:

For use in conditions associated with fluid and electrolyte loss, such as dehydration, shock, vomiting and diarrhea, particularly when an immediate source of energy is also indicated.

CONTRAINDICATIONS:

Do not administer intraperitoneally to horses.

CAUTION:

This product contains no preservatives. Use entire contents when first opened. Do not use if solution is not clear. Discard any unused solution.

DOSAGE AND ADMINISTRATION:

Warm solution to body temperature and administer slowly (10 to 30 mL per minute) by intravenous or intraperitoneal injection, using strict aseptic procedures.

Adult Cattle and **Horses** - 1000 to 2000 mL **Calves, Ponies** and **Foals** - 500 to 1000 mL**Adult Sheep** and **Swine** - 500 to 1000 mL

These are suggested dosages. The actual amount and rate of fluid administration must be judged by the veterinarian in relation to the condition being treated and the clinical response of the animal, being careful to avoid overhydration.

TAKE TIME OBSERVE LABEL DIRECTIONS

RMS 92-1086 Rev. 07-15

18-804

Lot No. Exp. Date

Manufactured by Nova-Tech, Inc. Grand Island, NE 68801

Trademarks are property of Clipper Distributing Company, LLC

Manufactured for:

Clipper Distributing Company, LLCSt. Joseph, MO 64507

Net Contents: 500 mL

ELECTROLYTES - DEXTROSE

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

Caution:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

COMPOSITION:

Store between 15 degrees C and 30 degrees C (59 degrees F and 86 degrees F).

INDICATIONS: For use in conditions associated with fluid and electrolyte loss, such as dehydration, shock, vomiting and diarrhea, particularly when an immediate source of energy is also indicated.

 $\begin{tabular}{ll} \textbf{CONTRAINDICATIONS:} Do not administer intraperitoneally to horses. \end{tabular}$

CAUTION: This product contains no preservatives. Use entire contents when first opened. Do not use if solution is not clear. Discard any unused solution.

DOSAGE AND ADMINISTRATION: Warm solution to body temperature and administer slowly (10 to 30 mL per minute) by intravenous or intraperitoneal injection, using strict aseptic procedures.

Adult Cattle and Horses – 1000 to 2000 mL Calves, Ponies and Foals – 500 to 1000 mL Adult Sheep and Swine – 500 to 1000 mL

These are suggested dosages. The actual amount and rate of fluid administration must be judged by the veterinarian in relation to the condition being treated and the clinical response of the animal, being careful to avoid overhydration.



RMS 92-1086 18-804 Rev. 07-15

Lot No.

Exp. Date



ELECTROLYTES – DEXTROSE

FOR ANIMAL USE ONLY
KEEP OUT OF REACH OF CHILDREN

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian. Net Contents: 500 mL

PHOENIX

Manufactured for:
Clipper Distributing Company, LLC
St. Joseph, MO 64507

Manufactured by Nova-Tech, Inc. Grand Island, NE 68801

Trademarks are property of Clipper Distributing Company, LLC



COMPOSITION: Each 500 mL of sterile aqueous

Milliequivalents per liter		
Cations SodiumPotassium	9	mEq/L
Magnesium	4	mEq/L
ChlorideLactate		
Osmolarity (calc.) 602 mOsmol/L		

Store between 15°C and 30°C (59°F and 86°F).



DEXOLYTE SOLUTION

dexolyte injection, solution

Product Information				
Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:57319-617	
Route of Administration	INTRAVENOUS, INTRAPERITONEAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Dextrose Monohydrate (UNII: LX22YL083G) (Anhydrous Dextrose - UNII:5SL0G7R0OK)	Dextrose Monohydrate	12.50 g in 500 mL	
Sorbitol (UNII: 506T60A25R) (Sorbitol - UNII:506T60A25R)	Sorbitol	12.50 in 500 mL	
Sodium Lactate (UNII: TU7HW0W0QT) (Sodium Cation - UNII:LYR4M0NH37)	Sodium Lactate	3.95 in 500 mL	
Sodium Chloride (UNII: 451W47IQ8X) (Chloride Ion - UNII:Q32ZN48698)	Sodium Chloride	2.40 in 500 mL	
Potassium Chloride (UNII: 660 YQ98 I10) (Potassium Cation - UNII:295053K152)	Potassium Chloride	0.37 in 500 mL	
Magnesium Chloride (UNII: 02F3473H9O) (Magnesium Cation - UNII:T6V3LHY838)	Magnesium Chloride	0.21 in 500 mL	
Calcium Chloride (UNII: M4I0 D6 VV5M) (Calcium Cation - UNII:2M8 3C4R6 ZB)	Calcium Chloride	0.19 in 500 mL	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57319-617-07	500 mL in 1 BOTTLE, PLASTIC		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/01/2015	

Labeler - Clipper (150711039)

Registrant - Clipper (150711039)

Establishment				
Name	Address	ID/FEI	Business Operations	
Nova-Tech, Inc.		196078976	manufacture, api manufacture	

Revised: 12/2015 Clipper