

ZOLPIDEM TARTRATE- zolpidem tartrate tablet

Par Pharmaceutical, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ZOLPIDEMTARTRATE Sublingual Tablets safely and effectively. See full prescribing information for ZOLPIDEMTARTRATE Sublingual Tablets.

ZOLPIDEMTARTRATE Sublingual Tablets, for oral use C-IV

Initial U.S. Approval: 1992

----- INDICATIONS AND USAGE -----

Zolpidem Tartrate Sublingual Tablets, a gamma-aminobutyric acid (GABA) A agonist, is indicated for the short-term treatment of insomnia characterized by difficulties with sleep initiation. Zolpidem Tartrate has been shown to decrease sleep latency for up to 35 days in controlled clinical studies. (1) (1)

----- DOSAGE AND ADMINISTRATION -----

- Use the lowest dose effective for the patient (2.1)
- Recommended dose is 5 mg for women and 5 or 10 mg for men, immediately before bedtime (2.1)
- Geriatric patients and patients with hepatic impairment: Recommended dose is 5 mg for men and women (2.2)
- Lower doses of CNS depressants may be necessary when taken concomitantly with zolpidem tartrate sublingual tablets (2.3)
- Coadministration with CNS depressants: Recommended dose is 5 mg for men and women (2.3)
- The effect of Zolpidem Tartrate Sublingual Tablets may be slowed if taken with or immediately after a meal (2.4)
- Zolpidem Tartrate Sublingual Tablet should be placed under the tongue, where it will disintegrate (2.4). The tablet should not be swallowed and the tablet should not be taken with water (2.4)

----- DOSAGE FORMS AND STRENGTHS -----

Sublingual tablets: 5 mg and 10 mg. Tablets not scored. (3) (3)

----- CONTRAINDICATIONS -----

Known hypersensitivity to zolpidem (4) (4)

----- WARNINGS AND PRECAUTIONS -----

- CNS depressant effects: Impairs alertness and motor coordination. Instruct patients on correct use. (5.1)
- Need to evaluate for co-morbid diagnosis. Reevaluate if insomnia persists after 7 to 10 days of use. (5.2)
- Severe anaphylactic and anaphylactoid reactions: angioedema and anaphylaxis have been reported. Do not rechallenge if such reactions occur. (5.3)
- "Sleep-driving" and other complex behaviors while not fully awake. Risk increases with dose and use with other CNS depressants and alcohol. Immediately evaluate any new onset behavioral changes. (5.4)
- Depression: Worsening of depression or suicidal thinking may occur. Prescribe the least amount of tablets feasible to avoid intentional overdose. (5.5)
- Respiratory Depression: Consider this risk before prescribing in patients with compromised respiratory function. (5.6)
- Withdrawal effects: Symptoms may occur with rapid dose reduction or discontinuation. (5.7, 9.3)

----- ADVERSE REACTIONS -----

Most commonly observed adverse reactions were: (6)

Short-term (<10 nights): Drowsiness, dizziness and diarrhea (6)

Long-term (28 to 35 nights): Dizziness and drugged feelings (6.1) (6)

To report SUSPECTED ADVERSE REACTIONS, contact Par Pharmaceutical at 1-800-828-9393 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. (6)

----- DRUG INTERACTIONS -----

- CNS-depressants, including alcohol: Possible adverse additive CNS-depressant effects (5.1, 7.1)
- Imipramine: decreased alertness observed (7.1)
- Chlorpromazine: impaired alertness and psychomotor performance observed (7.1)
- Rifampin: combination use may decrease effects (7.2)
- Ketoconazole: combination use may increase effect (7.2)

----- USE IN SPECIFIC POPULATIONS -----

- Pregnancy: Based on animal data, may cause fetal harm. (8.1)
- Pediatric use: Safety and effectiveness not established. Hallucinations (incidence rate 7%) and other psychiatric and/or

nervous system adverse reactions were observed frequently in a study of pediatric patients with Attention-Deficit/Hyperactivity Disorder. (5.4, 8.4)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 11/2017

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS & USAGE

2 DOSAGE & ADMINISTRATION

- 2.1 Dosage in adults
- 2.2 Special Populations
- 2.3 Use with CNS depressants
- 2.4 Administration

3 DOSAGE FORMS & STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 CNS Depressant Effects and Next-Day Impairment
- 5.2 Need to Evaluate for Co-morbid Diagnoses
- 5.3 Severe Anaphylactic and Anaphylactoid Reactions
- 5.4 Abnormal Thinking and Behavioral Changes
- 5.5 Use in Patients with Depression
- 5.6 Respiratory Depression
- 5.7 Withdrawal Effects

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 CNS-active drugs
- 7.2 Drugs that Affect Drug Metabolism Via Cytochrome P450

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor & Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Gender Difference in Pharmacokinetics

9 DRUG ABUSE AND DEPENDENCE

- 9.1 Controlled Substance
- 9.2 Abuse
- 9.3 Dependence

10 OVERDOSAGE

- 10.1 Signs and Symptoms
- 10.2 Recommended Treatment

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis & Mutagenesis & Impairment Of Fertility

14 CLINICAL STUDIES

- 14.1 Transient Insomnia

14.2 Chronic Insomnia

14.3 Studies Pertinent to Safety Concerns for Sedative/Hypnotic Drugs

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS & USAGE

Zolpidem Tartrate Sublingual Tablets are indicated for the short-term treatment of insomnia characterized by difficulties with sleep initiation [see *CLINICAL STUDIES (14)*].

The clinical trials performed with zolpidem tartrate in support of efficacy were 4 to 5 weeks in duration with the final formal assessments of sleep latency performed at the end of treatment.

2 DOSAGE & ADMINISTRATION

2.1 Dosage in adults

Use the lowest effective dose for the patient. The recommended initial dose is 5 mg for women and either 5 or 10 mg for men, taken only once per night immediately before bedtime with at least 7 to 8 hours remaining before the planned time of awakening. If the 5 mg dose is not effective, the dose can be increased to 10 mg. In some patients, the higher morning blood levels following use of the 10 mg dose increase the risk of next day impairment of driving and other activities that require full alertness [see *Warnings and Precautions (5.1)*]. The total dose of Zolpidem Tartrate Sublingual Tablets should not exceed 10 mg once daily immediately before bedtime.

The recommended initial doses for women and men are different because zolpidem clearance is lower in women.

2.2 Special Populations

Elderly or debilitated patients may be especially sensitive to the effects of zolpidem tartrate. Patients with hepatic insufficiency do not clear the drug as rapidly as normal subjects. The recommended dose of zolpidem tartrate sublingual tablets in both of these patient populations is 5 mg once daily immediately before bedtime [see *Warnings and Precautions (5.1)*; *Use in Specific Populations (8.5)*].

2.3 Use with CNS depressants

Dosage adjustment may be necessary when Zolpidem Tartrate Sublingual Tablets are combined with other CNS-depressant drugs because of the potentially additive effects [see *Warnings and Precautions (5.1)*].

2.4 Administration

The effect of Zolpidem Tartrate Sublingual Tablets may be slowed by ingestion with or immediately after a meal.

Zolpidem Tartrate Sublingual Tablets should be placed under the tongue, where it will disintegrate. The

tablet should not be swallowed and the tablet should not be taken with water.

3 DOSAGE FORMS & STRENGTHS

Zolpidem Tartrate Sublingual Tablets are available in 5 mg and 10 mg strength tablets for sublingual administration. Tablets are not scored.

Zolpidem tartrate sublingual tablets 5 mg are white, round tablets, flat-faced, bevel-edged with debossed “P1” on one side and plain on the other side

Zolpidem tartrate sublingual tablets 10 mg are white, round tablets, flat-faced, bevel-edged with debossed “P7” on one side and plain on the other side.

4 CONTRAINDICATIONS

Zolpidem tartrate is contraindicated in patients with known hypersensitivity to zolpidem. Observed reactions include anaphylaxis and angioedema [*see Warnings and Precautions (5.3)*].

5 WARNINGS AND PRECAUTIONS

5.1 CNS Depressant Effects and Next-Day Impairment

Zolpidem tartrate, like other sedative-hypnotic drugs, has central nervous system (CNS) depressant effects. Coadministration with other CNS depressants (e.g., benzodiazepines, opioids, tricyclic antidepressants, alcohol) increases the risk of CNS depression. Dosage adjustments of zolpidem tartrate and of other concomitant CNS depressants may be necessary when zolpidem tartrate is administered with such agents because of the potentially additive effects. The use of zolpidem tartrate with other sedative-hypnotics (including other zolpidem products) at bedtime or the middle of the night is not recommended [*see Dosage and Administration (2.3)*].

The risk of next-day psychomotor impairment, including impaired driving, is increased if zolpidem tartrate is taken with less than a full night of sleep remaining (7 to 8 hours); if a higher than the recommended dose is taken; if coadministered with other CNS depressants; or if coadministered with other drugs that increase the blood level of zolpidem. Patients should be cautioned against driving and other activities requiring complete mental alertness if zolpidem tartrate is taken in these circumstances.

5.2 Need to Evaluate for Co-morbid Diagnoses

Because sleep disturbances may be the presenting manifestation of a physical and/or psychiatric disorder, symptomatic treatment of insomnia should be initiated only after a careful evaluation of the patient. The failure of insomnia to remit after 7 to 10 days of treatment may indicate the presence of a primary psychiatric and/or medical illness that should be evaluated. Worsening of insomnia or the emergence of new thinking or behavior abnormalities may be the consequence of an unrecognized psychiatric or physical disorder. Such findings have emerged during the course of treatment with sedative/hypnotic drugs, including zolpidem.

5.3 Severe Anaphylactic and Anaphylactoid Reactions

Cases of angioedema involving the tongue, glottis or larynx have been reported in patients after taking the first or subsequent doses of sedative-hypnotics, including zolpidem tartrate. Some patients have had additional symptoms such as dyspnea, throat closing or nausea and vomiting that suggest anaphylaxis. Some patients have required medical therapy in the emergency department. If angioedema involves the throat, glottis or larynx, airway obstruction may occur and be fatal. Patients who develop angioedema after treatment with zolpidem tartrate should not be rechallenged with the drug.

5.4 Abnormal Thinking and Behavioral Changes

Abnormal thinking and behavior changes have been reported in patients treated with sedative/hypnotics, including zolpidem. Some of these changes included decreased inhibition (e.g., aggressiveness and extroversion that seemed out of character), bizarre behavior, agitation and depersonalization. Visual and auditory hallucinations have been reported.

In controlled trials of zolpidem tartrate 10 mg taken at bedtime, <1% of adults with insomnia who received zolpidem reported hallucinations. In a clinical trial, 7% of pediatric patients treated with zolpidem tartrate 0.25 mg/kg taken at bedtime reported hallucinations, versus 0% treated with placebo [*see Use in Specific Populations (8.4)*].

Complex behaviors such as “sleep-driving” (i.e., driving while not fully awake after ingestion of a sedative-hypnotic, with amnesia for the event) have been reported in sedative-hypnotic-naive as well as in sedative-hypnotic-experienced persons. Although behaviors such as “sleep-driving” have occurred with zolpidem alone at therapeutic doses, the coadministration of zolpidem with alcohol or other CNS depressants increases the risk of such behaviors, as does the use of zolpidem tartrate at doses exceeding the maximum recommended dose. Due to the risk to the patient and the community, discontinuation of zolpidem tartrate should be strongly considered for patients who report a “sleep-driving” episode.

Other complex behaviors (e.g., preparing and eating food, making phone calls, or having sex) have been reported in patients who are not fully awake after taking a sedative-hypnotic. As with “sleep-driving”, patients usually do not remember these events. Amnesia, anxiety and other neuro-psychiatric symptoms may also occur.

It can rarely be determined with certainty whether a particular instance of the abnormal behaviors listed above is drug induced, spontaneous in origin, or a result of an underlying psychiatric or physical disorder. Nonetheless, the emergence of any new behavioral sign or symptom of concern requires careful and immediate evaluation.

5.5 Use in Patients with Depression

In primarily depressed patients treated with sedative-hypnotics, worsening of depression, and suicidal thoughts and actions (including completed suicides), have been reported. Suicidal tendencies may be present in such patients and protective measures may be required. Intentional overdose is more common in this group of patients; therefore, the least amount of drug that is feasible should be prescribed for the patient at any one time.

5.6 Respiratory Depression

Although studies with 10 mg zolpidem tartrate did not reveal respiratory depressant effects at hypnotic doses in healthy subjects or in patients with mild-to-moderate chronic obstructive pulmonary disease (COPD), a reduction in the Total Arousal Index, together with a reduction in lowest oxygen saturation and increase in the time of oxygen desaturation below 80% and 90%, was observed in patients with mild-to-moderate sleep apnea when treated with zolpidem compared to placebo. Since sedative-hypnotics have the capacity to depress respiratory drive, precautions should be taken if zolpidem tartrate is prescribed to patients with compromised respiratory function. Post-marketing reports of respiratory insufficiency in patients receiving 10 mg of zolpidem tartrate, most of whom had pre-existing respiratory impairment, have been reported. The risks of respiratory depression should be considered prior to prescribing zolpidem tartrate in patients with respiratory impairment including sleep apnea and myasthenia gravis.

5.7 Withdrawal Effects

There have been reports of withdrawal signs and symptoms following the rapid dose decrease or abrupt discontinuation of zolpidem. Monitor patients for tolerance, abuse, and dependence [see *Drug Abuse and Dependence (9.2) and (9.3)*].

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the labeling:

- CNS-depressant effects and next-day impairment [see *Warnings and Precautions (5.1)*]
- Serious anaphylactic and anaphylactoid reactions [see *Warning and Precautions (5.3)*]
- Abnormal thinking and behavior changes, complex behaviors [see *Warning and Precautions (5.4)*]
- Withdrawal effects [see *Warning and Precautions (5.7)*]

6.1 Clinical Trials Experience

Associated with discontinuation of treatment:

Approximately 4% of 1,701 patients who received zolpidem tartrate at all doses (1.25 to 90 mg) in U.S. premarketing clinical trials discontinued treatment because of an adverse reaction. Reactions most commonly associated with discontinuation from U.S. trials were daytime drowsiness (0.5%), dizziness (0.4%), headache (0.5%), nausea (0.6%), and vomiting (0.5%).

Approximately 4% of 1,959 patients who received zolpidem tartrate at all doses (1 to 50 mg) in similar foreign trials discontinued treatment because of an adverse reaction. Reactions most commonly associated with discontinuation from these trials were daytime drowsiness (1.1%), dizziness/vertigo (0.8%), amnesia (0.5%), nausea (0.5%), headache (0.4%), and falls (0.4%).

Data from a clinical study in which selective serotonin reuptake inhibitor (SSRI)-treated patients were given zolpidem tartrate revealed that four of the seven discontinuations during double-blind treatment with zolpidem (n=95) were associated with impaired concentration, continuing or aggravated depression, and manic reaction; one patient treated with placebo (n=97) was discontinued after an attempted suicide.

Most commonly observed adverse reactions in controlled trials:

During short-term treatment (up to 10 nights) with zolpidem tartrate at doses up to 10 mg, the most commonly observed adverse reactions associated with the use of zolpidem and seen at statistically significant differences from placebo-treated patients were drowsiness (reported by 2% of zolpidem patients), dizziness (1%), and diarrhea (1%). During longer-term treatment (28 to 35 nights) with zolpidem tartrate at doses up to 10 mg, the most commonly observed adverse reactions associated with the use of zolpidem and seen at statistically significant differences from placebo-treated patients were dizziness (5%) and drugged feelings (3%).

Adverse reactions observed at an incidence of $\geq 1\%$ in controlled trials:

The following tables enumerate treatment-emergent adverse event frequencies that were observed at an incidence equal to 1% or greater among patients with insomnia who received zolpidem tartrate and at a greater incidence than placebo in U.S. placebo-controlled trials. Events reported by investigators were classified utilizing a modified World Health Organization (WHO) dictionary of preferred terms for the purpose of establishing event frequencies. The prescriber should be aware that these figures cannot be used to predict the incidence of side effects in the course of usual medical practice, in which patient characteristics and other factors differ from those that prevailed in these clinical trials. Similarly, the cited frequencies cannot be compared with figures obtained from other clinical investigators involving related drug products and uses, since each group of drug trials is conducted under a different set of

conditions. However, the cited figures provide the physician with a basis for estimating the relative contribution of drug and nondrug factors to the incidence of side effects in the population studied.

The following table was derived from a pool of 11 placebo-controlled short-term U.S. efficacy trials involving zolpidem in doses ranging from 1.25 to 20 mg. The table is limited to data from doses up to and including 10 mg, the highest dose recommended for use.

TABLE 1: Incidence of Treatment-Emergent Adverse Experiences in Placebo-Controlled Clinical Trials with Zolpidem Tartrate Lasting up to 10 nights (Percentage of Patients Reporting)

Body System/ Adverse Event*	Zolpidem Tartrate (≤ 10 mg) (N=685)	Placebo (N=473)
Central and Peripheral Nervous System		
Headache	7	6
Drowsiness	2	-
Dizziness	1	-
Gastrointestinal System		
Diarrhea	1	-

*Reactions reported by at least 1% of patients treated with oral zolpidem and at a greater frequency than placebo.

The following table was derived from a pool of three placebo-controlled long-term efficacy trials involving oral zolpidem. These trials involved patients with chronic insomnia who were treated for 28 to 35 nights with zolpidem at doses of 5, 10, or 15 mg. The table is limited to data from doses up to and including 10 mg, the highest dose recommended for use. The table includes only adverse events occurring at an incidence of at least 1% for zolpidem patients.

TABLE 2: Incidence of Treatment-Emergent Adverse Experiences in Placebo-Controlled Clinical Trials with Zolpidem Tartrate Lasting up to 35 nights (Percentage of Patients Reporting)

Body System/ Adverse Event*	Zolpidem Tartrate (≤ 10 mg) (N=152)	Placebo (N=161)
Autonomic Nervous System		
Dry mouth	3	1
Body as a Whole		
Allergy	4	1
Back pain	3	2
Influenza-like symptoms	2	-
Chest pain	1	-
Cardiovascular System		
Palpitation	2	-
Central and Peripheral Nervous System		
Drowsiness	8	5
Dizziness	5	1
Lethargy	3	1

Drugged feeling	3	-
Lightheadedness	2	1
Depression	2	1
Abnormal dreams	1	-
Amnesia	1	-
Sleep disorder	1	-
Gastrointestinal System		
Diarrhea	3	2
Abdominal pain	2	2
Constipation	2	1
Respiratory System		
Sinusitis	4	2
Pharyngitis	3	1
Skin and Appendages		
Rash	2	1

*Reactions reported by at least 1% of patients treated with oral zolpidem and at a greater frequency than placebo.

Dose relationship for adverse reactions associated with oral zolpidem:

There is evidence from dose comparison trials suggesting a dose relationship for many of the adverse reactions associated with oral zolpidem use, particularly for certain CNS and gastrointestinal adverse events.

Oral tissue-related adverse reactions to zolpidem tartrate:

The effect of chronic daily administration of zolpidem tartrate on oral tissue was evaluated in a 60-day open-label study in 60 insomniac patients. One patient developed transient sublingual erythema, and another transient paresthesia of the tongue.

Adverse event incidence across the entire preapproval oral zolpidem database:

Zolpidem was administered to 3,660 subjects in clinical trials throughout the U.S., Canada, and Europe. Treatment-emergent adverse events associated with clinical trial participation were recorded by clinical investigators using terminology of their own choosing. To provide a meaningful estimate of the proportion of individuals experiencing treatment-emergent adverse events, similar types of untoward events were grouped into a smaller number of standardized event categories and classified utilizing a modified World Health Organization (WHO) dictionary of preferred terms.

The frequencies presented, therefore, represent the proportions of the 3,660 individuals exposed to zolpidem, at all doses, who experienced an event of the type cited on at least one occasion while receiving zolpidem. All reported treatment-emergent adverse events are included, except those already listed in the table above of adverse events in placebo-controlled studies, those coding terms that are so general as to be uninformative, and those events where a drug cause was remote. It is important to emphasize that, although the events reported did occur during treatment with zolpidem, they were not necessarily caused by it.

Adverse events are further classified within body system categories and enumerated in order of decreasing frequency using the following definitions: frequent adverse events are defined as those occurring in greater than 1/100 subjects; infrequent adverse events are those occurring in 1/100 to 1/1,000 patients; rare events are those occurring in less than 1/1,000 patients.

Autonomic nervous system: Infrequent: increased sweating, pallor, postural hypotension, syncope. *Rare:*

abnormal accommodation, altered saliva, flushing, glaucoma, hypotension, impotence, increased saliva, tenesmus.

Body as a whole: Frequent: asthenia. *Infrequent:* edema, falling, fever, malaise, trauma.

Rare: allergic reaction, allergy aggravated, anaphylactic shock, face edema, hot flashes, increased ESR, pain, restless legs, rigors, tolerance increased, weight decrease.

Cardiovascular system: Infrequent: cerebrovascular disorder, hypertension, tachycardia.

Rare: angina pectoris, arrhythmia, arteritis, circulatory failure, extrasystoles, hypertension aggravated, myocardial infarction, phlebitis, pulmonary embolism, pulmonary edema, varicose veins, ventricular tachycardia.

Central and peripheral nervous system: Frequent: ataxia, confusion, euphoria, headache, insomnia, vertigo. *Infrequent:* agitation, anxiety, decreased cognition, detached, difficulty concentrating, dysarthria, emotional lability, hallucination, hypoesthesia, illusion, leg cramps, migraine, nervousness, paresthesia, sleeping (after daytime dosing), speech disorder, stupor, tremor. *Rare:* abnormal gait, abnormal thinking, aggressive reaction, apathy, appetite increased, decreased libido, delusion, dementia, depersonalization, dysphasia, feeling strange, hypokinesia, hypotonia, hysteria, intoxicated feeling, manic reaction, neuralgia, neuritis, neuropathy, neurosis, panic attacks, paresis, personality disorder, somnambulism, suicide attempts, tetany, yawning.

Gastrointestinal system: Frequent: dyspepsia, hiccup, nausea. *Infrequent:* anorexia, constipation, dysphagia, flatulence, gastroenteritis, vomiting. *Rare:* enteritis, eructation, esophagospasm, gastritis, hemorrhoids, intestinal obstruction, rectal hemorrhage, tooth caries.

Hematologic and lymphatic system: Rare: anemia, hyperhemoglobinemia, leukopenia, lymphadenopathy, macrocytic anemia, purpura, thrombosis.

Immunologic system: Infrequent: infection. *Rare:* abscess herpes simplex herpes zoster, otitis externa, otitis media.

Liver and biliary system: Infrequent: abnormal hepatic function, increased SGPT. *Rare:* bilirubinemia, increased SGOT.

Metabolic and nutritional: Infrequent: hyperglycemia, thirst. *Rare:* gout, hypercholesteremia, hyperlipidemia, increased alkaline phosphatase, increased BUN, periorbital edema.

Musculoskeletal system: Frequent: arthralgia, myalgia. *Infrequent:* arthritis. *Rare:* arthrosis, muscle weakness, sciatica, tendinitis.

Reproductive system: Infrequent: menstrual disorder, vaginitis. *Rare:* breast fibroadenosis, breast neoplasm, breast pain.

Respiratory system: Frequent: upper respiratory infection. *Infrequent:* bronchitis, coughing, dyspnea, rhinitis. *Rare:* bronchospasm, epistaxis, hypoxia, laryngitis, pneumonia.

Skin and appendages: Infrequent: pruritus. *Rare:* acne, bullous eruption, dermatitis, furunculosis, injection-site inflammation, photosensitivity reaction, urticaria.

Special senses: Frequent: diplopia, vision abnormal. *Infrequent:* eye irritation, eye pain, scleritis, taste perversion, tinnitus. *Rare:* conjunctivitis, corneal ulceration, lacrimation abnormal, parosmia, photopsia.

Urogenital system: Frequent: urinary tract infection. *Infrequent:* cystitis, urinary incontinence. *Rare:* acute renal failure, dysuria, micturition frequency, nocturia, polyuria, pyelonephritis, renal pain, urinary retention.

7 DRUG INTERACTIONS

7.1 CNS-active drugs

Coadministration of zolpidem with other CNS depressants increases the risk of CNS depression [see *Warnings and Precautions (5.1)*]. Zolpidem tartrate was evaluated in healthy volunteers in single-dose interaction studies for several CNS drugs.

Imipramine, Chlorpromazine

Imipramine in combination with zolpidem produced no pharmacokinetic interaction other than a 20% decrease in peak levels of imipramine, but there was an additive effect of decreased alertness. Similarly, chlorpromazine in combination with zolpidem produced no pharmacokinetic interaction, but there was an additive effect of decreased alertness and psychomotor performance [see *Clinical Pharmacology (12.3)*].

Haloperidol

A study involving haloperidol and zolpidem revealed no effect of haloperidol on the pharmacokinetics or pharmacodynamics of zolpidem. The lack of a drug interaction following single-dose administration does not predict the absence of an effect following chronic administration [see *Clinical Pharmacology (12.3)*].

Alcohol

An additive adverse effect on psychomotor performance between alcohol and oral zolpidem was demonstrated [see *Warnings and Precautions (5.1)*].

Sertraline

Concomitant administration of zolpidem and sertraline increases exposure to zolpidem and may increase the pharmacodynamics effect of zolpidem [see *Clinical Pharmacology (12.3)*].

Fluoxetine

After multiple doses of zolpidem tartrate and fluoxetine an increase in the zolpidem half-life (17%) was observed. There was no evidence of an additive effect in psychomotor performance [see *Clinical Pharmacology (12.3)*].

7.2 Drugs that Affect Drug Metabolism Via Cytochrome P450

Some compounds known to inhibit CYP3A may increase exposure to zolpidem. The effect of other P450 enzymes on the exposure to zolpidem is not known.

Rifampin

Rifampin, a CYP3A4 inducer, significantly reduced the exposure to and the pharmacodynamics effects of zolpidem. Use of Rifampin in combination with zolpidem may decrease the efficacy of zolpidem.

Ketoconazole

Ketoconazole, a potent CYP3A4 inhibitor, increased the pharmacodynamics effects of zolpidem. Consideration should be given to using a lower dose of zolpidem when ketoconazole and zolpidem are given together.

8 USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal data, may cause fetal harm. (8.1)
- Pediatric use: Safety and effectiveness not established. Hallucinations (incidence rate 7%) and other psychiatric and/or nervous system adverse reactions were observed frequently in a study of pediatric patients with Attention-Deficit/Hyperactivity Disorder. (5.4, 8.4)

8.1 Pregnancy

Pregnancy Category C:

There are no adequate and well-controlled studies of zolpidem tartrate in pregnant women. Studies in children to assess the effects of prenatal exposure to zolpidem have not been conducted; however, cases of severe neonatal respiratory depression have been reported when zolpidem was used at the end of pregnancy, especially when taken with other CNS-depressants. Children born to mothers taking sedative-hypnotic drugs may be at risk for withdrawal symptoms during the postnatal period. Neonatal flaccidity has also been reported in infants born to mothers who received sedative-hypnotic drugs during pregnancy. Zolpidem tartrate should be used during pregnancy only if the potential benefit outweighs the potential risk to the fetus.

Administration of zolpidem to pregnant rats and rabbits resulted in adverse effects on offspring development at doses greater than the maximum recommended human dose (MRHD) of 10 mg/day (approximately 8 mg/day zolpidem base); however, teratogenicity was not observed.

When zolpidem was administered at oral doses of 4, 20, and 100 mg base/kg to pregnant rats during the period of organogenesis, dose-related decreases in fetal skull ossification occurred at all but the lowest dose, which is approximately 5 times the MRHD on a mg/m² basis. In rabbits treated during organogenesis with zolpidem at oral doses of 1, 4, and 16 mg base/kg, increased embryo-fetal death and incomplete fetal skeletal ossification occurred at the highest dose. The no-effect dose for embryo-fetal toxicity in rabbits is approximately 10 times the MRHD on a mg/m² basis. Administration of zolpidem to rats at oral doses of 4, 20, and 100 mg base/kg during the latter part of pregnancy and throughout lactation produced decreased offspring growth and survival at all but the lowest dose, which is approximately 5 times the MRHD on a mg/m² basis.

8.2 Labor & Delivery

Zolpidem tartrate has no established use in labor and delivery [*see Pregnancy (8.1)*].

8.3 Nursing Mothers

Zolpidem is excreted in human milk. Caution should be exercised when zolpidem tartrate is administered to a nursing woman.

8.4 Pediatric Use

Zolpidem tartrate is not recommended for use in children. Safety and effectiveness in pediatric patients have not been established in pediatric patients below the age of 18.

In an 8-week controlled study in 201 pediatric patients (aged 6 to 17 years) with insomnia associated with attention-deficit/hyperactivity disorder (ADHD), an oral solution of zolpidem tartrate dosed at 0.25mg/kg at bedtime did not decrease sleep latency compared to placebo. Ten patients on zolpidem (7.4%) discontinued treatment due to an adverse reaction.

Psychiatric and nervous system disorders comprised the most frequent (>5%) treatment emergent adverse reactions observed with zolpidem versus placebo and included dizziness (23.5% vs. 1.5%), headache (12.5% vs. 9.2%), and hallucinations reported in 7% of the pediatric patients who received

zolpidem; none of the pediatric patients who received placebo reported hallucinations [see *Warnings and Precautions (5.4)*].

8.5 Geriatric Use

A total of 154 patients in U.S. controlled clinical trials and 897 patients in non-U.S. clinical trials who received oral zolpidem were ≥ 60 years of age. For a pool of U.S. patients receiving zolpidem tartrate at doses of ≤ 10 mg or placebo, there were three adverse events occurring at an incidence of at least 3% for zolpidem and for which the zolpidem incidence was at least twice the placebo incidence (i.e., they could be considered drug-related).

Adverse Event	Zolpidem	Placebo
Dizziness	3%	0%
Drowsiness	5%	2%
Diarrhea	3%	1%

A total of 30/1,959 (1.5%) non-U.S. patients receiving zolpidem tartrate reported falls, including 28/30 (93%) who were ≥ 70 years of age. Of these 28 patients, 23 (82%) were receiving zolpidem doses > 10 mg. A total of 24/1,959 (1.2%) non-U.S. patients receiving zolpidem reported confusion, including 18/24 (75%) who were ≥ 70 years of age. Of these 18 patients, 14 (78%) were receiving zolpidem doses > 10 mg.

The dose of zolpidem tartrate sublingual tablets in elderly patients is 5 mg to minimize adverse effects related to impaired motor and/or cognitive performance and unusual sensitivity to sedative/hypnotic drugs [see *DOSAGE AND ADMINISTRATION (2)*, *WARNINGS AND PRECAUTIONS (5) CLINICAL PHARMACOLOGY (12)* and *CLINICAL STUDIES (14)*].

8.6 Gender Difference in Pharmacokinetics

Women clear zolpidem tartrate from the body at a lower rate than men, C_{max} and AUC parameters of zolpidem were approximately 45% higher at the same dose in female subjects compared with male subjects. Given the higher blood levels of zolpidem tartrate in women compared to men at a given dose, the recommended dose of zolpidem tartrate for adult women is 5 mg, and the recommended dose for adult men is 5 or 10 mg.

In geriatric patients, clearance of zolpidem is similar in men and women. The recommended dose of zolpidem tartrate in geriatric patients is 5 mg regardless of gender.

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Zolpidem tartrate sublingual tablets contains the same active substance, zolpidem tartrate, as zolpidem tartrate oral tablets and is classified as a Schedule IV controlled substance by federal regulation.

9.2 Abuse

Abuse and addiction are separate and distinct from physical dependence and tolerance. Abuse is characterized by misuse of the drug for non-medical purposes, often in combination with other

psychoactive substances. Tolerance is a state of adaptation in which exposure to a drug induces changes that result in a diminution of one or more of the drug effects over time. Tolerance may occur to both desired and undesired effects of drugs and may develop at different rates for different effects.

Addiction is a primary, chronic, neurobiological disease with genetic, psychosocial, and environmental factors influencing its development and manifestations. It is characterized by behaviors that include one or more of the following: impaired control over drug use, compulsive use, continued use despite harm, and craving. Drug addiction is a treatable disease, using a multidisciplinary approach, but relapse is common.

Studies of abuse potential in former drug abusers found that the effects of single doses of zolpidem tartrate 40 mg were similar, but not identical, to diazepam 20 mg, while zolpidem tartrate 10 mg was difficult to distinguish from placebo.

Because persons with a history of addiction to or abuse of, drugs or alcohol are at increased risk for misuse, abuse, and addiction of zolpidem tartrate, they should be monitored carefully when receiving zolpidem tartrate or any other hypnotic.

9.3 Dependence

Physical dependence is a state of adaptation that is manifested by a specific withdrawal syndrome that can be produced by abrupt cessation, rapid dose reduction, decreasing blood level of the drug, and/or administration of an antagonist.

Sedative/hypnotics have produced withdrawal signs and symptoms following abrupt discontinuation. These reported symptoms range from mild dysphoria and insomnia to a withdrawal syndrome that may include abdominal and muscle cramps, vomiting, sweating, tremors, and convulsions. The following adverse events which are considered to meet the DSM-III-R criteria for uncomplicated sedative/hypnotic withdrawal were reported during U.S. clinical trials following placebo substitution occurring within 48 hours following last zolpidem tartrate treatment: fatigue, nausea, flushing, lightheadedness, uncontrolled crying, emesis, stomach cramps, panic attack, nervousness, and abdominal discomfort. These reported adverse events occurred at an incidence of 1% or less. However, available data cannot provide a reliable estimate of the incidence, if any, of dependence during treatment at recommended doses. Post-marketing reports of abuse, dependence and withdrawal have been received.

10 OVERDOSAGE

10.1 Signs and Symptoms

In post-marketing experience of overdose with zolpidem tartrate alone, or in combination with CNS-depressant agents, impairment of consciousness ranging from somnolence to coma, cardiovascular and/or respiratory compromise, and fatal outcomes have been reported.

10.2 Recommended Treatment


Based on data obtained for zolpidem tartrate, general symptomatic and supportive measures for overdose with zolpidem tartrate should be used along with immediate gastric lavage where appropriate. Intravenous fluids should be administered as needed. Zolpidem's sedative/hypnotic effect was shown to be reduced by flumazenil and therefore may be useful; however, flumazenil administration may contribute to the appearance of neurological symptoms (convulsions). As in all cases of drug overdose, respiration, pulse, blood pressure, and other appropriate signs should be monitored and general supportive measures employed. Hypotension and CNS depression should be monitored and treated by appropriate medical intervention. Sedating drugs should be withheld following zolpidem overdose,

even if excitation occurs. The value of dialysis in the treatment of overdose has not been determined, although hemodialysis studies in patients with renal failure receiving therapeutic doses have demonstrated that zolpidem is not dialyzable.

As with the management of all overdose, the possibility of multiple drug ingestion should be considered. The physician may wish to consider contacting a poison control center for up-to-date information on the management of hypnotic drug product overdose.

11 DESCRIPTION

Zolpidem Tartrate Sublingual Tablet is a non-benzodiazepine hypnotic of the imidazopyridine class and is available in 5 mg and 10 mg strength tablets for sublingual administration. Chemically, zolpidem tartrate is *N,N,6-trimethyl-2-p-tolylimidazo[1,2-a]pyridine-3-acetamide L-(+)-tartrate* (2:1). It has the following structure:



Zolpidem tartrate is a white to off-white crystalline powder that is sparingly soluble in water, alcohol, and propylene glycol. It has a molecular weight of 764.88.

Each zolpidem tartrate sublingual tablet includes the following inactive ingredients: colloidal silicon dioxide, croscopovidone saccharin sodium, magnesium stearate, corn starch, mannitol and pregelatinized starch.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Zolpidem, the active moiety of zolpidem tartrate, is a hypnotic agent with a chemical structure unrelated to benzodiazepines, barbiturates, or other drugs with known hypnotic properties. It interacts with a GABA-BZ receptor complex and shares some of the pharmacological properties of the benzodiazepines. In contrast to the benzodiazepines, which non-selectively bind to and activate all BZ receptor subtypes, zolpidem *in vitro* binds the BZ₁ receptor preferentially with a high affinity ratio of the α_1/α_5 subunits. This selective binding of zolpidem on the BZ₁ receptor is not absolute, but it may explain the relative absence of myorelaxant and anticonvulsant effects in animal studies as well as the preservation of deep sleep (stages 3 and 4) in human studies of zolpidem tartrate at hypnotic doses.

12.3 Pharmacokinetics

Absorption:

Zolpidem tartrate sublingual tablets are bioequivalent to Ambien[®] tablets (Sanofi-Aventis) with respect to C_{max} and AUC. Similar to zolpidem tartrate oral tablets, zolpidem tartrate sublingual tablets result in a pharmacokinetic profile characterized by rapid absorption.

Following administration of single 10 mg zolpidem tartrate sublingual tablets, in 18 healthy adult subjects (18 to 65 years of age), the mean peak concentration (C_{max}) of zolpidem was 106 ng/mL (range:

52 to 205 ng/mL) occurring at a median time (T_{\max}) of 82 minutes (range: 30 to 180 min).

A food-effect study in 18 healthy volunteers compared the pharmacokinetics of zolpidem tartrate 10 mg when administered while fasting or within 20 minutes after a high fat meal. The mean AUC and C_{\max} were decreased by 20% and 31%, respectively, while median T_{\max} was prolonged by 28% (from 82 to 105 min). The half-life remained unchanged. These results suggest that, for faster sleep onset, zolpidem tartrate should not be administered with or immediately after a meal.

Distribution:

Based on data obtained with oral zolpidem, the total protein binding was found to be $92.5 \pm 0.1\%$ and remained constant, independent of concentration between 40 and 790 ng/mL.

Metabolism:

Based on data obtained with oral zolpidem, zolpidem is converted to inactive metabolites that are eliminated primarily by renal excretion.

Elimination:

When zolpidem tartrate administered as a single 5 or 10 mg dose in healthy adult subjects, the mean zolpidem elimination half-life was 2.85 hours (range: 1.57 to 6.73 hr) and 2.65 hours (range: 1.75 to 3.77 hr) respectively.

Special Populations

Elderly:

In the elderly, the dose for zolpidem tartrate should be 5 mg [see **WARNINGS AND PRECAUTIONS (5) and DOSAGE AND ADMINISTRATION (2)**]. This recommendation is based on several studies with zolpidem tartrate in which the mean C_{\max} , $T_{1/2}$, and AUC were significantly increased when compared to results in young adults. In one study of eight elderly subjects (>70 years), the means for C_{\max} , $T_{1/2}$, and AUC significantly increased by 50% (255 vs. 384 ng/mL), 32% (2.2 vs. 2.9 hr), and 64% (955 vs. 1,562 ng•hr/mL), respectively, as compared to younger adults (20 to 40 years) following a single 20 mg oral dose. Zolpidem did not accumulate in elderly subjects following nightly oral dosing of 10 mg for 1 week.

Hepatic Impairment:

The pharmacokinetics of zolpidem tartrate in eight patients with chronic hepatic insufficiency were compared to results in healthy subjects. Following a single 20 mg oral zolpidem tartrate dose, mean C_{\max} and AUC were found to be two times (250 vs. 499 ng/mL) and five times (788 vs. 4,203 ng•hr/mL) higher, respectively, in hepatically-compromised patients. T_{\max} did not change. The mean half-life in cirrhotic patients of 9.9 hr (range: 4.1 to 25.8 hr) was greater than that observed in normals of 2.2 hr (range: 1.6 to 2.4 hr). Dosing with zolpidem tartrate sublingual tablets should be modified accordingly in patients with hepatic insufficiency [see **Dosage and Administration (2.2)**].

Renal Impairment:

The pharmacokinetics of zolpidem tartrate were studied in 11 patients with end-stage 4 renal failure (mean $Cl_{Cr} = 6.5 \pm 1.5$ mL/min) undergoing hemodialysis three times a week, who were dosed with zolpidem tartrate 10 mg orally each day for 14 or 21 days. No statistically significant differences were

observed for C_{\max} , T_{\max} , half-life, and AUC between the first and last day of drug administration when baseline concentration adjustments were made. Zolpidem was not hemodialyzable. No accumulation of unchanged drug appeared after 14 or 21 days. Zolpidem pharmacokinetics were not significantly different in renally-impaired patients. No dosage adjustment is necessary in patients with compromised renal function.

Drug Interactions

CNS-depressants:

Coadministration of zolpidem with other CNS depressants increases the risk of CNS depression [see *Warnings and Precautions (5.1)*]. Zolpidem tartrate was evaluated in healthy volunteers in single-dose interaction studies for several CNS drugs. Imipramine in combination with zolpidem produced no pharmacokinetic interaction other than a 20% decrease in peak levels of imipramine, but there was an additive effect of decreased alertness. Similarly, chlorpromazine in combination with zolpidem produced no pharmacokinetic interaction, but there was an additive effect of decreased alertness and psychomotor performance.

A study involving haloperidol and zolpidem revealed no effect of haloperidol on the pharmacokinetics or pharmacodynamics of zolpidem. The lack of a drug interaction following single-dose administration does not predict the absence of an effect following chronic administration.

An additive adverse effect on psychomotor performance between alcohol and oral zolpidem was demonstrated [see *Warnings and Precautions (5.1)*].

Following five consecutive nightly doses at bedtime of oral zolpidem tartrate 10 mg in the presence of sertraline 50 mg (17 consecutive daily doses, at 7:00 am, in healthy female volunteers), zolpidem C_{\max} was significantly higher (43%) and T_{\max} was significantly decreased (-53%). Pharmacokinetics of sertraline and N-desmethylsertraline were unaffected by zolpidem.

A single-dose interaction study with zolpidem tartrate 10 mg and fluoxetine 20 mg at steady-state levels in male volunteers did not demonstrate any clinically significant pharmacokinetic or pharmacodynamics interactions. When multiple doses of zolpidem and fluoxetine were given at steady-state and the concentrations evaluated in healthy females, an increase in the zolpidem half-life (17%) was observed. There was no evidence of an additive effect in psychomotor performance.

Drugs that Affect Drug metabolism via Cytochrome P450

Some compounds known to inhibit CYP3A may increase exposure to zolpidem. The effect of inhibitors of other P450 enzymes on the pharmacokinetics of zolpidem is unknown.

A single-dose interaction study with zolpidem tartrate 10 mg and itraconazole 200 mg at steady-state levels in male volunteers resulted in a 34% increase in $AUC_{0-\infty}$ of zolpidem tartrate. There were no pharmacodynamics effects of zolpidem detected on subjective drowsiness, postural sway, or psychomotor performance.

A single-dose interaction study with zolpidem tartrate 10 mg and rifampin 600 mg at steady-state levels in female subjects showed significant reductions of the AUC (-73%), C_{\max} (-58%), and $T_{1/2}$ (-36%) of zolpidem together with significant reductions in the pharmacodynamics effects of zolpidem tartrate. Rifampin, a CYP3A4 inducer, significantly reduced the exposure to and the pharmacodynamics effects of zolpidem.

A single-dose interaction study with zolpidem 5 mg and ketoconazole, a potent CYP3A4 inhibitor, given as 200 mg twice daily for 2 days increased C_{\max} of zolpidem (30%) and the total AUC of zolpidem (70%) compared to zolpidem alone and prolonged the elimination half-life (30%) along with an increase

in the pharmacodynamics effects of zolpidem. Consideration should be given to using a lower dose of zolpidem when ketoconazole and zolpidem are given together.

Other Drugs with No Interactions with Zolpidem

A study involving cimetidine/zolpidem tartrate and ranitidine/zolpidem tartrate combinations revealed no effect of either drug on the pharmacokinetics or pharmacodynamics of zolpidem.

Zolpidem tartrate had no effect on digoxin pharmacokinetics and did not affect prothrombin time when given with warfarin in healthy subjects.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis & Mutagenesis & Impairment Of Fertility

Carcinogenesis:

Zolpidem was administered to mice and rats for 2 years at oral doses of 4, 18, and 80 mg base/kg. In mice, these doses are approximately 2.5, 10, and 50 times the maximum recommended human dose (MRHD) of 10 mg/day (8 mg zolpidem base) on a mg/m² basis. In rats, these doses are approximately 5, 20, and 100 times the MRHD on a mg/m² basis. No evidence of carcinogenic potential was observed in mice. In rats, renal tumors (lipoma, liposarcoma) were seen at the mid- and high doses.

Mutagenesis:

Zolpidem was negative in *in vitro* (bacterial reverse mutation, mouse lymphoma, and chromosomal aberration) and *in vivo* (mouse micronucleus) genetic toxicology assays.

Impairment of fertility:

Oral administration of zolpidem (doses of 4, 20, and 100 mg base/kg to rats prior to and during mating, and continuing in females through postpartum day 25, resulted in irregular estrus cycles and prolonged pre-coital intervals at the highest dose tested. The no-effect dose for these findings is approximately 24 times the MRHD on a mg/m² basis. There was no impairment of fertility at any dose tested.

14 CLINICAL STUDIES

14.1 Transient Insomnia

Normal adults experiencing transient insomnia (n = 462) during the first night in a sleep laboratory were evaluated in a double-blind, parallel group, single night trial comparing two doses of zolpidem tartrate oral tablets (7.5 and 10 mg) and placebo. Both zolpidem doses were superior to placebo on objective (polysomnographic) measures of sleep latency, sleep duration, and number of awakenings.

Normal elderly adults (mean age 68) experiencing transient insomnia (n = 35) during the first two nights in a sleep laboratory were evaluated in a double-blind, crossover, 2-night trial comparing four doses of zolpidem (5, 10, 15, and 20 mg) and placebo. All zolpidem doses were superior to placebo on the two primary PSG parameters (sleep latency and efficiency) and all four subjective outcome measures (sleep duration, sleep latency, number of awakenings, and sleep quality).

14.2 Chronic Insomnia

Zolpidem was evaluated in two controlled studies for the treatment of patients with chronic insomnia (most closely resembling primary insomnia, as defined in the APA Diagnostic and Statistical Manual of Mental Disorders, DSM-IV™). Adult outpatients with chronic insomnia (n = 75) were evaluated in a

double-blind, parallel group, 5-week trial comparing two doses of zolpidem tartrate and placebo. On objective (polysomnographic) measures of sleep latency and sleep efficiency, zolpidem 10 mg was superior to placebo on sleep latency for the first 4 weeks and on sleep efficiency for weeks 2 and 4. Zolpidem was comparable to placebo on number of awakenings at both doses studied.

Adult outpatients (n=141) with chronic insomnia were also evaluated, in a double-blind, parallel group, 4-week trial comparing two doses of zolpidem and placebo. Zolpidem 10 mg was superior to placebo on a subjective measure of sleep latency for all 4 weeks, and on subjective measures of total sleep time, number of awakenings, and sleep quality for the first treatment week.

Increased wakefulness during the last third of the night as measured by polysomnography has not been observed in clinical trials with zolpidem tartrate.

14.3 Studies Pertinent to Safety Concerns for Sedative/Hypnotic Drugs

Next-day residual effects:

Next-day residual effects of zolpidem tartrate were evaluated in seven studies involving normal subjects. In three studies in adults (including one study in a phase advance model of transient insomnia) and in one study in elderly subjects, a small but statistically significant decrease in performance was observed in the Digit Symbol Substitution Test (DSST) when compared to placebo. Studies of zolpidem tartrate in non-elderly patients with insomnia did not detect evidence of next-day residual effects using the DSST, the Multiple Sleep Latency Test (MSLT), and patient ratings of alertness.

Rebound effects:

There was no objective (polysomnographic) evidence of rebound insomnia at recommended doses seen in studies evaluating sleep on the nights following discontinuation of zolpidem tartrate. There was subjective evidence of impaired sleep in the elderly on the first post-treatment night at doses of zolpidem tartrate above the recommended elderly dose of 5 mg.

Memory impairment:

Controlled studies in adults utilizing objective measures of memory yielded no consistent evidence of next-day memory impairment following the administration of zolpidem tartrate. However, in one study involving zolpidem doses of 10 and 20 mg, there was a significant decrease in next-morning recall of information presented to subjects during peak drug effect (90 minutes post-dose), i.e., these subjects experienced anterograde amnesia. There was also subjective evidence from adverse event data for anterograde amnesia occurring in association with the administration of zolpidem tartrate, predominantly at doses above 10 mg.

Effects on sleep stages:

In studies that measured the percentage of sleep time spent in each sleep stage, zolpidem tartrate has generally been shown to preserve sleep stages. Sleep time spent in stages 3 and 4 (deep sleep) was found comparable to placebo with only inconsistent, minor changes in REM (paradoxical) sleep at the recommended dose.

16 HOW SUPPLIED/STORAGE AND HANDLING

Zolpidem tartrate sublingual tablets are supplied in two dosage strengths: Tablets are not scored.

Zolpidem tartrate sublingual tablets 5 mg are white, round tablets, flat-faced, bevel-edged with debossed "P1" on one side and plain on the other side and supplied as:

NDC Number	Size
49884-903-52	10 tablets in one blister card
49884-903-55	Carton of 30 tablets in a 3 blister cards

The blister packs consist of aluminum/aluminum Child Resistant Control (CRC) blisters.

Zolpidem tartrate sublingual tablets 10 mg are white, round tablets, flat-faced, bevel-edged with debossed "P7" on one side and plain on the other side and supplied as:

NDC Number	Size
49884-904-52	10 tablets in one blister card
49884-904-55	Carton of 30 tablets in a 3 blister cards

The blister packs consist of aluminum/aluminum Child Resistant Control (CRC) blisters.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Medication Guide).

Inform patients and their families about the benefits and risks of treatment with zolpidem tartrate. Inform patients of the availability of a Medication Guide and instruct them to read the Medication Guide prior to initiating treatment with zolpidem tartrate and with each prescription refill. Review the zolpidem tartrate Medication Guide with every patient prior to initiation of treatment. Instruct patients or caregivers that zolpidem tartrate should be taken only as prescribed.

CNS-depressant Effects and Next-Day Impairment

Tell patients that zolpidem tartrate has the potential to cause next-day impairment, and that this risk is increased if dosing instructions are not carefully followed. Tell patients to wait for at least 8 hours after dosing before driving or engaging in other activities requiring full mental alertness. Inform patients that impairment can be present despite feeling fully awake.

Severe Anaphylactic and Anaphylactoid Reactions

Inform patients that severe anaphylactic and anaphylactoid reactions have occurred with zolpidem. Describe the signs/symptoms of these reactions and advise patients to seek medical attention immediately if any of them occur.

Sleep-Driving and Other Complex Behaviors

Instruct patients and their families that sedative hypnotics can cause abnormal thinking and behavior change, including "sleep-driving" and other complex behaviors while not being fully awake (preparing and eating food, making phone calls, or having sex). Tell patients to call you immediately if they develop any of these symptoms.

Suicide

Tell patients to immediately report any suicidal thoughts.

Alcohol and Other Drugs

Ask patients about alcohol consumption, medicines they are taking, and drugs they may be taking without

a prescription. Advise patients not to use zolpidem tartrate if they drank alcohol that evening or before bed.

Tolerance, Abuse, and Dependence

Tell patients not to increase the dose of zolpidem tartrate on their own, and to inform you if they believe the drug “does not work”.

Administration Instructions

Patients should be counseled to take zolpidem tartrate right before they get into bed and only when they are able to stay in bed a full night (7 to 8 hours) before being active again. Zolpidem tartrate tablets should not be taken with or immediately after a meal. Advise patients NOT to take zolpidem tartrate when drinking alcohol that evening or before bed. Zolpidem tartrate sublingual tablet should be placed under the tongue, where it will disintegrate. The tablet should not be swallowed and the tablet should not be taken with water.

MEDICATION GUIDE

Medication Guide

Zolpidem Tartrate (Zole-PI-dem TAR-trate) Sublingual Tablets C-IV

Read this Medication Guide that comes with zolpidem tartrate sublingual tablets before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment.

What is the most important information I should know about zolpidem tartrate sublingual tablets?

- **Do not take more zolpidem tartrate sublingual tablets than prescribed.**
- **Do not take zolpidem tartrate sublingual tablets unless you are able to stay in bed a full night (7 to 8 hours) before you must be active again.**
- **Take zolpidem tartrate sublingual tablets right before you get in bed, not sooner**

Zolpidem tartrate sublingual tablets may cause serious side effects, including:

- **After taking zolpidem tartrate sublingual tablets, you may get up out of bed while not being fully awake and do an activity that you do not know you are doing. The next morning, you may not remember that you did anything during the night.**

You have a higher chance for doing these activities if you drink alcohol or take other medicines that make you sleepy with zolpidem tartrate sublingual tablets. Reported activities include:

- driving a car (“sleep-driving”)
- making and eating food
- talking on the phone
- having sex
- sleep-walking

Call your healthcare provider right away if you find out that you have done any of the above activities after taking zolpidem tartrate sublingual tablets.

Do not take zolpidem tartrate sublingual tablets if you:

- drank alcohol that evening or before bed
- took another medicine to help you sleep

What is zolpidem tartrate sublingual tablets?

Zolpidem tartrate sublingual tablets is a sedative-hypnotic (sleep) medicine. Zolpidem tartrate sublingual

tablets is used in adults for the short-term treatment of a sleep problem called insomnia (trouble falling asleep).

It is not known if zolpidem tartrate sublingual tablets is safe and effective in children under the age of 18 years.

Zolpidem tartrate is a class four (C-IV) federally controlled substance because it can be abused or lead to dependence. Keep zolpidem tartrate sublingual tablets in a safe place to prevent misuse and abuse. Selling or giving away zolpidem tartrate sublingual tablets may harm others, and is against the law. Tell your doctor if you have ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take zolpidem tartrate sublingual tablets?

- Do not take zolpidem tartrate sublingual tablets if you are allergic to zolpidem or any other ingredients in zolpidem tartrate sublingual tablets. See the end of this Medication Guide for a complete list of ingredients in zolpidem tartrate sublingual tablets.
- Do not take zolpidem tartrate sublingual tablets if you have had an allergic reaction to drugs containing zolpidem, such as Ambien, Ambien CR, Zolpimist, or Intermezzo. Symptoms of a serious allergic reaction to zolpidem can include:
 - swelling of your face, lips, and throat that may cause difficulty breathing or swallowing
 - nausea and vomiting

What should I tell my healthcare provider before taking zolpidem tartrate sublingual tablets?

Zolpidem tartrate sublingual tablets may not be right for you. Before starting zolpidem tartrate sublingual tablets, tell your healthcare provider about all of your health conditions, including if you:

- have a history of depression, mental illness or, suicidal thoughts
- have a history of drug or alcohol abuse or addiction
- have kidney or liver disease
- have lung disease or breathing problems
- are pregnant or planning to become pregnant. It is not known if zolpidem tartrate sublingual tablets will harm your unborn baby.
- are breastfeeding or plan to breastfeed. Zolpidem tartrate can pass into your breast milk. It is not known if zolpidem tartrate will harm your baby. Talk to your healthcare provider about the best way to feed your baby while you take zolpidem tartrate sublingual tablets.

Tell your healthcare provider about all of the medicines you take, including prescription and nonprescription medicines, vitamins and herbal supplements.

Medicines can interact with each other, sometimes causing serious side effects.

Do not take zolpidem tartrate sublingual tablets with other medicines that can make you sleepy, unless directed by your healthcare provider

Know the medicines you take. Keep a list of your medicines with you to show your healthcare provider and pharmacist each time you get a new medicine.

How should I take zolpidem tartrate sublingual tablets?

- See **“What is the most important information I should know about zolpidem tartrate sublingual tablets?”**
- Take zolpidem tartrate sublingual tablets exactly as prescribed. Only take 1 zolpidem tartrate sublingual tablet a night and only if needed.
- Do not take zolpidem tartrate sublingual tablets if you drank alcohol that evening or before bed.
- You should not take zolpidem tartrate sublingual tablets with or right after a meal. Zolpidem tartrate sublingual tablets may help you fall asleep faster if you take it on an empty stomach.
- Do not use the tablet if the seal on the childproof blister pack is broken, or if the blister holding the tablet is broken.
- To open the blister pack, separate the individual blisters at the perforations. Peel off the top layer of

paper, and push the tablet through the foil. Alternatively, use scissors to open the blister.

- Place the tablet under the tongue, where it will disintegrate. Do not swallow or take with water.
- Call your healthcare provider if your insomnia worsens or is not better within 7 to 10 days. This may mean that there is another condition causing your sleep problem.
- If you take too much zolpidem tartrate sublingual tablets or overdose, get emergency treatment.

What are the possible side effects of zolpidem tartrate sublingual tablets?

Zolpidem tartrate sublingual tablets may cause serious side effects, including:

- **getting out of bed while not being fully awake and doing an activity that you do not know you are doing.** See "What is the most important information I should know about zolpidem tartrate sublingual tablets?"
- **abnormal thoughts and behavior.** Symptoms include more outgoing or aggressive behavior than normal, confusion, agitation, hallucinations, worsening of depression, suicidal thoughts or actions.
- **memory loss**
- **anxiety**
- **severe allergic reactions.** Symptoms include swelling of the tongue or throat, trouble breathing. Get emergency medical help if you get these symptoms after taking zolpidem tartrate sublingual tablets.

Call your healthcare provider right away if you have any of the above side effects or any other side effects that worry you while using zolpidem tartrate sublingual tablets.

The most common side effects of zolpidem tartrate sublingual tablets are:

- drowsiness
- dizziness
- diarrhea
- grogginess or feeling as if you have been drugged
- fatigue
- headache

You may still feel drowsy the next day after taking zolpidem tartrate sublingual tablets.

After you stop taking a sleep medicine, you may have symptoms for 1 to 2 days such as:

- trouble sleeping
- nausea
- flushing
- lightheadedness
- uncontrolled crying
- vomiting
- stomach cramps
- panic attack
- nervousness
- stomach area pain

These are not all the side effects of zolpidem tartrate sublingual tablets. Ask your doctor or pharmacist for more information.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store zolpidem tartrate sublingual tablets?

- Store zolpidem tartrate sublingual tablets at room temperature, between 68° and 77°F (20° to 25°C).
- Protect from light and moisture.

Keep zolpidem tartrate sublingual tablets and all medicines out of reach of children.

General information about the safe and effective use of zolpidem tartrate sublingual tablets

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide.

Do not use zolpidem tartrate sublingual tablets for a condition for which it was not prescribed.

Do not share zolpidem tartrate sublingual tablets with other people, even if you think they have the same symptoms that you have. It may harm them and it is against the law.

This Medication Guide summarizes the most important information about zolpidem tartrate sublingual tablets.

If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about zolpidem tartrate sublingual tablets that is written for healthcare professionals.

For more information about zolpidem tartrate sublingual tablets, call Par Pharmaceutical. at 1-800-828-9393.

What are the ingredients in zolpidem tartrate sublingual tablets?

Active Ingredient: zolpidem tartrate

Inactive ingredients: colloidal silicon dioxide, crospovidone, corn starch, magnesium stearate, mannitol, pregelatinized starch and saccharin sodium.

This Medication Guide has been approved by U.S. Food and Drug Administration.

All brand names listed are the registered trademarks of their respective owners and are not trademarks of Par Pharmaceutical.

Rx only

Dist. by:

Par Pharmaceutical

Chestnut Ridge, NY 10977 U.S.A.

Mfg. by:

Par Formulations Private Limited,

9/215, Pudupakkam, Kelambakkam - 603 103.

Made in India

Mfg. Lic. No.: TN00002121

OS903-01-74-02

Revised: 05/2017

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**

5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01

NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**

5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01



NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**

5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01

NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**

5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01



NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**

5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01

NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**

5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01



NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**

5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01

NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**


5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01



NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**

5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01

NDC 49884-903-52
**Zolpidem Tartrate
Sublingual Tablet**

5 mg 

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.
Made in India
Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL903-54-74-01

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

Peel and
push

Peel and
push

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

Peel and
push

Peel and
push

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

Peel and
push

Peel and
push

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

Peel and
push

Peel and
push

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

NDC 49884-904-52
**Zolpidem Tartrate
Sublingual Tablet**

10 mg 

Dist. by: Par Pharmaceutical
Chestnut Ridge, NY 10977 U.S.A.
Made in India

Mfg. Lic. No.: TN00002121



Space for Lot & Exp

BL904-54-74-01

NDC 49884-903-55

Zolpidem Tartrate Sublingual Tablets

5 mg



Rx Only
30 Sublingual Tablets

NDC 49884-903-55

Zolpidem Tartrate Sublingual Tablets

5 mg



Attention Pharmacist: Federal Law requires dispensing the accompanying Medication Guide to each patient.

Rx Only

30 Sublingual Tablets



NDC 49884-903-55
Zolpidem Tartrate Sublingual Tablets
5 mg
Rx Only
30 Sublingual Tablets

Un varnished Area

Each sublingual tablet (to be placed under the tongue) contains 5 mg of zolpidem tartrate.

Usual Dosage: See prescribing information for dosage information.

KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN.

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature]. Protect from light and moisture.
Do not use if blisters are torn, or seal is broken.

Dist. by: **Par Pharmaceutical**
Chestnut Ridge, NY 10977 U.S.A.

Mfg. by: **Par Formulations Private Limited,**
9/215, Pudupakkam, Kelambakkam - 603103.
Made in India

Mfg. Lic. No.: TN00002121
UC903-55-74-02
R05/17



Zolpidem Tartrate Sublingual Tablets
5 mg
Rx Only
30 Sublingual Tablets
NDC 49884-903-55



ZOLPIDEM TARTRATE

zolpidem tartrate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:49884-903
Route of Administration	SUBLINGUAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZOLPIDEM TARTRATE (UNII: WY6W63843K) (ZOLPIDEM - UNII:7K3830Q123)	ZOLPIDEM TARTRATE	5 mg

Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 2S7830E561)	
SACCHARIN SODIUM ANHYDROUS (UNII: I4807BK602)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND (white round tablets, flat faced, bevel-edged tablet)	Size	7mm
Flavor		Imprint Code	P1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49884-903-52	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	11/04/2016	
2	NDC:49884-903-55	30 in 1 CARTON; Type 0: Not a Combination Product	11/04/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201509	11/04/2016	

ZOLPIDEM TARTRATE

zolpidem tartrate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:49884-904
Route of Administration	SUBLINGUAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZOLPIDEM TARTRATE (UNII: WY6W63843K) (ZOLPIDEM - UNII:7K383OQI23)	ZOLPIDEM TARTRATE	10 mg

Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 2S7830E561)	
SACCHARIN SODIUM ANHYDROUS (UNII: I4807BK602)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND (white, round tablets, flat-faced, bevel-edged tablet)	Size	7mm
Flavor		Imprint Code	P7
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49884-904-52	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	11/04/2016	
2	NDC:49884-904-55	30 in 1 CARTON; Type 0: Not a Combination Product	11/04/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201509	11/04/2016	

Labeler - Par Pharmaceutical, Inc. (092733690)

Revised: 11/2017

Par Pharmaceutical, Inc.