

**CHILDRENS BENADRYL ALLERGY PLUS CONGESTION- diphenhydramine hydrochloride and phenylephrine hydrochloride solution**  
Kenvue Brands LLC

-----  
**Childrens Benadryl Allergy Plus Congestion**

***Drug Facts***

<b><i>Active ingredients (in each 5 mL)</i></b>	<b><i>Purposes</i></b>
Diphenhydramine HCl 12.5 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
  - nasal congestion
  - stuffy nose
- temporarily relieves these symptoms due to the common cold:
  - runny nose
  - sneezing
  - nasal congestion
  - stuffy nose
- temporarily relieves sinus congestion and pressure

**Warnings**

**Do not use**

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes

- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dose**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- find right dose on chart below
- mL = milliliter
- take every 4 hours
- do not take more than 6 doses in 24 hours

<b>Age (yr)</b>	<b>Dose (mL)</b>
children under 4 years	do not use
children 4 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL
adults and children 12 years and over	10 mL

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other information**

- each 5 mL contains: **sodium 10 mg**
- store between 20-25°C (68-77°F)
- **do not use if carton tape or bottle wrap imprinted with "Sealed For Your Safety" is broken or missing**

## **Inactive ingredients**

anhydrous citric acid, carboxymethylcellulose sodium, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

## **Questions or comments?**

call **1-877-717-2824** (toll-free) or **215-273-8755** (collect)

## **PRINCIPAL DISPLAY PANEL**

NDC 50580-341-01

Children's  
Benadryl®

ALLERGY PLUS  
CONGESTION

Diphenhydramine HCl / antihistamine  
Phenylephrine HCl / nasal decongestant  
oral solution

4 Hours/Dose

RELIEF OF:

- Runny Nose
- Sneezing
- Itchy, Watery Eyes
- Itchy Throat or Nose
- Sinus Congestion
- Stuffy Nose

Alcohol Free  
Sugar Free

ALLERGY + CONGESTION RELIEF

Grape!  
Flavored Liquid

4 fl oz (118 mL)

Children's  
**Benadryl**<sup>®</sup>

**ALLERGY PLUS**  
**CONGESTION**

How can we help?  
☎ **1-877-717-2824**  
benadryl.com

NDC 50580-341-01

Children's  
**Benadryl**<sup>®</sup>

**ALLERGY PLUS**  
**CONGESTION**

Diphenhydramine HCl / antihistamine  
Phenylephrine HCl / nasal decongestant  
oral solution

**4 Hours/Dose**

**RELIEF OF:**

- ✔ Runny Nose
- ✔ Sneezing
- ✔ Itchy, Watery Eyes
- ✔ Itchy Throat or Nose
- ✔ Sinus Congestion
- ✔ Stuffy Nose

Alcohol Free  
Sugar Free



Actual Size

**Tips for Outdoor Allergens:**



- Avoid being outside when the pollen count is high, typically in the morning
- Shower and change clothing after spending time outdoors

**Tips for Indoor Allergens:**



- Wash sheets and blankets in hot water and use the dryer weekly



• Minimize contact and keep pets out of the bedroom to help with pet allergies



**Drug Facts (continued)**

- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

---

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

---

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help

Children's  
**Benadryl**<sup>®</sup>

**ALLERGY PLUS**

# CONGESTION

## Drug Facts

Active ingredients (in each 5 mL)	Purposes
Diphenhydramine HCl 12.5 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

## Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
  - nasal congestion
  - stuffy nose
- temporarily relieves these symptoms due to the common cold:
  - runny nose
  - sneezing
  - nasal congestion
  - stuffy nose
- temporarily relieves sinus congestion and pressure

## Warnings

### Do not use

- to make a child sleepy
- with any other product containing diphenhydramine, even one used on skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

### When using this product

- do not exceed recommended dose
- marked drowsiness may occur
- avoid alcoholic drinks

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- find right dose on chart below
- mL = milliliter
- take every 4 hours
- do not take more than 6 doses in 24 hours

Age (yr)	Dose (mL)
children under 4 years	do not use
children 4 to 5 years	do not use unless directed by a doctor
children 6 to 11 years	5 mL
adults and children 12 years and over	10 mL

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

## Other information

- each 5 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F)
- do not use if carton tape or bottle wrap imprinted with "Sealed For Your Safety" is broken or missing

## Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

## Questions or comments?

call 1-877-717-2824 (toll-free) or 215-273-8755 (collect)

Distributed by:

**JOHNSON & JOHNSON CONSUMER INC.**

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

©J&JCI 2019

30046178

For more information, visit:  
**www.benadryl.com**

# CHILDRENS BENADRYL ALLERGY PLUS CONGESTION

diphenhydramine hydrochloride and phenylephrine hydrochloride solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50580-341
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DIPHENHYDRAMINE HYDROCHLORIDE</b> (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

## Product Characteristics

<b>Color</b>	purple	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-341-01	1 in 1 CARTON	09/30/2014	
1		118 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/30/2014	

**Labeler** - Kenvue Brands LLC (118772437)

Revised: 11/2024

Kenvue Brands LLC