

**ASPIRIN- aspirin tablet, film coated**  
**Geiss, Destin & Dunn Inc.**

-----  
**GoodSense 44-157**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

**Inactive ingredients**

corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?**

**1-800-426-9391**

**Principal display panel**

**GoodSense®**

NDC 50804-157-12

**Aspirin**

*Pain Reliever/Fever Reducer*  
**(NSAID)**

**Safe Pain Relief**

*actual size*

†Compare to the active ingredient of  
Genuine Bayer® Aspirin

**100** Tablets - **325 mg** each

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY  
SEAL UNDER CAP IS BROKEN OR MISSING

†This product is not manufactured or distributed by Bayer AG, owner of the registered  
trademark Genuine Bayer® Aspirin.            50844            ORG012215712

Distributed by:

Perrigo Direct, Inc.,  
Peachtree City, GA 30269  
www.PerrigoDirect.com  
(1-800-426-9391)

GoodSense® is a registered  
trademark of L. Perrigo  
Company.

**GOODSENSE**® NDC 50804-157-12

# Aspirin

Pain Reliever/Fever Reducer  
(NSAID)

Safe Pain Relief actual size

**†Compare to the active ingredient of Genuine Bayer® Aspirin**

**100 Tablets - 325 mg each**

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine Bayer® Aspirin. 50844 ORG012215712

Distributed by:  
Perrigo Direct, Inc., 30269  
Peachtree City, GA 30269  
www.PerrigoDirect.com  
(1-800-426-9391)  
GoodSense® is a registered trademark of L. Perrigo Company.

8 46036 00955 2

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 325 mg (NSAID)\* ..... Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves: ■ minor pain of arthritis  
■ headache ■ muscle pain ■ menstrual pain  
■ toothache ■ pain and fever of colds

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling  
■ shock ■ hives ■ asthma (wheezing)  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug ■ have 3 or more alcoholic drinks every day while using this product  
■ have had stomach ulcers or bleeding problems  
■ take more or for a longer time than directed  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

No print/No varnish  
Lot & Exp date

**PEEL HERE FOR MORE DRUG FACTS**

**STOP PEELING**

**Drug Facts (continued)**

**Do not use** ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if** ■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools  
■ feel faint ■ have stomach pain that does not get better  
■ ringing in the ears or a loss of hearing occurs  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ do not take more than directed  
■ drink a full glass of water with each dose  
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor  
■ children under 12 years: do not use unless directed by a doctor

**Other information**  
■ use by expiration date on package  
■ store at 25°C (77°F); excursions permitted between 15-30°C (59-86°F)

**Inactive ingredients** corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?** 1-800-426-9391

GoodSense 44-157

## ASPIRIN

aspirin tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50804-157
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	

<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;157;ASPIRIN
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50804-157-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/12/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/12/2023	

**Labeler** - Geiss, Destin & Dunn Inc. (076059836)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50804-157)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50804-157)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50804-157)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50804-157)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50804-157)

