

PALONOSETRON - palonosetron injection, solution
Fresenius Kabi USA, LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PALONOSETRON HYDROCHLORIDE INJECTION safely and effectively. See full prescribing information for PALONOSETRON HYDROCHLORIDE INJECTION.

PALONOSETRON HYDROCHLORIDE Injection, for intravenous use
Initial U.S. Approval: 2003

----- **INDICATIONS AND USAGE** -----

Palonosetron Hydrochloride Injection is a serotonin-3 (5-HT₃) receptor antagonist indicated in:

Adults for prevention of:

- acute and delayed nausea and vomiting associated with initial and repeat courses of moderately emetogenic cancer chemotherapy (MEC). (1)
- acute nausea and vomiting associated with initial and repeat courses of highly emetogenic cancer chemotherapy (HEC). (1)

Pediatric patients aged 1 month to less than 17 years for prevention of:

- acute nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy, including highly emetogenic cancer chemotherapy (HEC). (1)

----- **DOSAGE AND ADMINISTRATION** -----

Chemotherapy-Induced Nausea and Vomiting (2.1)

Age	Dose*	Infusion Time
Adults	0.25 mg as a single dose	Infuse over 30 seconds beginning approximately 30 minutes before the start of chemotherapy
Pediatrics (1 month to less than 17 years)	20 micrograms per kilogram (maximum 1.5 mg) as a single dose	Infuse over 15 minutes beginning approximately 30 minutes before the start of chemotherapy

*Note different dosing units in pediatrics

Instructions for Intravenous Administration

- For a dose of 0.25 mg, use the entire contents (5 mL) of the prefilled syringe. Do not use the prefilled syringe to administer a dose of less than 0.25 mg (5 mL). (2.2)

----- **DOSAGE FORMS AND STRENGTHS** -----

0.25 mg (free base) per 5 mL (concentration: 0.05 mg per mL, 50 mcg per mL) single-dose prefilled syringe (3)

----- **CONTRAINDICATIONS** -----

Hypersensitivity to palonosetron or any of its components (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Hypersensitivity reactions, including anaphylaxis and anaphylactic shock; reported in patients with or without known hypersensitivity to other selective 5-HT₃ receptor antagonists. If symptoms occur, discontinue Palonosetron Hydrochloride Injection and initiate appropriate medical treatment. (5.1)
- Serotonin syndrome; reported with 5-HT₃ receptor antagonists alone, but particularly with concomitant use of serotonergic drugs. (5.2, 7.1)

----- **ADVERSE REACTIONS** -----

Most common adverse reactions in

- chemotherapy-induced nausea and vomiting in adults (≥5%) are: headache and constipation (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **DRUG INTERACTIONS** -----

Serotonergic Drugs: Monitor for serotonin syndrome; if symptoms occur, discontinue Palonosetron Hydrochloride Injection and initiate supportive treatment. (7.1)

FULL PRESCRIBING INFORMATION: CONTENTS*

1. INDICATIONS AND USAGE

2. DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosage
- 2.2 Instructions for Intravenous Administration

3. DOSAGE FORM AND STRENGTHS

4. CONTRAINDICATIONS

5. WARNINGS AND PRECAUTIONS

- 5.1 Hypersensitivity Reactions
- 5.2 Serotonin Syndrome

6. ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7. DRUG INTERACTIONS

- 7.1 Serotonergic Drugs

8. USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

10. OVERDOSAGE

11. DESCRIPTION

12. CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13. NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14. CLINICAL STUDIES

- 14.1 Prevention of Nausea and Vomiting Associated with MEC and HEC in Adults
- 14.2 Prevention of Nausea and Vomiting Associated with Emetogenic Chemotherapy, Including HEC in Pediatric Patients

16. HOW SUPPLIED/STORAGE AND HANDLING

17. PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1. INDICATIONS AND USAGE

Palonosetron Hydrochloride Injection is indicated in adults for prevention of:

- acute and delayed nausea and vomiting associated with initial and repeat courses of moderately emetogenic cancer chemotherapy (MEC).
- acute nausea and vomiting associated with initial and repeat courses highly emetogenic cancer

chemotherapy (HEC).

Palonosetron Hydrochloride Injection is indicated in pediatric patients 1 month to less than 17 years of age for prevention of:

- acute nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy, including highly emetogenic cancer chemotherapy.

2. DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

Prevention of Chemotherapy-Induced Nausea and Vomiting

The recommended dosage of Palonosetron Hydrochloride Injection for prevention of nausea and vomiting associated with HEC and MEC in adults and associated with emetogenic chemotherapy, including HEC in pediatric patients 1 month to less than 17 years of age is shown in Table 1.

Table 1: Recommended Dosage of Palonosetron Hydrochloride Injection for the Prevention of Nausea and Vomiting Associated with Chemotherapy in Adults and Pediatric Patients 1 Month to Less than 17 Years

Age	Dose*	Infusion Time
Adults	0.25 mg as a single dose	Infuse over 30 seconds beginning approximately 30 minutes before the start of chemotherapy
Pediatrics (1 month to less than 17 years)	20 micrograms per kilogram (max 1.5 mg) as a single dose	Infuse over 15 minutes beginning approximately 30 minutes before the start of chemotherapy

*Note different dosing units in pediatrics

2.2 Instructions for Intravenous Administration

- Palonosetron Hydrochloride Injection is supplied ready for intravenous administration at a concentration of 0.05 mg/mL (50 mcg/mL).
- Do not mix Palonosetron Hydrochloride Injection with other drugs.
- Flush the infusion line with normal saline before and after administration of Palonosetron Hydrochloride Injection.
- Inspect Palonosetron Hydrochloride Injection visually for particulate matter and discoloration before administration.
- Expel air from syringe prior to administration. For a dose of 0.25 mg, use the entire contents (5mL) of the prefilled syringe.
- Do not use the prefilled syringe to administer a dose less than 0.25mg (5mL)
- Use aseptic technique while handling the syringe

3. DOSAGE FORM AND STRENGTHS

Palonosetron Hydrochloride Injection is supplied as a single-dose sterile, clear, colorless solution in a prefilled syringe that provides:

- 0.25 mg (free base) per 5 mL (concentration: 0.05 mg per mL, 50 mcg per mL)

4. CONTRAINDICATIONS

Palonosetron Hydrochloride Injection is contraindicated in patients known to have hypersensitivity to

palonosetron [see *Warnings and Precautions (5.1)*].

5. WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

Hypersensitivity reactions, including anaphylaxis and anaphylactic shock, have been reported with administration of Palonosetron Hydrochloride Injection [see *Adverse Reactions (6.2)*]. These reactions occurred in patients with or without known hypersensitivity to other 5-HT₃ receptor antagonists. If hypersensitivity reactions occur, discontinue Palonosetron Hydrochloride Injection and initiate appropriate medical treatment. Do not reinitiate Palonosetron Hydrochloride Injection in patients who have previously experienced symptoms of hypersensitivity [see *Contraindications (4)*].

5.2 Serotonin Syndrome

The development of serotonin syndrome has been reported with 5-HT₃ receptor antagonists. Most reports have been associated with concomitant use of serotonergic drugs (e.g., selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), monoamine oxidase inhibitors, mirtazapine, fentanyl, lithium, tramadol, and intravenous methylene blue). Some of the reported cases were fatal. Serotonin syndrome occurring with overdose of another 5-HT₃ receptor antagonist alone has also been reported. The majority of reports of serotonin syndrome related to 5-HT₃ receptor antagonist use occurred in a post-anesthesia care unit or an infusion center.

Symptoms associated with serotonin syndrome may include the following combination of signs and symptoms: mental status changes (e.g. agitation, hallucinations, delirium, and coma), autonomic instability (e.g., tachycardia, labile blood pressure, dizziness, diaphoresis, flushing, hyperthermia), neuromuscular symptoms (e.g., tremor, rigidity, myoclonus, hyperreflexia, incoordination), seizures, with or without gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Patients should be monitored for the emergence of serotonin syndrome, especially with concomitant use of Palonosetron Hydrochloride Injection and other serotonergic drugs. If symptoms of serotonin syndrome occur, discontinue Palonosetron Hydrochloride Injection and initiate supportive treatment. Patients should be informed of the increased risk of serotonin syndrome, especially if Palonosetron Hydrochloride Injection is used concomitantly with other serotonergic drugs [see *Drug Interactions (7.1)*].

6. ADVERSE REACTIONS

Serious or otherwise clinically significant adverse reactions reported in other sections of labeling:

- Hypersensitivity Reactions [see *Warnings and Precautions (5.1)*]
- Serotonin Syndrome [see *Warnings and Precautions (5.2)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Chemotherapy-Induced Nausea and Vomiting

Adults

In double-blind randomized clinical trials for the prevention of nausea and vomiting induced by MEC or HEC, 1,374 adult patients received a single dose of Palonosetron Hydrochloride Injection, ondansetron (Studies 1 and 3) or dolasetron (Study 2) administered 30 minutes prior to chemotherapy [see *Clinical Studies (14.1)*].

Adverse reactions were similar in frequency and severity in all 3 treatment groups. Common adverse reactions reported in at least 2% of patients in these trials are shown in Table 2.

Table 2: Common Adverse Reactions* in Adults with Receiving MEC (Studies 1 and 2) or HEC (Study 3)

Adverse Reaction	Palonosetron Hydrochloride Injection 0.25 mg intravenously (N=633)	Ondansetron 32 mg intravenously (N=410)	Dolasetron 100 mg intravenously (N=194)
Headache	9%	8%	16%
Constipation	5%	2%	6%
Diarrhea	1%	2%	2%
Dizziness	1%	2%	2%
Fatigue	< 1%	1%	2%
Abdominal Pain	< 1%	< 1%	2%
Insomnia	< 1%	1%	2%

* Reported in at least 2% of patients in any treatment group

Less common adverse reactions, reported in 1% or less of patients, in Studies 1, 2 and 3 were:

- Cardiovascular: non-sustained tachycardia, bradycardia, hypotension, hypertension, myocardial ischemia, extrasystoles, sinus tachycardia, sinus arrhythmia, supraventricular extrasystoles and QT prolongation.
- Dermatological: allergic dermatitis, rash
- Hearing and Vision: motion sickness, tinnitus, eye irritation and amblyopia
- Gastrointestinal System: diarrhea, dyspepsia, abdominal pain, dry mouth, hiccups and flatulence
- General: weakness, fatigue, fever, hot flash, flu-like syndrome
- Liver: transient, asymptomatic increases in AST and/or ALT and bilirubin. These changes occurred predominantly in patients receiving highly emetogenic chemotherapy
- Metabolic: hyperkalemia, electrolyte fluctuations, hyperglycemia, metabolic acidosis, glycosuria, appetite decrease, anorexia
- Musculoskeletal: arthralgia
- Nervous System: dizziness, somnolence, insomnia, hypersomnia, paresthesia
- Psychiatric: anxiety, euphoric mood
- Urinary System: urinary retention
- Vascular: vein discoloration, vein distention

In other studies, 2 subjects experienced severe constipation following a single Palonosetron Hydrochloride Injection dose of approximately 0.75 mg (three times the recommended dose).

Pediatrics Aged 2 Months to 17 Years

In a pediatric clinical trial, 163 pediatric cancer patients with a mean age of 8 years received a single 20 mcg/kg (maximum 1.5 mg) intravenous infusion of Palonosetron Hydrochloride Injection 30 minutes before beginning the first cycle of emetogenic chemotherapy [see *Clinical Studies (14.2)*]. Adverse reactions were evaluated in pediatric patients receiving Palonosetron Hydrochloride Injection for up to 4 chemotherapy cycles. The following adverse reactions were reported in less than 1% of patients:

- Nervous System: headache, dizziness, dyskinesia.
- General: infusion site pain.
- Dermatological: allergic dermatitis, skin disorder.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of palonosetron HCl. Because these reactions are reported voluntarily from a population of uncertain size, it is not always

possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Hypersensitivity reactions: including dyspnea, bronchospasm, swelling/edema, erythema, pruritus, rash, urticaria, anaphylaxis and anaphylactic shock [see *Warnings and Precautions (5.1)*]
- Injection site reactions: including burning, induration, discomfort and pain

7. DRUG INTERACTIONS

7.1 Serotonergic Drugs

Serotonin syndrome (including altered mental status, autonomic instability, and neuromuscular symptoms) has been described following the concomitant use of 5-HT₃ receptor antagonists and other serotonergic drugs, including selective serotonin reuptake inhibitors (SSRIs) and serotonin and noradrenaline reuptake inhibitors (SNRIs). Monitor for the emergence of serotonin syndrome. If symptoms occur, discontinue Palonosetron Hydrochloride Injection and initiate supportive treatment [see *Warnings and Precautions (5.2)*].

8. USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no available data on palonosetron HCl use in pregnant women to inform a drug-associated risk.

In animal reproduction studies, no effects on embryo-fetal development were observed with the administration of oral palonosetron HCl during the period of organogenesis at doses up to 1,894 and 3,789 times the recommended human intravenous dose in rats and rabbits, respectively (*see Data*).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

In animal reproduction studies, no effects on embryo-fetal development were observed in pregnant rats given oral palonosetron HCl at doses up to 60 mg/kg/day (1,894 times the recommended human intravenous dose based on body surface area) or pregnant rabbits given oral doses up to 60 mg/kg/day (3,789 times the recommended human intravenous dose based on body surface area) during the period of organogenesis.

8.2 Lactation

Risk Summary

There are no data on the presence of palonosetron in human milk, the effects of palonosetron on the breastfed infant, or the effects of palonosetron on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Palonosetron Hydrochloride Injection and any potential adverse effect on the breastfed infant from palonosetron or from the underlying maternal condition.

8.4 Pediatric Use

Chemotherapy-Induced Nausea and Vomiting

Safety and effectiveness of Palonosetron Hydrochloride Injection have been established in pediatric

patients aged 1 month to less than 17 years for the prevention of acute nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy, including HEC. Use is supported by a clinical trial where 165 pediatric patients aged 2 months to less than 17 years were randomized to receive a single dose of Palonosetron Hydrochloride Injection 20 mcg/kg (maximum 1.5 mg) administered as an intravenous infusion 30 minutes prior to the start of emetogenic chemotherapy [see *Clinical Studies (14.2)*]. While this study demonstrated that pediatric patients require a higher palonosetron dose than adults to prevent chemotherapy-induced nausea and vomiting, the safety profile is consistent with the established profile in adults [see *Adverse Reactions (6.1)*].

Safety and effectiveness of Palonosetron Hydrochloride Injection in neonates (less than 1 month of age) have not been established.

8.5 Geriatric Use

Of the 1,374 adult cancer patients in clinical studies of intravenously administered palonosetron HCl, 316 (23%) were 65 years and over, while 71 (5%) were at least 75 years and over. Of the 1,520 adult patients in clinical studies of intravenously administered palonosetron HCl, 73 (5%) were at least 65 years old [see *Clinical Studies (14.1)*]. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, but greater sensitivity in some older individuals cannot be ruled out. Population pharmacokinetics analysis did not reveal any differences in palonosetron pharmacokinetics between cancer patients 65 years of age and older compared to younger patients [see *Clinical Pharmacology (12.3)*]. No dose adjustment is required for geriatric patients.


10. OVERDOSAGE

There is no known antidote to palonosetron. Overdose should be managed with supportive care.

Dialysis studies have not been performed, however, due to the large volume of distribution, dialysis is unlikely to be an effective treatment for palonosetron overdose. A single intravenous dose of palonosetron (HCl) at 30 mg/kg (947 and 474 times the human dose for rats and mice, respectively, based on body surface area) was lethal to rats and mice. The major signs of toxicity were convulsions, gasping, pallor, cyanosis and collapse.

11. DESCRIPTION

Palonosetron Hydrochloride Injection contains palonosetron as palonosetron HCl, an antiemetic and antinauseant agent. It is a serotonin-3 (5-HT₃) receptor antagonist with a strong binding affinity for this receptor. Chemically, palonosetron hydrochloride is: (3aS)-2-[(S)-1-Azabicyclo [2.2.2]oct-3-yl]-2,3,3a,4,5,6-hexahydro-1-oxo-1Hbenz[de]isoquinoline hydrochloride. Palonosetron hydrochloride exists as a single isomer and has the following structural formula:



C₁₉ H₂₄ N₂O•HCl

M.W. 332.87

Palonosetron Hydrochloride is a white to off-white crystalline powder. It is freely soluble in water, soluble in propylene glycol, and slightly soluble in ethanol and 2-propanol.

Palonosetron Hydrochloride Injection is a sterile, clear, colorless, non-pyrogenic, isotonic, buffered solution for intravenous administration. Palonosetron Hydrochloride Injection is available as a 5 mL

single-dose prefilled syringe.

Each 5 mL syringe contains: 0.25 mg palonosetron (equivalent to 0.28 mg palonosetron HCl), 202.4 mg mannitol, 2.5 mg edetate disodium dihydrate, 18 mg trisodium citrate dihydrate, and 7 mg citric acid anhydrous in water for intravenous administration.

The pH of the solution in the 5 mL syringe is 4.5 to 5.5.

12. CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Palonosetron is a 5-HT₃ receptor antagonist with a strong binding affinity for this receptor and little or no affinity for other receptors.

Cancer chemotherapy may be associated with a high incidence of nausea and vomiting, particularly when certain agents, such as cisplatin, are used. 5-HT₃ receptors are located on the nerve terminals of the vagus in the periphery and centrally in the chemoreceptor trigger zone of the area postrema. It is thought that chemotherapeutic agents produce nausea and vomiting by releasing serotonin from the enterochromaffin cells of the small intestine and that the released serotonin then activates 5-HT₃ receptors located on vagal afferents to initiate the vomiting reflex.

12.2 Pharmacodynamics

Cardiac Electrophysiology

The effect of intravenous palonosetron on blood pressure, heart rate, and ECG parameters including QTc were comparable to intravenous ondansetron and dolasetron in CINV clinical trials. In non-clinical studies palonosetron possesses the ability to block ion channels involved in ventricular de- and re-polarization and to prolong action potential duration.

At a dose of 9 times the maximum recommended adult dose, Palonosetron Hydrochloride Injection does not prolong the QT interval to any clinically relevant extent.

12.3 Pharmacokinetics

After intravenous dosing of palonosetron HCl in healthy subjects and cancer patients, an initial decline in palonosetron plasma concentrations is followed by a slow elimination from the body. Mean maximum plasma concentration (C_{max}) and area under the concentration-time curve (AUC_{0-∞}) are generally dose-proportional over the dose range of 0.3 to 90 mcg/kg in healthy subjects and in cancer patients.

Following a single intravenous dose of palonosetron HCl at 3 mcg/kg (or 0.21 mg/70 kg) to six cancer patients, mean (±SD) maximum plasma concentration was estimated to be 5,630 ± 5,480 ng/L and mean AUC was 35.8 ± 20.9 h•mcg/L.

Following intravenous administration of Palonosetron Hydrochloride Injection 0.25 mg once every other day for 3 doses in 11 cancer patients, the mean increase in plasma palonosetron concentration from Day 1 to Day 5 was 42±34%. Following intravenous administration of Palonosetron Hydrochloride Injection 0.25 mg once daily for 3 days in 12 healthy subjects, the mean (±SD) increase in plasma palonosetron concentration from Day 1 to Day 3 was 110±45%.

Distribution

Palonosetron has a volume of distribution of approximately 8.3 ± 2.5 L/kg. Approximately 62% of palonosetron is bound to plasma proteins.

Elimination

After a single intravenous dose of 10 mcg/kg [¹⁴C]-palonosetron, approximately 80% of the dose was recovered within 144 hours in the urine with palonosetron representing approximately 40% of the administered dose. In healthy subjects, the total body clearance of palonosetron was 0.160 ± 0.035

L/h/kg and renal clearance was 0.067 ± 0.018 L/h/kg. Mean terminal elimination half-life is approximately 40 hours.

Metabolism

Palonosetron is eliminated by multiple routes with approximately 50% metabolized to form two primary metabolites: N-oxide-palonosetron and 6-S-hydroxy-palonosetron. These metabolites each have less than 1% of the 5-HT₃ receptor antagonist activity of palonosetron. *In vitro* metabolism studies have suggested that CYP2D6 and to a lesser extent, CYP3A4 and CYP1A2 are involved in the metabolism of palonosetron. However, clinical pharmacokinetic parameters are not significantly different between poor and extensive metabolizers of CYP2D6 substrates.

Specific Populations

Pediatric Patients

Pharmacokinetic data was obtained from a subset of pediatric cancer patients that received 10 mcg/kg or 20 mcg/kg as a single intravenous dose of Palonosetron Hydrochloride Injection. When the dose was increased from 10 mcg/kg to 20 mcg/kg a dose-proportional increase in mean AUC was observed. Peak plasma concentrations (C_T) reported at the end of the 15-minute infusion of 20 mcg/kg were highly variable in all age groups and tended to be lower in patients less than 6 years than in older patients as shown in Table 4. The median half-life was 30 hours in overall age groups and ranged from about 20 to 30 hours across age groups after administration of 20 mcg/kg.

The total body clearance (L/h/kg) in patients 12 to 17 years old was similar to that in healthy adults. There are no apparent differences in volume of distribution when expressed as L/kg.

Table 4: Pharmacokinetics Parameters in Pediatric Cancer Patients following Intravenous Infusion of 20 mcg/kg Palonosetron Hydrochloride Injection Over 15 minutes

PK Parameter ^a	Pediatric Age Group			
	Less than 2 years	2 years to less than 6 years	6 years to less than 12 years	12 years to less than 17 years
	N=12	N=42	N=38	N=44
C _T ^b , ng/L	9025 (197)	9414 (252)	16275 (203)	11831 (176)
		N=5	N=7	N=10
AUC _{0-∞} , h·mcg/L		103.5 (40.4)	98.7 (47.7)	124.5 (19.1)
	N=6	N=14	N=13	N=19
Clearance ^c , L/h/kg	0.31 (34.7)	0.23 (51.3)	0.19 (46.8)	0.16 (27.8)
V _{ss} ^c , L/kg	6.08 (36.5)	5.29 (57.8)	6.26 (40.0)	6.20 (29.0)

^a Geometric Mean (CV) except for t_{1/2} which is median values

^b C_T is the plasma palonosetron concentration at the end of the 15-minute infusion

^c Clearance and V_{ss} calculated from 10 and 20 mcg/kg and are weight adjusted

Racial or Ethnic Groups

The pharmacokinetics of palonosetron were characterized in 24 healthy Japanese subjects over an intravenous dose range of 3 to 90 mcg/kg. Total body clearance was 25% higher in Japanese subjects compared to Whites, however, this increase is not considered to be clinically meaningful.

Patients with Renal Impairment

Mild to moderate renal impairment does not significantly affect palonosetron pharmacokinetic parameters. Total systemic exposure increased by approximately 28% in patients with severe renal impairment relative to healthy subjects. This increase is not considered clinically meaningful.

Patients with Hepatic Impairment

Hepatic impairment does not significantly affect total body clearance of palonosetron compared to the healthy subjects.

Drug Interaction Studies

In vitro studies indicated that palonosetron is not an inhibitor of CYP1A2, CYP2A6, CYP2B6, CYP2C9, CYP2D6, CYP2E1 and CYP3A4/5 (CYP2C19 was not investigated) nor does it induce the activity of CYP1A2, CYP2D6, or CYP3A4/5. Therefore, the potential for clinically significant drug interactions with palonosetron appears to be low.

Dexamethasone

Coadministration of 0.25 mg Palonosetron Hydrochloride Injection and 20 mg dexamethasone administered intravenously in healthy subjects revealed no pharmacokinetic drug-interactions between palonosetron and dexamethasone.

Oral Aprepitant

In an interaction study in healthy subjects where a single 0.25 mg intravenous dose of Palonosetron Hydrochloride Injection was administered on day 1 and oral aprepitant for 3 days (125 mg/80 mg/80 mg), the pharmacokinetics of palonosetron were not significantly altered (AUC: no change, C_{max} : 15% increase).

Metoclopramide

A study in healthy subjects involving a single 0.75 mg intravenous dose of Palonosetron Hydrochloride Injection and steady state oral metoclopramide (10 mg four times daily) demonstrated no significant pharmacokinetic interaction.

Corticosteroids, Analgesics, Antiemetics/Antinauseants, Antispasmodics and Anticholinergic Agents

In controlled clinical trials, Palonosetron Hydrochloride Injection has been safely administered with corticosteroids, analgesics, antiemetics/antinauseants, antispasmodics and anticholinergic agents.

13. NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 104-week carcinogenicity study in CD-1 mice, animals were treated with oral doses of palonosetron HCl at 10, 30 and 60 mg/kg/day. Treatment with palonosetron was not tumorigenic. The highest tested dose produced a systemic exposure to palonosetron (Plasma AUC) of about 150 to 289 times the human exposure (AUC= 29.8 h•mcg/L) at the recommended intravenous dose of 0.25 mg. In a 104-week carcinogenicity study in Sprague-Dawley rats, male and female rats were treated with oral doses of 15, 30 and 60 mg/kg/day and 15, 45 and 90 mg/kg/day, respectively. The highest doses produced a systemic exposure to palonosetron (Plasma AUC) of 137 and 308 times the human exposure at the recommended dose. Treatment with palonosetron produced increased incidences of adrenal benign pheochromocytoma and combined benign and malignant pheochromocytoma, increased incidences of pancreatic Islet cell adenoma and combined adenoma and carcinoma and pituitary adenoma in male rats. In female rats, it produced hepatocellular adenoma and carcinoma and increased the incidences of thyroid C-cell adenoma and combined adenoma and carcinoma.

Palonosetron was not genotoxic in the Ames test, the Chinese hamster ovarian cell (CHO/HGPRT) forward mutation test, the *ex vivo* hepatocyte unscheduled DNA synthesis (UDS) test or the mouse micronucleus test. It was, however, positive for clastogenic effects in the Chinese hamster ovarian (CHO) cell chromosomal aberration test.

Palonosetron HCl at oral doses up to 60 mg/kg/day (about 1,894 times the recommended human intravenous dose based on body surface area) was found to have no effect on fertility and reproductive

performance of male and female rats.

14. CLINICAL STUDIES

14.1 Prevention of Nausea and Vomiting Associated with MEC and HEC in Adults

Efficacy of a single intravenous dose of Palonosetron Hydrochloride Injection in preventing acute and delayed nausea and vomiting associated with MEC or HEC were studied in 4 trials. In these double-blind studies, complete response rates (no emetic episodes and no rescue medication) and other efficacy parameters were assessed through at least 120 hours after administration of chemotherapy.

The safety and efficacy of Palonosetron Hydrochloride Injection in repeated courses of chemotherapy was also assessed.

Moderately Emetogenic Chemotherapy

Two double-blind trials (Study 1 and Study 2) involving 1,132 patients compared a single dose of Palonosetron Hydrochloride Injection with either a single-dose of ondansetron (Study 1) or dolasetron (Study 2) given 30 minutes prior to MEC, including carboplatin, cisplatin $\leq 50 \text{ mg/m}^2$, cyclophosphamide $< 1,500 \text{ mg/m}^2$, doxorubicin $> 25 \text{ mg/m}^2$, epirubicin, irinotecan, and methotrexate $> 250 \text{ mg/m}^2$. Concomitant corticosteroids were not administered prophylactically in Study 1 and were only used by 4 to 6% of patients in Study 2. The majority of patients in these studies were women (77%), White (65%) and naïve to previous chemotherapy (54%). The mean age was 55 years.

Highly Emetogenic Chemotherapy

A double-blind, dose-ranging trial evaluated the efficacy of a single intravenous dose of Palonosetron Hydrochloride Injection from 0.3 to 90 mcg/kg (equivalent to $< 0.1 \text{ mg}$ to 6 mg fixed dose) in 161 chemotherapy-naïve adult cancer patients receiving HEC, either cisplatin $\geq 70 \text{ mg/m}^2$ or cyclophosphamide $> 1,100 \text{ mg/m}^2$. Concomitant corticosteroids were not administered prophylactically. Analysis of data from this trial indicates that 0.25 mg is the lowest effective dose in preventing acute nausea and vomiting associated with HEC.

A double-blind trial involving 667 patients compared a single intravenous dose of Palonosetron Hydrochloride Injection with a single intravenous dose of ondansetron (Study 3) given 30 minutes prior to HEC, including cisplatin $\geq 60 \text{ mg/m}^2$, cyclophosphamide $> 1,500 \text{ mg/m}^2$, and dacarbazine. Corticosteroids were co-administered prophylactically before chemotherapy in 67% of patients. Of the 667 patients, 51% were women, 60% White, and 59% naïve to previous chemotherapy. The mean age was 52 years.

Efficacy Results

Studies 1, 2 and 3 show that Palonosetron Hydrochloride Injection was effective in the prevention of nausea and vomiting associated with initial and repeat courses of MEC and HEC in the acute phase (0 to 24 hours) [Table 5]. Clinical superiority over other 5-HT₃ receptor antagonists has not been adequately demonstrated in the acute phase. In Study 3, efficacy was greater when prophylactic corticosteroids were administered concomitantly.

Studies 1 and 2 show that Palonosetron Hydrochloride Injection was effective in the prevention of nausea and vomiting associated with initial and repeat course of MEC in the delayed phase (24 to 120 hours) [Table 6] and overall phase (0 to 120 hours) [Table 7].

Table 5: Prevention of Acute Nausea and Vomiting (0 to 24 Hours) in Adults with Nausea and Vomiting Associated with MEC or HEC in Studies 1, 2 and 3: Complete Response Rates

Chemotherapy	Study	Treatment Group	N ^a	% with Complete Response	P-value ^b
--------------	-------	-----------------	----------------	--------------------------	----------------------

Moderately Emetogenic	1	Palonosetron Hydrochloride Injection 0.25 mg intravenously	189	81	0.009	97.5% Confidence Interval Palonosetron Hydrochloride Injection minus Comparator^c
		Ondansetron 32 mg intravenously	185	69		
	2	Palonosetron Hydrochloride Injection 0.25 mg intravenously	189	63	NS	
		Dolasetron 100 mg intravenously	191	53		
Highly Emetogenic	3	Palonosetron Hydrochloride Injection 0.25 mg intravenously	223	59	NS	
		Ondansetron 32 mg intravenously	221	57		

^a Intent-to-treat cohort

^b 2-sided Fisher's exact test. Significance level at $\alpha=0.025$.

^c These studies were designed to show non-inferiority. A lower bound greater than -15% demonstrates non-inferiority between Palonosetron Hydrochloride Injection and comparator.

Table 6: Prevention of Delayed Nausea and Vomiting (24 to 120 Hours) Associated with MEC in Adults in Studies 1 and 2: Complete Response Rates

Chemotherapy	Study	Treatment Group	N ^a	% with Complete Response	P-value ^b	97.5% Confidence Interval Palonosetron Hydrochloride Injection minus Comparator ^c
Moderately Emetogenic	1	Palonosetron Hydrochloride Injection 0.25 mg intravenously	189	74	<0.001	
		Ondansetron 32 mg intravenously ^d	185	55		
	2	Palonosetron Hydrochloride Injection 0.25 mg intravenously	189	54	0.004	
		Dolasetron 100 mg intravenously	191	39		

^a Intent-to-treat cohort

^b 2-sided Fisher's exact test. Significance level at $\alpha=0.025$.

^c These studies were designed to show non-inferiority. A lower bound greater than -15% demonstrates non-inferiority between Palonosetron Hydrochloride Injection and comparator.

^d Ondansetron 32 mg intravenous was used in the clinical trial. Although this dose was used in the trial, it is no longer the currently recommended dose. Refer to the ondansetron prescribing information for the current recommended dose.

Table 7: Prevention of Overall Nausea and Vomiting (0 to 120 Hours) Associated with MEC in Adults in Studies 1 and 2: Complete Response Rates

Chemotherapy	Study	Treatment Group	N ^a	% with Complete Response	P-value ^b	97.5% Confidence Interval Palonosetron Hydrochloride Injection minus Comparator ^c	
Moderately Emetogenic	1	Palonosetron Hydrochloride Injection 0.25 mg intravenously	189	69	<0.001		
		Ondansetron 32 mg intravenously ^d	185	50			
	2	Palonosetron Hydrochloride Injection 0.25 mg intravenously	189	46	0.021		
		Dolasetron 100 mg intravenously	191	34			

^a Intent-to-treat cohort

^b 2-sided Fisher's exact test. Significance level at $\alpha=0.025$.

^c These studies were designed to show non-inferiority. A lower bound greater than -15% demonstrates non-inferiority between Palonosetron Hydrochloride Injection and comparator.

^d Ondansetron 32 mg intravenously was used in the clinical trial. Although this dose was used in the trial, it is no longer the currently recommended dose. Refer to the ondansetron prescribing information for the current recommended dose.

14.2 Prevention of Nausea and Vomiting Associated with Emetogenic Chemotherapy, Including HEC in Pediatric Patients

One double-blind, active-controlled clinical trial was conducted in pediatric cancer patients. The total population (N = 327) had a mean age of 8.3 years (range 2 months to 16.9 years) and were 53% male; and 96% white. Patients were randomized and received a 20 mcg/kg (maximum 1.5 mg) intravenous infusion of Palonosetron Hydrochloride Injection 30 minutes prior to the start of emetogenic chemotherapy (followed by placebo infusions 4 and 8 hours after the dose of Palonosetron Hydrochloride injection) or 0.15 mg/kg of intravenous ondansetron 30 minutes prior to the start of emetogenic chemotherapy (followed by ondansetron 0.15 mg/kg infusions 4 and 8 hours after the first dose of ondansetron, with a maximum total dose of 32 mg). Emetogenic chemotherapies administered included doxorubicin, cyclophosphamide (< 1,500 mg/m²), ifosfamide, cisplatin, dactinomycin, carboplatin, and daunorubicin. Adjuvant corticosteroids, including dexamethasone, were administered with chemotherapy in 55% of patients.

Complete Response in the acute phase of the first cycle of chemotherapy was defined as no vomiting, no retching, and no rescue medication in the first 24 hours after starting chemotherapy. Efficacy was based on demonstrating non-inferiority of intravenous Palonosetron Hydrochloride injection compared to intravenous ondansetron. Non-inferiority criteria were met if the lower bound of the 97.5% confidence interval for the difference in Complete Response rates of intravenous Palonosetron Hydrochloride injection minus intravenous ondansetron was larger than -15%. The non-inferiority margin was 15%.

Efficacy Results

As shown in Table 8, intravenous Palonosetron Hydrochloride Injection 20 mcg/kg (maximum 1.5 mg) demonstrated non-inferiority to the active comparator during the 0 to 24-hour time interval.

Table 8: Prevention of Acute Nausea and Vomiting (0 to 24 Hours) Associated with Emetogenic Chemotherapy in Pediatric Patients: Complete Response Rates

Palonosetron	Ondansetron	Difference [97.5% Confidence Interval] ^a :
--------------	-------------	---

Hydrochloride Injection 20 mcg/kg intravenously (N=165)	0.15 mg/kg for 3 intravenous doses (N=162)	Palonosetron Hydrochloride Injection minus intravenous Ondansetron Comparator
59.4%	58.6%	0.36% [-11.7%, 12.4%]

^a To adjust for multiplicity of treatment groups, a lower-bound of a 97.5% confidence interval was used to compare to -15%, the negative value of the non-inferiority margin.

In patients that received Palonosetron Hydrochloride Injection at a lower dose than the recommended dose of 20 mcg/kg, non-inferiority criteria were not met.

16. HOW SUPPLIED/STORAGE AND HANDLING

Palonosetron Hydrochloride Injection 0.25 mg/5 mL (free base) single-dose prefilled syringe is available as follows:

Product No.	NDC No.	Strength	Package
673189	63323-673-89	0.25 mg per 5 mL (0.05 mg per mL)	5 mL single-dose prefilled syringe, individually packaged in cartons of ten.

Storage

- Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].
- Protect from freezing.
- Protect from light.
- Discard unused portion

17. PATIENT COUNSELING INFORMATION

Advise the patient or caregiver to read the FDA-approved patient labeling (Patient Information).

Hypersensitivity Reactions

Advise patients that hypersensitivity reactions, including anaphylaxis and anaphylactic shock, have been reported in patients with or without known hypersensitivity to other 5-HT₃ receptor antagonists. Advise patients to seek immediate medical attention if any signs or symptoms of a hypersensitivity reaction occur with administration of Palonosetron Hydrochloride Injection [see *Warnings and Precautions (5.1)*].

Serotonin Syndrome

Advise patients of the possibility of serotonin syndrome, especially with concomitant use of Palonosetron Hydrochloride injection and another serotonergic agent such as medications to treat depression and migraines. Advise patients to seek immediate medical attention if the following symptoms occur: changes in mental status, autonomic instability, neuromuscular symptoms with or without gastrointestinal symptoms [see *Warnings and Precautions (5.2)*].



Lake Zurich, IL 60047

Made in Austria

www.fresenius-kabi.com/us

451400B

(PAL-oh-NOE-se-tron HYE-dro-KLOR-ide) Injection
for Intravenous Use

Read this Patient Information before you receive Palonosetron Hydrochloride Injection and each time you receive Palonosetron Hydrochloride Injection. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment.

What is Palonosetron Hydrochloride Injection?

Palonosetron Hydrochloride Injection is a prescription medicine called an “antiemetic.” Palonosetron Hydrochloride Injection is used in adults to help prevent the nausea and vomiting that happens:

- right away or later with certain anti-cancer medicines (chemotherapy)

Palonosetron Hydrochloride Injection is used in children 1 month old to less than 17 years of age to help prevent the nausea and vomiting that happens right away with certain anti-cancer medicines (chemotherapy).

- It is not known if Palonosetron Hydrochloride Injection is safe and effective in children less than 1 month old to help prevent nausea and vomiting after chemotherapy.

Who should not receive Palonosetron Hydrochloride Injection?

Do not receive Palonosetron Hydrochloride Injection if you are allergic to palonosetron hydrochloride or any of the ingredients in Palonosetron Hydrochloride Injection. See the end of this leaflet for a complete list of ingredients in Palonosetron Hydrochloride Injection.

What should I tell my doctor before receiving Palonosetron Hydrochloride Injection?

Before receiving Palonosetron Hydrochloride Injection, tell your doctor about all of your medical conditions, including if you:

- have had an allergic reaction to another medicine for nausea or vomiting
- are pregnant or plan to become pregnant. It is not known if Palonosetron Hydrochloride Injection will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if Palonosetron Hydrochloride Injection passes into your breast milk or if it will affect your baby or your breast milk. Talk to your doctor about the best way to feed your baby if you will receive Palonosetron Hydrochloride Injection.

Tell your doctor about all of the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

Palonosetron Hydrochloride Injection and certain other medicines can affect each other, causing serious side effects.

How will I receive Palonosetron Hydrochloride injection?

- Palonosetron Hydrochloride Injection will be given to you in your vein by intravenous (I.V.) injection.
- Palonosetron Hydrochloride Injection is usually given about 30 minutes before you receive your anti-cancer medicine (chemotherapy).

What are the possible side effects of Palonosetron Hydrochloride Injection? Palonosetron Hydrochloride Injection may cause serious side effects, including:

- **Serious allergic reactions**, such as anaphylaxis. Get emergency medical help right away if you get any of the following symptoms.
 - hives
 - swollen face
 - breathing trouble
 - chest pain
- **Serotonin Syndrome.** A possible life threatening problem called serotonin syndrome can happen with medicines called 5-HT₃ receptor antagonists, including Palonosetron Hydrochloride Injection, especially when used with medicines used to treat depression and migraine headaches called serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), monoamine oxidase inhibitors (MAOIs) and certain other medicines. Tell your doctor or nurse right

away if you have any of the following symptoms of serotonin syndrome:

- agitation, seeing things that are not there (hallucinations), confusion, or coma
- dizziness, sweating, flushing, or fever
- seizures
- fast heartbeat or unusual and frequent changes in your blood pressure
- tremors, stiff muscles, muscle twitching, overactive reflexes, or loss of coordination
- nausea, vomiting, or diarrhea

The most common side effects in adults who receive Palonosetron Hydrochloride Injection to help prevent nausea and vomiting that happens with **certain anti-cancer medicine (chemotherapy)** include: headache and constipation.

These are not all the possible side effects from Palonosetron Hydrochloride Injection.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of Palonosetron Hydrochloride Injection Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. You can ask your doctor or pharmacist for information about Palonosetron Hydrochloride Injection that is written for health professionals.

What are the ingredients in Palonosetron Hydrochloride Injection?

Active ingredient: palonosetron hydrochloride

Inactive ingredients: mannitol, disodium edetate, and citrate buffer in water



Lake Zurich, IL 60047

Made in Austria

For more information, go to www.fresenius-kabi.com/us or call **1-800-551-7176**.

This Patient Information has been approved by the U.S. Food and Drug Administration.

451401B

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Palonosetron Hydrochloride 0.25 mg Single-Dose Prefilled Syringe Label

NDC 63323-673-21

Palonosetron Hydrochloride Injection

0.25 mg* per 5 mL

(0.05 mg* per mL)


For Intravenous Injection Only.

Discard Unused Portion.

5 mL

Single-Dose Syringe

Rx only



PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Palonosetron Hydrochloride 0.25 mg Single-Dose Prefilled Syringe Individual Carton Panel

NDC 63323-673-21

Palonosetron Hydrochloride Injection


0.25 mg* per 5 mL

(0.05 mg* per mL)

For Intravenous Injection Only.

Discard Unused Portion. Rx only

1x 5 mL Single-Dose Syringe



PACKAGE LABEL - PRINCIPAL DISPLAY PANEL - Palonosetron Hydrochloride 0.25 mg Single-Dose Prefilled Syringe Carton Panel

NDC 63323-673-89

Palonosetron Hydrochloride Injection

0.25 mg* per 5 mL

(0.05 mg* per mL)

For Intravenous Injection Only.

10 x 5 mL
Single-Dose Syringes

Rx only

NDC 63323-673-89

Palonosetron Hydrochloride
Injection

0.25 mg* per 5 mL

(0.05 mg* per mL)

For Intravenous Injection Only.

10 x 5 mL
Single-Dose Syringes
Rx only



PALONOSETRON

palonosetron injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63323-673
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PALONOSETRON HYDROCHLORIDE (UNII: 23310D4I19) (PALONOSETRON - UNII:5D06587D6R)	PALONOSETRON	0.25 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	202.4 mg in 5 mL
EDETATE DISODIUM (UNII: 7FLD91C86K)	2.5 mg in 5 mL
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	18 mg in 5 mL
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	7 mg in 5 mL

Packaging

Marketing Start Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63323-673-89	10 in 1 CARTON	09/19/2018	
1	NDC:63323-673-21	5 mL in 1 SYRINGE, PLASTIC; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206801	09/19/2018	

Labeler - Fresenius Kabi USA, LLC (608775388)

Establishment

Name	Address	ID/FEI	Business Operations
Fresenius Kabi Austria		300206604	ANALYSIS(63323-673) , MANUFACTURE(63323-673) , PACK(63323-673)

Revised: 1/2021

Fresenius Kabi USA, LLC