

**FLU HBP MAXIMUM STRENGTH- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated**  
**Cardinal Health 110, LLC. DBA Leader**

-----  
**Leader 44-738**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg  
Chlorpheniramine maleate 2 mg  
Dextromethorphan HBr 10 mg

***Purpose***

Pain reliever/fever reducer  
Antihistamine  
Cough suppressant

***Uses***

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - sore throat
  - sneezing
  - headache
  - runny nose
  - cough
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
- children under 12 years: ask a doctor

### ***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from excessive moisture
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, FD&C red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

### ***Questions or comments?***

**1-800-200-6313**

### ***Principal Display Panel***

**LEADER™**

NDC 70000-0050-1

Maximum Strength

Multi-Symptom

**Flu HBP**

#### **Acetaminophen**

Chlorpheniramine Maleate | Dextromethorphan HBr

Pain Reliever / Fever Reducer

Antihistamine | Cough Suppressant

#### **Relieves:**

Body Aches, Pains and Headache,

Fever, Cough, Runny Nose,

Sore Throat, Sneezing

For People with

High Blood Pressure

Decongestant-Free

**COMPARE TO**

**CORICIDIN® HBP**

**MULTI-SYMPATOM FLU**

## **MAXIMUM STRENGTH**

active ingredients\*

100% Money Back Guarantee

20 CAPLETS

Actual Size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS OF TAMPERING**

50844

REV1019A73809

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Multi-Symptom Flu Maximum Strength.

©2022 Cardinal Health. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

DISTRIBUTED BY CARDINAL HEALTH

DUBLIN, OHIO 43017

[www.myleader.com](http://www.myleader.com)

1-800-200-6313

**100%**

**Money Back**

**Guarantee**

Return to place of purchase.

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Multi-Symptom Flu Maximum Strength.  
©2022 Cardinal Health, CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

DISTRIBUTED BY CARDINAL HEALTH  
DUBLIN, OHIO 43017  
www.myleader.com  
1-800-200-6313

**100% Money Back Guarantee**  
Return to place of purchase.

**PARENTS:**  
Learn about teen medicine abuse  
www.StopMedicineAbuse.org

**LEADER™**

NDC 70000-0050-1

**Maximum Strength Multi-Symptom**

**Flu HBP**

**Acetaminophen**  
**Chlorpheniramine Maleate | Dextromethorphan HBr**  
**Pain Reliever / Fever Reducer**  
**Antihistamine | Cough Suppressant**

**Relieves:**  
Body Aches, Pains and Headache,  
Fever, Cough, Runny Nose,  
Sore Throat, Sneezing

For People with  
High Blood Pressure  
Decongestant-Free

**20 CAPLETS**

Actual Size

**COMPARE TO CORICIDIN® HBP MULTI-SYMPOM FLU MAXIMUM STRENGTH active ingredients\***

100% Money Back Guarantee

**Drug Facts (continued)**

F&C red #40 aluminum lake, lactin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium stearoyl glycolate, stearic acid, tac, titanium dioxide

**Questions or comments? 1-800-200-6313**

**Drug Facts (continued)**

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from excessive moisture ■ see end flap for expiration date and lot number

**Inactive ingredients** com starch, croscopolone,

**Drug Facts (continued)**

**Uses**  
■ temporarily relieves these common cold and flu symptoms:  
■ minor aches and pains ■ sore throat ■ cough  
■ sneezing ■ headache ■ runny nose  
■ temporarily reduces fever

**Drug Facts (continued)**

**Active ingredients (in each caplet)** Purpose  
Acetaminophen 325 mg ..... Pain reliever/fever reducer  
Chlorpheniramine maleate 2 mg ..... Antihistamine  
Dextromethorphan HBr 10 mg ..... Cough suppressant

**Drug Facts (continued)**

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ blisters ■ rash ■ skin reddening  
If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
■ difficulty in urination due to enlargement of the prostate gland  
■ liver disease ■ glaucoma  
■ cough that occurs with too much phlegm (mucus)  
■ a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use if you are**  
■ taking sedatives or tranquilizers  
■ taking the blood thinning drug warfarin

**When using this product**  
■ avoid alcoholic beverages  
■ marked drowsiness may occur  
■ alcohol, sedatives, and tranquilizers may increase drowsiness

■ excitability may occur, especially in children  
■ use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**  
■ pain or cough gets worse or lasts more than 7 days ■ new symptoms occur  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not take more than directed  
■ adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.  
■ children under 12 years: ask a doctor

**Other information**  
■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**

CIN 5621453 REV. 3/22



**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

50844  
REV1019A73809

No print/No varnish Lot & Exp date

8-0225-738-09  
REV1019A73809

**Leader 44-738**

**FLU HBP MAXIMUM STRENGTH**

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70000-0050
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

<b>CROSPROVIDONE</b> (UNII: 2S7830E561)
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)

Product Characteristics			
<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	44;738
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0050-1	2 in 1 CARTON	06/17/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/17/2020	

**Labeler** - Cardinal Health 110, LLC. DBA Leader (063997360)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(70000-0050) , pack(70000-0050)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(70000-0050)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(70000-0050)

Revised: 3/2024

Cardinal Health 110, LLC. DBA Leader