LORATADINE AND PSEUDOEPHEDRINE- loratadine and pseudoephedrine sulfate tablet, extended release Rite Aid

allergy relief & nasal decongestant

Drug Facts

Active ingredients (in each tablet)	Purpose
Loratadine, USP 10 mg	Antihistamine
Pseudoephedrine sulfate, USP	
240 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - sneezing
 - itchy, watery eyes
 - runny nose
 - itching of the nose or throat
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

• do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet daily with a full glass of water; not more than 1 tablet in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- sodium: contains 10 mg/tablet
- calcium: contains 25 mg/tablet
- TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.
- store between 20° C to 25° C (68° F to 77° F).
- protect from light and store in a dry place

Inactive ingredients

calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium alginate, sodium citrate, talc and titanium dioxide

Questions?

call **1-800-406-7984**

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011

PRINCIPAL DISPLAY PANEL - 15 Tablet Blister Pack Carton

24 HOUR

Compare to the active ingredients of Claritin-D[®] 24 Hour**

NDC 11822-9990-5

ORIGINAL PRESCRIPTION STRENGTH
ALLERGY RELIEF &
NASAL DECONGESTANT
LORATADINE, USP 10 mg / PSEUDOEPHEDRINE SULFATE, USP 240 mg

ANTIHISTAMINE & NASAL DECONGESTANT

NON-DROWSY*
INDOOR / OUTDOOR ALLERGIES
24 HOUR RELIEF OF

Nasal & sinus congestion due to colds or allergies

Sneezing • Runny nose • Itchy, watery eyes Itchy throat or nose due to allergies

*WHEN TAKEN AS DIRECTED. SEE DRUG FACTS PANEL.

ACTUAL SIZE

15 EXTENDED-RELEASE TABLETS

Non Varnish Area

ORIGINAL PRESCRIPTION STRENGTH

ALLERGY RELIEF & NASAL DECONGESTANT

LORATADINE, USP 10 mg / PSEUDOEPHEDRINE SULFATE, USP 240 mg

ANTIHISTAMINE & NASAL DECONGESTANT

24 HOUR

Compare to the active ingredients of Claritin-D° 24 Hour**

NDC 11822-9990-5

ORIGINAL PRESCRIPTION STRENGTH

ALLERGY RELIEF &

LORATADINE, USP 10 mg / PSEUDOEPHEDRINE SULFATE, USP 240 mg

ANTIHISTAMINE & NASAL DECONGESTANT

24 HOUR RELIEF OF

Nasal & sinus congestion due to colds or allergies

Sneezing • Runny nose • Itchy, watery eyes Itchy throat or nose due to allergies

EXTENDED-RELEASE TABLETS Keep the carton, It contains important information, See end panel for expiration date.



**AI trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Claritin-D*.

*WHEN TAKEN AS DIRECTED, SEE DRUG FACTS PANEL.

Questions? call 1-800-406-7984

Inactive ingredients calcum carbonate, coabidal silicon dioxide, hydroxypropyl calludose, hyprometace, iron oxide black, factose monohydrate, magnesium stearate, microcrystaline cellulose, polyethytene glycol, povidone, pregelatinized starch, propylene glycol, sheller glace, sodium alginate, sodium cirrate, talc and titanium dioxide pregelatinized starch, propylene glycol, sheller glace, sodium alginate, sodium cirrate, talc and titanium dioxide

- protect from light and store in a dry place
- 12MPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.

 - Other information

 sodium: contains 10 mg/tablet
 calcium: contains 25 mg/tablet

Drug Facts (continued)

or kidney disease ask a doctor consumers with Inver 12 years of age SEK S COCIOL children under SINON PS IN 191061 I I tablet daily with a full glass of water; not more than 12 years and over adults and children ■ do not divide, crush, chew or dissolve the tablet Directions

Keep out of reach of children. In case of overdose, get medical, help or contact a Poison Control Center right away (1-800-222-1222).

I pregnant or breast-feeding, ask a health professional ■ nervousness, dizziness or sleeplessness occurs accompanied by a fever

ain allergic reaction to this product occurs. Seek
medical help right away.
 symptoms do not improve within 7 days or are
symptoms do not improve within 7 days or are

Stop use and ask a doctor if

When using this product do not take more than directed. Taking more than directed may cause

 Industring due to an enlarged prostate gland
 Inver or fiddney disease. Your doctor should
 determine if you need a different dose. thyroid disease
 diabetes
 diabetes eunsseud poord uffilu ■ heart disease Ask a doctor before use if you have

Drug Facts (continued)

In our meeting the section to the product or early of the ingredients in our modelling of the ingredients in the product or early of the ingredients of the individual (NAM) (certain drugs for depression, psychiatric, or emotional conditions, between the individual product stopping the MACI drug. If you do not know if your prescription drug contains an MACI, ask a doctor or pharmacist before taking this product.

- Warnings

 Do not use

 Do not use

 If you have ever had an allergic reaction to this

- temporarily restores freer breathing through the common cold, hay fever or other upper respiratory
- fever or other upper respiratory allergies:

 E sneezing it ichty, watery eyes


 Fromosorally releves sinus congestion and pressure

 Remoorarly releves raisal congestion due to the

 Remoorarly releves raisal congestion due to the
 - temporarity refeves these symptoms due to hay รอรก

USP 240 mg. ...gm 01 9SU ,anibatano. ,atalius anihadiqaobuas^c (in each tablet) Purpose Active ingredients

Drug Facts



PRINCIPAL DISPLAY PANEL - 10 Tablet Blister Pack Carton

NDC 11822-9990-0 24 HOUR

Compare to the active ingredients in Claritin-D[®] 24 Hour**

ORIGINAL PRESCRIPTION STRENGTH

ALLERGY RELIEF & NASAL DECONGESTANT

LORATADINE, USP 10 mg / PSEUDOEPHEDRINE SULFATE, USP 240 mg

ANTIHISTAMINE & NASAL DECONGESTANT

NON-DROWSY*
INDOOR / OUTDOOR ALLERGIES
24 HOUR RELIEF OF
Nasal & sinus congestion due to colds or allergies

Sneezing • Runny nose • Itchy, watery eyes Itchy throat or nose due to allergies

*WHEN TAKEN AS DIRECTED. SEE DRUG FACTS PANEL.

ACTUAL SIZE

10 EXTENDED-RELEASE TABLETS

directed. Taking more than directed may cause When using this product do not take more than determine it you need a different dose. ■ liver or kidney disease. Your doctor should ■ trouble urinating due to an enlarged prostate gland ■ heart disease
■ thyroid disease
■ high blood pressure
■ disbetes
■ tsouch Ask a doctor before use if you have before taking this product. the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist Parkinson's disease), or for 2 weeks after stopping depression, psychiatric, or emotional conditions, or

product or any of its ingredients

If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

■ if you have ever had an allergic reaction to this Do not use

Warnings

■ temporarily restores freer breathing through the allergies

common cold, hay fever or other upper respiratory ■ temporarily relieves nasal congestion due to the ■ temporarily relieves sinus congestion and pressure

■ reduces swe ling of nasal passages

חגות וויים בינג (couriuned)

■ protect from light and store in a dry place ■ store between 20° C to 25° C (68° F to 77° F) TAMPERING. ARE TORN, BROKEN OR SHOW ANY SIGNS OF ■ TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS

■ sodium: contains 10 mg/tablet ■ calcium: contains 25 mg/tablet

Other information

. ,	
consumers with liver or kidney disease	ask a doctor
children under 12 years of age	ask a doctor
Directions ■ do not divide, crush adults and children 2 years and over	, chew or dissolve the tablet tables a tablet daily with a full glass of water; not more than 1 tablet in 24 hours

(1-800-222-1222) medical help or contact a Poison Control Center right away Keep out of reach of children. In case of overdose, get

It pregnant or breast-feeding, ask a health professional ■ nervousness, dizziness or sleeplessness occurs

■ symptoms do not improve within 7 days or are medical help right away an allergic reaction to this product occurs. Seek Stop use and ask a doctor if

Drug Facts (continued)

вссошраліва ру з течег

Iching of the nose or throat ■ runny nose ■ Itcny, watery eyes 6uizəəus ■ USes Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

Loratadine, USP 10 mg. Antihistamine. Purpose Active ingredients (in each tablet)

orug Facts

R1021

Compare to the active ingredients in Claritin-D® 24 Hour**

NDC 11822-9990-0 **24 HOUR**

ORIGINAL PRESCRIPTION STRENGTH

LORATADINE, USP 10 mg / PSEUDOEPHEDRINE SULFATE, USP 240 mg

ANTIHISTAMINE & NASAL DECONGESTANT

NON-DROWSY* INDOOR / OUTDOOR ALLERGIES

24 HOUR RELIEF OF

Nasal & sinus congestion due to colds or allergies

Sneezing • Runny nose • Itchy, watery eyes Itchy throat or nose due to allergies

ACTUAL SIZE

EXTENDED-

*WHEN TAKEN AS D**i**rected. See drug facts panel.

RELEASE TABLETS

Keep the carton. It contains important information. See end panel for expiration date,

Questions? call 1-800-406-7984

oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glazse, sodium alginate, sodium citrate, talc and titanium dioxide Inactive ingredients calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron

DISTRIBUTED BY

SATISFACTION GUARANTEE: ppily refund your money /ou're not satisfied, we'll



LEM 331426



LORATADINE AND PSEUDOEPHEDRINE

loratadine and pseudoephedrine sulfate tablet, extended release

Product Information				
Product Type HUMAN	OTC DRUG Item	em Code (Source)	NDC:11822-9990	

Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	10 mg		
PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE - UNII: 7CUC9DDI9F)	- PSEUDOEPHEDRINE SULFATE	240 mg		

Inactive Ingredients	
Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: 08232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B710)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERROSOFERRIC OXIDE (UNII: XM0M87F357)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics				
Color	white	Score	no score	
Shape	OVAL	Size	17mm	
Flavor		Imprint Code	RX724	
Contains				

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822- 9990-5	1 in 1 CARTON	11/17/2004	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822- 9990-0	1 in 1 CARTON	11/17/2004	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076557	11/17/2004	

Labeler - Rite Aid (014578892)

Establishment				
Name	Address	ID/FEI	Business Operations	
Ohm Laboratories Inc.		184769029	manufacture(11822-9990)	

Revised: 5/2022 Rite Aid