# MILK OF MAGNESIA- magnesium hydroxide suspension Meijer, Inc.

-----

meijer mom CHRY

# Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

# **Purpose**

Saline laxative

#### Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

# Warnings

# Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug.

This product may interact with certain prescription drugs.

# Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose

age	dose
adults and children 12 years and over	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

# Other information

- each 15 mL contains: magnesium 500 mg, sodium 6 mg
- do not freeze
- store at room temperature tightly closed

# **Inactive ingredients**

carboxymethylcellulose sodium, citric acid, D & C red no. 28, flavor, glycerin, microcrystalline cellulose, purified water, sodium citrate, sodium hypochlorite, sucrose, xanthan gum

**Questions or comments?** 

1-800-540-3765

package Label



#### Drug Facts

TAMPER EVIDENT: Do not use if imprinted neckband is missing or broken.

Active ingredient (in each 15 mL)
Magnesium hydroxide 1200 mg ......

Purpose

• relieves occasional constipation (irregularity)
• generally produces bowel movement in ½ to 6 hours

#### Warnings

Ask a doctor before use if you have • kidney disease

a magnesium-restricted diet

· stomach pain, nausea, or vomiting

. a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with prescription drugs.

#### Stop use and ask a doctor if

you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
 you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

· shake well before use

- do not exceed the maximum recommended daily dose in a 24 hour period
   dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
   drink a full glass (8 oz) of liquid with each dose

armit a rail grace (e ex) or india mar exert acce		
age	dose	
adults and children 12 years and over	30 mL to 60 mL	
children 6 to 11 years	15 mL to 30 mL	
children under 6 years	ask a doctor	

#### Other information

### Drug Facts (continued)

#### Inactive ingredients

carboxymethylcellulose sodium, citric acid, D&C red no. 28, flavor, glycerin, microcrystalline cellulose, purified water, sodium citrate, sodium hypochlorite, sucrose, xanthan gum

Questions or comments? 1-800-540-3765

\*This product is not manufactured or distributed by Bayer Healthcare LLC, the owner of the registered trademark Phillips ®

DIST. BY MELJER DISTRIBUTION, INC. GRAND RAPIDS, MI 48544





### **MILK OF MAGNESIA**

magnesium hydroxide suspension

# **Product Information**

**HUMAN OTC DRUG** NDC:79481-0648 **Product Type Item Code (Source)** 

**ORAL Route of Administration** 

# **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION -	MAGNESIUM HYDROXIDE	1200 mg

# **Inactive Ingredients**

Ingredient Name	Strength
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
GLYCERIN (UNII: PDC6A3C0OX)	
XANTHAN GUM (UNII: TTV12P4NEE)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
WATER (UNII: 059QF0KO0R)	
SODIUM HYPOCHLORITE (UNII: DY38VHM5OD)	
SUCROSE (UNII: C151H8M554)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Product Characteristics		
Color		Score
Shape		Size
Flavor	CHERRY	Imprint Code
Contains		

ı	Packaging				
	#	# Item Code Package Description		Marketing Start Date	Marketing End Date
	1	NDC:79481- 0648-1	769 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/01/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	02/01/2021	

# **Labeler -** Meijer, Inc. (006959555)

# **Registrant -** GCP Laboratories (965480861)

Establishment				
Name	Address	ID/FEI	Business Operations	
GCP Laboratories		965480861	manufacture(79481-0648)	

Revised: 11/2023 Meijer, Inc.