

**CREAM- titanium dioxide, zinc oxide lotion**  
**OXYGEN DEVELOPMENT, LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**LIGHT-SKIN TREAT BLURRING SKIN TINT BROAD SPECTRUM SPF 20**  
**SUNSCREEN -**

***Active Ingredients***

Titanium Dioxide 4.55%

Zinc Oxide 3.5%

***Purpose***

Sunscreen

***Uses***

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see Directions) decreases the risk of skin cancer and early skin aging caused by the sun.

***Warnings***

**For external use only**

**Stop use and ask doctor** if rash occurs.

**Do not use** on damaged or broken skin.

**When using this product** keep out of eyes. Rinse with water to remove.

**Keep out of the reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water-resistant sunscreen of swimming or sweating.
- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.
- **Sun Protection Measures.** Spending time in the sun increases our risk of skin cancer and early skin aging. To decrease this risk regularly use a sunscreen with a Broad-Spectrum SPF value of 15 or higher and other sun protection measures including:

Limit time in the sun, especially from 10 a.m.- 2p.m.

Wear long-sleeved shirts, pants, hats and sunglasses.

### Other information

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.

### Inactive Ingredients

Cyclopentasiloxane, Isododecane, Mica, Polysilicone-11, Polymethylsilsesquioxane, Hexyl Laurate, PEG-10 Dimethicone, Polyglyceryl-4 Isostearate, Stearic Acid, Cetyl PEG/PPG-10/1 Dimethicone, Alumina, Triethoxycaprylylsilane, Dipalmitoyl Hydroxyproline, Diamond Powder, Iron Oxides.

### Primary Package (Inner)



F: front panel width      B: back panel width  
CW: circumferential width

F: front panel length (Circumferential length)

L: tube length (from shoulder to bottom)  
 Eyemark (3×6mm): mark for end sealing  
 Upper edge(7mm): space for end sealing  
 Bottom edge(5mm): non-printing space

Dia 25×L110	F/B 39.25mm	CW 78.5mm	L(changeable) 110mm
shadow area:	printing area		

Attention of artwork: Use CorelDraw,Freehand or Illustrator, and do not compress or change the file's type, then outline the file and send us.

sugar  rush	
SUB-BRAND: SUGAR RUSH	FILE NAME: SKIN TREAT
COMPONENT: PRIMARY	SHADE: SPF
DATE OPENED: 5/4/18	DATE REVISED: 10/19/18
REASON FOR CHANGE: PID REVISED	
<b>COLORS</b>	
■ BLACK	■ 182C
■ 225C	■ 304C

## Secondary Package (Outer)

30ml, 61354-51-14

**Drug Facts**  
(continued)

**Inactive Ingredients**

Cyclopentasiloxane, isododecane, polysilicone-11, polymethylsilsesquioxane, hexyl laurate, PEG-10 dimethicone, polyglyceryl-4 isostearate, mica, stearic acid, cetyl PEG/PPG-10/1 dimethicone, alumina, triethoxycaprylylsilane, dipalmitoyl hydroxyproline, diamond powder, iron oxides.  
OX100FZ000081.

LOT: 0453  
EXP: 05/21

net 30 mL / 1 FL Oz. Made in USA. Dist. by tarte, Inc. NY 10018. All rights reserved. Biorius Sprl, Rue de la Croeyere 10A, B-7170 Manage, BE.



light

sugar  rush  
tarte™

*skin  
treat*

@sugarrush  
sugarrushbeauty.com



blurring skin tint  
Broad Spectrum  
SPF 20 Sunscreen

**Drug Facts**

**Active Ingredients**

Titanium Dioxide 4.55%... Sunscreen  
Zinc Oxide 3.50%... Sunscreen

**Purpose**

**Uses**  
• Helps prevent sunburn.  
• If used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun.

**Warnings**

For external use only.

Stop use and ask doctor if rash occurs.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Keep out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- Shake before use.
- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- Children under 6 months: Ask a doctor.
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
  - Limit time in the sun, especially from 10 a.m.-2 p.m.
  - Wear long-sleeved shirts, pants, hats and sunglasses

**Other information**

- Protect the product in this container from excessive heat and direct sunlight.
- You may report a serious adverse reaction to: tarte c/o Report Reaction, LLC, P.O. Box 22, Plainsboro, NJ 08536-0222.



SP08821

# CREAM

titanium dioxide, zinc oxide lotion

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:61354-051
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	3.5 mg in 100 mg
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	4.55 mg in 100 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>POLYMETHYLSILSESQUIOXANE (11 MICRONS)</b> (UNII: Z570VEV8XK)	
<b>HEXYL LAURATE</b> (UNII: 4CG9F9W01Q)	
<b>ISODODECANE</b> (UNII: A8289P68Y2)	
<b>MICA</b> (UNII: V8A1AW0880)	
<b>POLYGLYCERYL-4 ISOSTEARATE</b> (UNII: 820DPX33S7)	
<b>PEG-10 DIMETHICONE (600 CST)</b> (UNII: 8PR7V1SVM0)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE)</b> (UNII: 9E4CO0W6C5)	
<b>CYCLOMETHICONE 5</b> (UNII: 0THT5PCI0R)	

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:61354-051-14	1 in 1 CARTON	02/23/2021	
1		30 mg in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part352	02/23/2021	

**Labeler** - OXYGEN DEVELOPMENT, LLC (137098492)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
OXYGEN DEVELOPMENT, LLC		137098492	manufacture(61354-051)

Revised: 2/2023

OXYGEN DEVELOPMENT, LLC