

**NEW LIFE 3% HYDROGEN PEROXIDE FIRST AID ANTISEPTIC- hydrogen peroxide solution**  
**New Life Products, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**new life™ 3% Hydrogen Peroxide**

**Drug Facts**

**Active Ingredient**

Hydrogen Peroxide(stabilized) 3%

**Purpose**

First aid antiseptic/oral debriding agent

**Uses**

• First aid to help prevent the risk of infection in minor cuts, scrapes and burns • aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth.

**Warnings**

• **For external use only. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.**

**Do not use:** In the eyes or apply over large areas of the body • Longer than one week

**Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns.**

**Stop use and ask a doctor if** swelling, rash, or fever develops • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, redness persists or worsens

**KEEP OUT OF REACH OF CHILDREN.**

**Directions**

• **For use as a first aid antiseptic:**

• clean affected area • apply small amount of product on affected area 1 to 3 times a day • if bandaged, let dry first • may be covered with a sterile bandage

**For use as an oral debriding agent (oral rinse):**

- adults and children 2 years of age and over:
- mix with an equal amount of water
- swish around in the mouth over the affected area for at least 1 minute and then spit out. Do not swallow.
- use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor

**Children under 12 years should be supervised in the use of this product.**

**Children under 2 years of age: consult a doctor or dentist.**

**Other Information**

- Keep tightly closed and in a dark place at room temperature • Do not shake bottle. Hold away from face when opening.

**Inactive Ingredients:**

Purified water.

***Topical Solution USP***

**FIRST AID ANTISEPTIC**

- + **Treats minor cuts and abrasions**
- + **Oral debriding agent**
- + **Use as a gargle or rinse**

**DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**New Life Products Inc.** • Chicago, IL 60629

Watford WD24 7GN, UK • 800-262-2326

[www.newlifesanitizer.com](http://www.newlifesanitizer.com)

ISO 9001, 22716 and cGMP compliant

**MADE IN USA**

**Packaging**

PACKAGE - 118 mL



# 3% Hydrogen Peroxide Topical Solution USP

## FIRST AID ANTISEPTIC

- + Treats minor cuts and abrasions
- + Oral debriding agent
- + Use as a gargle or rinse


DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

4 Fl. Oz e 118 mL

Drug Facts	Purpose
<b>Active Ingredient</b> Hydrogen Peroxide (activated) 3%.....	First aid antiseptic/oral debriding agent
<b>Uses</b> • First aid to help prevent the risk of infection in minor cuts, scrapes, and burns • aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth.	
<b>Warnings</b> • For external use only. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away. <b>KEEP OUT OF REACH OF CHILDREN.</b>	
<b>Do not use:</b> In the eyes or apply over large areas of the body • Longer than one week	
<b>Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns.</b>	
<b>Stop use and ask doctor if</b> swelling, rash, or fever develops • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, redness persists or worsens	
<b>Directions</b> • For use as a first aid antiseptic: • clean affected area • apply small amount of product on affected area 1-3 times a day • if bandaged, let dry first • may be covered with a sterile bandage	
<b>For use as an oral debriding agent (oral rinse):</b> • adults and children 2 years of age and over: • mix with an equal amount of water • swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. • use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor	
<b>Children under 12 years should be supervised in the use of this product. Children under 2 years of age: consult a doctor or dentist.</b>	
<b>Other information</b> • Keep tightly closed and in a dark place at room temperature • Do not shake bottle. Hold away from face when opening.	
<b>Inactive ingredients:</b> Purified water.	

Item # NL9445 NDC #  
**New Life Products Inc.** • Chicago, IL 60629  
 Watford WD24 7GN, UK • 800-262-2326  
[www.newlifesanitizer.com](http://www.newlifesanitizer.com)  
 ISO 9001, 22716 and cGMP compliant





**3%**  
**Hydrogen Peroxide**  
*Topical Solution USP*

**FIRST AID ANTISEPTIC**

- + Treats minor cuts and abrasions
- + Oral debriding agent
- + Use as a gargle or rinse

**DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

16 Fl. Oz e 473 mL

Drug Facts	Purpose
<b>Active Ingredient</b> Hydrogen Peroxide(stabilized) 3%.....	First aid antiseptic/oral debriding agent
<b>Uses</b> • First aid to help prevent the risk of infection in minor cuts, scrapes, and burns • aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth.	
<b>Warnings</b> • For external use only. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away. <b>KEEP OUT OF REACH OF CHILDREN.</b>	
<b>Do not use:</b> In the eyes or apply over large areas of the body • Longer than one week	
<b>Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns.</b>	
<b>Stop use and ask doctor if</b> swelling, rash, or fever develops • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, redness persists or worsens	
<b>Directions</b> • For use as a first aid antiseptic: • clean affected area • apply small amount of product on affected area 1-3 times a day • if bandaged, let dry first • may be covered with a sterile bandage	
<b>For use as an oral debriding agent (oral rinse):</b> • adults and children 2 years of age and over: • mix with an equal amount of water • swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. • use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor	
<b>Children under 12 years should be supervised in the use of this product. Children under 2 years of age: consult a doctor or dentist.</b>	
<b>Other Information</b> • Keep tightly closed and in a dark place at room temperature • Do not shake bottle. Hold away from face when opening.	
<b>Inactive Ingredients:</b> Purified water.	

Item # NL9184      NDC #  
**New Life Products Inc.** • Chicago, IL 60629  
 Watford WD24 7GN, UK • 800-262-2326  
 www.newlifesanitizer.com  
 ISO 9001, 22716 and cGMP compliant



PACKAGE - 946 mL

<b>Drug Facts</b>	<b>Purpose</b>
<b>Active Ingredient</b> Hydrogen Peroxide(stabilized) 3%.....	First aid antiseptic/oral debriding agent
<b>Uses</b> • First aid to help prevent the risk of infection in minor cuts, scrapes, and burns • aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth.	
<b>Warnings</b> • <b>For external use only. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away. KEEP OUT OF REACH OF CHILDREN.</b>	
<b>Do not use:</b> In the eyes or apply over large areas of the body • Longer than one week	
<b>Ask a doctor before use if you have deep or puncture wounds, animal bites or serious burns.</b>	
<b>Stop use and ask doctor if</b> swelling, rash, or fever develops • the condition persists or gets worse • sore mouth symptoms do not improve in 7 days • irritation, pain, redness persists or worsens	
<b>Directions</b> • <b>For use as a first aid antiseptic:</b> • clean affected area • apply small amount of product on affected area 1-3 times a day • if bandaged, let dry first • may be covered with a sterile bandage	
<b>For use as an oral debriding agent (oral rinse):</b> • adults and children 2 years of age and over: • mix with an equal amount of water • swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. • use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor	
<b>Children under 12 years should be supervised in the use of this product. Children under 2 years of age; consult a doctor or dentist.</b>	
<b>Other Information</b> • Keep tightly closed and in a dark place at room temperature • Do not shake bottle. Hold away from face when opening.	
<b>Inactive Ingredients:</b> Purified water.	

Item # NL9185    NDC #  
**New Life Products Inc.** • Chicago, IL 60629  
 Watford WD24 7GN, UK • 800-262-2326  
 www.newlifesanitizer.com  
 ISO 9001, 22716 and cGMP compliant

## NEW LIFE 3% HYDROGEN PEROXIDE FIRST AID ANTISEPTIC

hydrogen peroxide solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49681-016
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>HYDROGEN PEROXIDE</b> (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	3 mL in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

**WATER** (UNII: 059QF0KO0R)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49681-016-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/05/2021	
2	NDC:49681-016-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/05/2021	
3	NDC:49681-016-32	946 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/05/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	02/05/2021	

**Labeler** - New Life Products, Inc. (117815140)

### Establishment

Name	Address	ID/FEI	Business Operations
New Life Products, Inc.		117815140	manufacture(49681-016)

Revised: 2/2023

New Life Products, Inc.