COLD AND FLU RELIEF DAYTIME- acetaminohpen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled QUALITY CHOICE (Chain Drug Marketing Association)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethoprhan HBr 10 mg

Phenlyephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - fever
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-

800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients

butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow#6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

Call 1-800-935-2362 Monday-Friday 9AM-5PM EST

Principal Display Panel

†Compare to the active ingredients in VICKS® DAYQUIL® COLD & FLU LIQUILCAPS® Non-Drowsy

Daytime

Multi-Symptom Relief For Cold/Flu

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

Pain Reliever | Fever Reducer

Cough Suppressant | Nasal Decongestant

Non-drowsy | Alcohol-free | Antihistamine-free

Softgels

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of the Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Distributed by C.D.M.A., Inc.© 43157 W. 9 Mile Rd Novi, MI 48376-0995 www.qualitychoice.com

Package Label

When using this product, do not exceed recommended dosage.

thinning drug warfarin.

Ask a doctor or pharmacist before use if you are taking the blood

- condu trat occurs with too much phiegm (mucus) ешриуѕета
- bersistent or chronic cough such as occurs with smoking, asthma, or ■ trouble unnating due to an enlarged prostate gland
- µiðµ pjoog bussanus ■ heart disease ■ thyroid disease Ask a doctor before use if you have miliver disease malabetes

pharmacist before taking this product. do not know if your prescription drug contains an MAOI, ask a doctor or

- Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you (certain drugs for depression, psychiatric, or emotional conditions, or If you are now taking a prescription monoamine oxidase inhibitor (MAOI)
 - acetaminophen, ask a doctor or pharmacist. nonprescription). If you are not sure whether a drug contains
 - with any other drug containing acetaminophen (prescription or

nomiting, consult a doctor promptly.

days, is accompanied or followed by fever, headache, rash, nausea, or Sore throat warning: If sore throat is severe, persists for more than 2 It a skin reaction occurs, stop use and seek medical help right away. may include:

skin reddening

bilsters

rash

- Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms
 - 3 or more alcoholic drinks every day while using this product
 - with other drugs containing acetaminophen
 - more than 4,000 mg of acetaminophen in 24 hours
 - may occur if you take:

Liver warning: This product contains acetaminophen. Severe liver damage Warnings

Drug Facts (continued)

Call 1-800-935-2362 Monday-Friday 9AM-5PM EST Questions or comments?

povidone, propylene glycol, purfied water, sorbitan, sorbitol, white ink hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyetrylene glycol, INACTIVE INGREDIES butylated hydroxyanisole, butylated

> ■ store between 15-30°C (59-86°F)
> ■ avoid excessive heat поиътиоти тэпи

label to ensure correct dosing

- when using other Daytime or Nighttime products, carefully read each
 - children under 12 years: do not use
 - m swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 4 hours ■ do not take more than 4 doses in 24 hours

DIRECTIONS = do not take more than directed (see Overdose warning)

adults as well as for children even if you do not notice any signs or symptoms. Center (1-800-222-1222) right away. Quick medical attention is critical for damage. In case of overdose, get medical help or contact a Poison Control Overdose warning: Taking more than the recommended dose can cause liver Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use.

These could be signs of a serious condition.

- cond) comes pack or occurs with rash or headache that lasts.
 - redness or swelling is present new symptoms occur
 - fever gets worse or lasts more than 3 days
- uervousness, dizziness, or sleeplessness occur
- bain, cough, or nasal congestion gets worse or lasts more than 7 days Stop use and ask a doctor if

Drug Facts (continued)

Nasal decongestant Phenylephrine HCl 5 mg. nondu auppressant Dextromethorphan HBr 10 mg. Pain reliever/fever reducer Acetaminophen 325 mg Active ingredients (in each softgel) Purposes Drug Facts

 cough due to minor throat and bronchial irritation ■ usaal congestion ■ fever ■ minor aches and pains ■ headache ■ sore throat NSGS = temporarily relieves common cold and flu symptoms:

Drug Facts (continued)

NDC 63868-409-24



†Compare to the Active Ingredients in VICKS® DAYQUIL®

COLD & FLU LIQUICAPS®

DayTime

Non-Drowsy

Multi-Symptom Relief For Cold/Flu

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCI

Pain Reliever | Fever Reducer Cough Suppressant | Nasal Decongestant

Non-Drowsy | Alcohol Free | Antihistamine Free



KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMAT

EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

actual size

[†]This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.







QUALITY CHOICE Non-Drowsy Daytime Cold and Flu Relief

COLD AND FLU RELIEF DAYTIME

acetaminohpen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information	
	n

Product Type HUMAN OTC DRUG Item Code (Source) NDC:63868-409

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

-		
Ingredient Name	Strength	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)		
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)		

MANNITOL (UNII: 30WL53L36A)

Inactive Ingredients

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)

GELATIN (UNII: 2G86QN327L)
GLYCERIN (UNII: PDC6A3C0OX)

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3MJQ0SDW1A)

POVIDONE (UNII: FZ989GH94E)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

WATER (UNII: 059QF0KO0R)
SORBITAN (UNII: 6092ICV9RU)
SORBITOL (UNII: 506T60A25R)

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	20mm
Flavor		Imprint Code	P19

Contains

Packaging						
# Item Code	Package Description	Marketing Start Date	Marketing End Date			
NDC:63868-409- 24	24 in 1 CARTON	09/30/2019				
1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product					

Marketing InformationMarketing CategoryApplication Number or Monograph CitationMarketing Start DateMarketing End DateOTC monograph finalpart34109/30/2019

Labeler - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 5/2023 QUALITY CHOICE (Chain Drug Marketing Association)