

IBUPROFEN- ibuprofen tablet, film coated
Walgreen Company

Walgreens 44-392

Active ingredient (in each orange tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- any new symptoms appear
- redness or swelling is present in the painful area

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, cellulose, corn starch, FD&C yellow #6 aluminum lake, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or Comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

Well at

Walgreens

WALGREENS PHARMACIST RECOMMENDED[□]

NDC 0363-0392-14

Ibuprofen 200

Ibuprofen Tablets

USP, 200 mg / Pain Reliever /

Fever Reducer (**NSAID**)

500

TABLETS

Compare to Motrin[®] IB Tablets active ingredient^{□□}

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

[□]Walgreens Pharmacist Survey Study, November 2014.

^{□□}This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Motrin[®] IB Tablets.


50844 ORG041139214 ORG0515-F

DISTRIBUTED BY: WALGREEN CO. 200 WILMOT RD. DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com

© 2015 Walgreen Co.




WALGREENS PHARMACIST RECOMMENDED*

Ibuprofen 200

Ibuprofen Tablets
USP, 200 mg / Pain Reliever /
Fever Reducer (NSAID)

500

TABLETS



Actual Size

Compare to Motrin® IB Tablets active ingredient**

UNVARNISHED AREA
LOT NO. & EXP. DATE

NDC 0363-0392-14

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL
UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each orange tablet) Purpose
Ibuprofen USP 200 mg (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - minor pain of arthritis
- headache
- backache
- toothache
- menstrual cramps
- muscular aches
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other pain reliever/fever reducer


† Walgreens Pharmacist Survey Study, November 2014.

** This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark, Motrin® IB Tablets.

50844 ORG041139214

ORG0515-F

ITEM 907106 W10048-0615-F



DISTRIBUTED BY:
WALGREEN CO.
200 WILMOT RD
DEERFIELD, IL 60015
100% SATISFACTION
GUARANTEED
walgreens.com
©2015 Walgreen Co.

PEEL HERE FOR MORE DRUG FACTS

FRONT (S1)

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin or heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk, if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- any new symptoms appear
- redness or swelling is present in the painful area

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless it is definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20-25°C (68-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients carnauba wax, cellulose, corn starch, FD&C yellow #6 aluminum lake, humed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-800-425-9391
8:30 AM-4:00 PM ET, Monday-Friday

STOP PEELING

BACK (S2)

Walgreens 44-392

IBUPROFEN			
ibuprofen tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0392
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg
Inactive Ingredients			

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	44;392
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0392-15	1 in 1 CARTON	03/30/2002	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0392-12	1 in 1 CARTON	03/30/2002	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0363-0392-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/30/2002	
4	NDC:0363-0392-16	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/30/2002	
5	NDC:0363-0392-17	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/30/2002	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075139	03/30/2002	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(0363-0392)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(0363-0392)