

**IBUPROFEN- ibuprofen tablet, film coated**  
**Walgreen Company**

-----  
**Walgreens 44-392**

***Active ingredient (in each orange tablet)***

Ibuprofen USP, 200 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - menstrual cramps
  - toothache
  - the common cold
  - backache
  - headache
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin reddening
- asthma (wheezing)
- rash
- facial swelling
- shock
- blisters
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart

failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- right before or after heart surgery
- if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
  - chest pain
  - trouble breathing
  - leg swelling
  - slurred speech
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist

- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### ***Other information***

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

### ***Questions or Comments?***

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

### ***Principal Display Panel***

Well at

**Walgreens**

WALGREENS PHARMACIST RECOMMENDED<sup>†</sup>

NDC 0363-0392-14

**Ibuprofen**

**200**

**Ibuprofen** Tablets USP, 200 mg /  
Pain Reliever / Fever Reducer (**NSAID**)

**50**

**TABLETS**

**Compare to Motrin<sup>®</sup> IB Tablets**  
**active ingredient<sup>‡‡</sup>**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED**  
**SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

<sup>†</sup>Walgreens Pharmacist Survey Study, November 2016.

<sup>‡‡</sup>This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin<sup>®</sup> IB Tablets.

50844 REV1116A39215

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015

**100% SATISFACTION GUARANTEED**

walgreens.com ©2017 Walgreen Co.

**Drug Facts (continued)**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- store between 20-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

**Inactive ingredients** carnauba wax, colloidal silicon dioxide, corn starch, FD&C yellow #6 aluminum lake, hypromellose, lactose anhydrous

**Drug Facts (continued)**

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- right before or after heart surgery
- if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

**Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke
- chest pain
- leg swelling
- slurred speech
- weakness in one part or side of body

**Drug Facts (continued)**

magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, sodium starch glycolate, stearic acid, titanium dioxide

**(Questions or comments?)**  
Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

\*Walgreens Pharmacist Survey Study, November 2016.  
\*\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Motrin® IB Tablets.  
50644 REV1116A39215

**Drug Facts**

**Active ingredient**  
Ibuprofen USP 200 mg (NSAID)\* (in each orange tablet)

**Other ingredients**  
monobasic anti-inflammatory drug

**Uses**

- temporarily relieves minor aches and pains due to:
  - menstrual cramps
  - toothache
  - backache
  - headache
  - muscular aches
- minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- skin redness
- facial swelling
- rash
- hives
- asthma (wheezing)
- shock
- anisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

NDC 0363-0392-15

**Ibuprofen**

**200**

WALGREENS PHARMACIST RECOMMENDED!

**Ibuprofen Tablets USP 200 mg / Pain Reliever / Fever Reducer (NSAID)**

**50 TABLETS**

Compare to Motrin® IB Tablets active ingredient\*\*

ORGI1012-F\_R01

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

DISTRIBUTED BY: WALGREEN CO.  
200 WILLMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2017 Walgreen Co.  
ITEM 500730  
11917 089333  
3 1

No Print Area  
Lot no. & Exp. Date

B-2201-392-15-R  
REV1116A39215

No Print  
Glue Area

No Print  
Glue Area

Walgreens 44-392

**IBUPROFEN**

ibuprofen tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-0392
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name		Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg	
<b>Inactive Ingredients</b>				
Ingredient Name		Strength		
CARNAUBA WAX (UNII: R12CBM0EIZ)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
STARCH, CORN (UNII: O8232NY3SJ)				
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
<b>Product Characteristics</b>				
Color	ORANGE	Score	no score	
Shape	ROUND	Size	10 mm	
Flavor		Imprint Code	44;392	
Contains				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0392-15	1 in 1 CARTON	03/01/1999	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0363-0392-12	1 in 1 CARTON	03/01/1999	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0363-0392-37	1 in 1 CARTON	03/01/1999	
3		75 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0363-0392-14	500 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/1999	
5	NDC:0363-0392-16	1000 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/1999	
6	NDC:0363-0392-17	300 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/1999	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA075139	03/01/1999		

**Labeler** - Walgreen Company (008965063)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		038154464	PACK(0363-0392)

### **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867894	MANUFACTURE(0363-0392)

### **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867837	PACK(0363-0392)

### **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		967626305	PACK(0363-0392)

### **Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		868734088	PACK(0363-0392)