

EPHEDRINE HYDROCHLORIDE- ephedrine hydrochloride tablet
Walgreen Company

1298P-Walgreen

Drug Facts

Active ingredient (in each tablet)

Ephedrine Hydrochloride 12.5 mg

Purpose

Bronchodilator

Use

☐ for temporary relief of mild symptoms of intermittent asthma: ☐ wheezing ☐ tightness of chest ☐ shortness of breath

Asthma alert: Because asthma may be life threatening, see a doctor if you

☐ are not better in 60 minutes

☐ get worse

☐ need more than 12 tablets in 24 hours

☐ use more than 8 tablets in 24 hours for 3 or more days a week

☐ have more than 2 asthma attacks in a week

These may be signs that your asthma is getting worse.

☐ This product will not give you asthma relief as quickly as an inhaled bronchodilator

Do not use

☐ unless a doctor said you have asthma

☐ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs taken for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

☐ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

☐ ever been hospitalized for asthma

☐ heart disease

☐ high blood pressure

☐ diabetes

☐ thyroid disease

☐ seizures

☐ narrow angle glaucoma

- a psychiatric or emotional condition
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking prescription drugs for asthma, obesity, weight control, depression, or psychiatric or emotional conditions
- taking any drug that contains phenylephrine, pseudoephedrine, ephedrine, or caffeine (such as for allergy, cough-cold, or pain)

When using this product

your blood pressure or heart rate may go up. This could increase your risk of heart attack or stroke, which may cause death.

your risk of heart attack or stroke increases if you:

have a history of high blood pressure or heart disease

take this product more frequently or take more than the recommended dose

avoid foods or beverages that contain caffeine

avoid dietary supplements containing ingredients reported or claimed to have a stimulant effect

Stop use and ask a doctor if

- your asthma is getting worse (see Asthma alert)
- you have difficulty sleeping
- you have a rapid heart beat
- you have tremors, nervousness, or seizure

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed

adults and children 12 years of age and over	Take 1-2 tablets every 4 hours as needed. Do not take more than 12 tablets in 24 hours
children under 12 years of age	Ask a doctor

Other information

- store in a dry place at 15° – 30°C (59° – 86°F).

crospovidone, D&C yellow #10 lake, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid.

Questions or comments?

Call 1-800-925-4733

TAMPER EVIDENT: DO NOT USE IF CARTON HAS BEEN OPENED OR BLISTER UNITS ARE TORN OR BROKEN

RETAIN THIS CARTON FOR COMPLETE PRODUCT INFORMATION

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

††This product is not manufactured or distributed by ARMSTRONG PHARMACEUTICALS, INC., owner of the registered trademark Primatene®.

Walgreens

NDC 0363-9812-07

Compare to the active ingredient in Primatene®††

Asthma Relief

EPHEDRINE HCl 12.5 mg / BRONCHODILATOR

- Relief of mild intermittent asthma symptoms
- Restores free breathing

60 Tablets





EPHEDRINE HYDROCHLORIDE

ephedrine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9812
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EPHEDRINE HYDROCHLORIDE (UNII: NLJ6390P1Z) (EPHEDRINE - UNII:GN83C131XS)	EPHEDRINE HYDROCHLORIDE	12.5 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

STEARIC ACID (UNII: 4ELV7Z 65AP)

CROSPROVIDONE (UNII: 2S7830E561)

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	1298
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9812-07	4 in 1 CARTON	12/02/2021	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/02/2021	

Labeler - Walgreen Company (008965063)

Revised: 12/2023

Walgreen Company