

MAJOR NASAL DECONGESTANT- oxymetazoline hydrochloride spray

Major Pharmaceuticals

Major Nasal Decongestant Spray No Drip

Drug facts

Active Ingredient

Oxymetazoline 0.05%

Purpose

Nasal Decongestant

Uses

Temporarily relieves nasal congestion due to:

- common cold
- hay fever
- upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- Shrinks swollen membranes so you can breathe more freely

Warnings

Ask a Doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to enlarged prostate gland.

When using this product

Do not use more than directed

- do not use for more than three days, Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur.
- use of this container by more than one person may spread infection.

Stop use and ask a Doctor if

Symptoms persist

if pregnant or breast-feeding

ask a health professional before use.

Keep out of the reach of children

if swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Before using for the first time, remove the protective cap from the tip and prime the pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm and even stroke, and sniff deeply.
- Wipe nozzle clean after use.
- Adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- Children under 6 years of age consult a doctor.

Other information

- store between 20° - 25°C (68° to 77°F)
- retain carton for future reference on full labeling

Inactive ingredients

Benzalkonium Chloride, Disodium EDTA, Glycerin, Microcrystalline Cellulose and Carboxymethylcellulose Sodium, Polyethylene Glycol 300, Povidone, Propylene Glycol, Purified Water, Sodium Phosphate Dibasic, Sodium Phosphate Monobasic

Questions? Call 800-222-1807

Principal Display Panel

Major ® No Drip Nasal Mist NDC 0904-7427-30
Nasal Spray Oxymetazoline HCL 0.05%
12 hour Nasal Spray
1 FL OZ (30 mL)

MAJOR
12 Hour
Nasal Decongestant Spray
Oxymetazoline HCl 0.05%
0.05%

Child Resistant Cap
(see side panel)

DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

Drug Facts

Active ingredient Oxymetazoline hydrochloride 0.05% **Purpose** Nasal decongestant

Uses
 • temporarily relieves nasal congestion due to:
 • common cold • hay fever • upper respiratory allergies
 • temporarily relieves sinus congestion and pressure
 • shrinks swollen membranes so you can breathe more freely

Warnings
 Ask a doctor before use if you have
 • heart disease • high blood pressure
 • thyroid disease • diabetes
 • trouble urinating due to an enlarged prostate gland

When using this product
 • do not use more than directed
 • do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
 • temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
 • use of this container by more than one person may spread infection

Stop use and ask a doctor if symptoms persist

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
 • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
 • children under 6 years of age: ask a doctor
 To Use: Shake well before use. Before using the first time, prime metered pump by depressing pump firmly several times. Squeeze both sides of the cap and twist counterclockwise. Align the lines on the groove at the base of the pump as shown in picture. Without tilting head, insert nozzle into nostril. Depress and spray. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use. Secure cap after use.

Drug Facts (continued)

Other information
 • Store at a controlled room temperature 20°-25°C (68°-77°F)
 • retain carton for future reference on full labeling

Inactive ingredients
 Benzalkonium Chloride, Edetate Disodium, Glycerin, Microcrystalline Cellulose and Carboxymethylcellulose Sodium, Polyethylene Glycol 300, Povidone, Propylene Glycol, Sodium Phosphate Dibasic, Sodium Phosphate Monobasic, Water (Purified)

Questions? Call 800-222-1807

IMPORTANT: KEEP OUTER CARTON FOR FUTURE REFERENCE ON FULL LABELING

Distributed by:
MAJOR® PHARMACEUTICALS
 Indianapolis, IN 46268
 Questions or comments?
 Call (800) 616-2471
 www.majorpharmaceuticals.com

This product is not manufactured or distributed by ©Bayer HealthCare LLC, owner of the registered trademark Afrin® Original No Drip.

MADE IN THE U.S.A. OF U.S. AND IMPORTED INGREDIENTS AND COMPONENTS

MAJOR®

NDC 0904-7427-30
 Compare to the active ingredient in Afrin® Original No Drip®

12 Hour Nasal Decongestant Spray

Oxymetazoline HCl 0.05%

0.05%


Won't drip from nose or down throat

Relief of:
 Nasal Congestion
 Sinus Pressure & Congestion

1 FL OZ (30 mL)



Directions: Squeeze both sides of the cap and twist counterclockwise. Align the lines on the groove at the base of the pump as shown in picture. Depress and spray.



Package Contains One Bottle

Actual Size

560110

LOT

EXP

Rev. 10/23 M-101 Re-order No. 701386



NDC 0904-7427-30

MAJOR[®]

12 Hour

**Nasal Decongestant
Spray**

Oxymetazoline HCl 0.05%

0.05%**1 FL OZ (30 mL)****SEE CARTON FOR FULL LABELING****DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING**

Active ingredient Oxymetazoline hydrochloride 0.05%

Uses • See Carton.

Warnings Ask a doctor before use if you have

• heart disease • high blood pressure • thyroid disease

• diabetes • trouble urinating due to an enlarged prostate

gland. When using this product • do not use more than

directed • do not use for more than 3 days. Use only as

directed. Frequent or prolonged use may cause nasal

congestion to recur or worsen. • Temporary discomfort

such as burning, stinging, sneezing or an increase in nasal

discharge may occur • use of this container by more than

one person may spread infection. Stop use and ask a

doctor if symptoms persist. If pregnant or

breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical

help or contact a Poison Control Center (1-800-222-1222)

right away. Directions • adults and children 6 to under

12 years of age (with adult supervision): 2 or 3 sprays in

each nostril not more often than every 10 to 12 hours. Do

not exceed 2 doses in any 24-hour period • children under

6 years of age: ask a doctor. To use SHAKE WELL

BEFORE USE. Squeeze both sides of the cap and

twist counterclockwise. Align the lines on the

groove at the base of the pump as shown in

picture. Depress and spray. Secure cap after use.

Store between 20°-25°C (68°-77°F)

Questions? Call 800-222-1807

Distributed by:

MAJOR[®] PHARMACEUTICALS

Indianapolis, IN 46268

Questions or comments?

Call (800) 616-2471

www.majorpharmaceuticals.com

MADE IN THE U.S.A. OF U.S. AND IMPORTED

INGREDIENTS AND COMPONENTS

Rev. 10/23 M-101 Re-order No. 701386

3

Lot and Exp Area

MAJOR NASAL DECONGESTANT

oxymetazoline hydrochloride spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-7427
Route of Administration	NASAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYMETAZOLINE HYDROCHLORIDE (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZOLINE HYDROCHLORIDE	0.5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 300 (UNII: 5655G9Y8AQ)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SODIUM PHOSPHATE, MONOBASIC (UNII: 3980JH2SW)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0904-7427-30	1 in 1 CARTON	12/31/2023	
1		30 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/31/2023	

Labeler - Major Pharmaceuticals (191427277)

Revised: 12/2023

Major Pharmaceuticals