

**DEXTROMETHORPHAN POLISTIREX EXTENDED RELEASE- dextromethorphan polistirex suspension  
Bryant Ranch Prepack**

-----  
**Perrigo Dextromethorphan Polistirex Extended-Release Oral Suspension Drug Facts**

**Active ingredient (in each 5 mL)**

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide

**Purpose**

Cough suppressant

**Uses**

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Allergy Alert:** Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

**Ask a doctor before use if you have**

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

**Stop use and ask a doctor if**

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **shake bottle well before use**
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by doctor

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

### **Other information**

- **each 5 mL contains:** sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- dosing cup provided

### **Inactive ingredients**

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

### **Questions or comments?**

**1-800-719-9260**

### **HOW SUPPLIED**

Dextromethorphan Polistirex Extended-Release Oral Suspension

- NDC 72162-2281-2: 89 mL in a BOTTLE

Repackaged/Relabeled by:  
Bryant Ranch Prepack, Inc.

Burbank, CA 91504

# Dextromethorphan Polistirex Extended-Release Oral Suspension



GTIN 00363629106424  
 Lot L0T123  
 Exp 09/12/2025  
 SN 1234567890

Drug Facts	
<b>Active Ingredient (in each 5 mL)</b> Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide	<b>Purpose</b> Cough suppressant
<b>Uses</b> temporarily relieves -cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants -the impulse to cough to help you get to sleep.	
<b>Warnings</b> Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions. Ask a doctor before use if you have -chronic cough that lasts as occurs with smoking, asthma or emphysema -cough that occurs with too much phlegm (mucus) Stop use and ask a doctor if -side effects occur. You may report side effects to FDA at 1-800-FDA-1088. -cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).	
<b>Other Information</b> -each 5 mL contains: sodium 5 mg *store at 20° to 25°C (68° to 77°F) *dosing cup provided. -12 Hour Cough Relief *Day or Night *Alcohol-free *Orange Flavored Liquid.	
<b>Directions</b> -shake bottle well before use -measure only with dosing cup provided -do not use dosing cup with other products -dose as follows or as directed by doctor. (adults and children 12 years of age and over: 10 mL every 12 hours, not to exceed 20 mL in 24 hours)(children 6 to under 12 years of age: 5 mL every 12 hours, not to exceed 10 mL in 24 hours) (children 4 to under 6 years of age: 2.5 mL every 12 hours, not to exceed 5 mL in 24 hours) (children under 4 years of age: do not use).	
<b>Inactive Ingredients</b> D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum.	

NDC 72162-2281-2

## Dextromethorphan Polistirex Extended-Release Oral Suspension

30 mg/5 mL



3 FL OZ (89 mL)

Relabeled by:  
 Bryant Ranch Prepack, Inc.  
 Burbank, CA 91504 USA



### Extended Label

DOSING SHAKE WELL BEFORE USE. Measure only with dosing cup provided. Do not use dosing cup with other products.	
Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

## DEXTROMETHORPHAN POLISTIREX EXTENDED RELEASE

dextromethorphan polistirex suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72162-2281(NDC:45802-433)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII: 7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
POLISTIREX (UNII: 5H9W9GTW27)	
D&C RED NO. 30 (UNII: 2S42T2808B)	

<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)
<b>POLYVINYL ACETATE</b> (UNII: 32K497ZK2U)
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>SODIUM METABISULFITE</b> (UNII: 4VON5FNS3C)
<b>SODIUM POLYSTYRENE SULFONATE</b> (UNII: 1699G8679Z)
<b>SUCROSE</b> (UNII: C151H8M554)
<b>TARTARIC ACID</b> (UNII: W4888I119H)
<b>TRAGACANTH</b> (UNII: 2944357O2O)
<b>TRIACETIN</b> (UNII: XHX3C3X673)
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)

### Product Characteristics

<b>Color</b>	ORANGE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72162-2281-2	1 in 1 CARTON	04/30/2024	
1		89 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	09/10/2012	

**Labeler** - Bryant Ranch Prepack (171714327)

**Registrant** - Bryant Ranch Prepack (171714327)

### Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(72162-2281) , RELABEL(72162-2281)

