ANTACID - aluminum hydroxide, magnesium hydroxide, simethicone liquid Rugby Laboratories

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rugby Regular Strength Antacid Liquid

ACTIVE INGREDIENTS (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel, USP) 400 mg Magnesium hydroxide 400 mg Simethicone 40 mg

PURPOSE

Antacid Antacid Antigas

USE(S)

relieves:

- · acid indigestion
- heartburn
- sour stomach
- upset stomach and gas associated with these symptoms

WARNINGS

Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

ASK A DOCTOR BEFORE USE IF YOU HAVE

- kidnev disease
- a magnesium-restricted diet

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

• presently taking a prescription drug. Antacids may interact with certain prescription drugs.

STOP USE AND ASK DOCTOR

KEEP OUT OF REACH OF CHILDREN

.

DIRECTIONS

- shake well before use
- mL = milliliter
- adults and children 12 years and older: take 10 mL to 20 mL four times a day, or as directed by a doctor
- children under 12 years: consult a doctor

OTHER INFORMATION

- each 10 mL contains: magnesium 170 mg, sodium 5 mg
- store at controlled room temperature 20°C-25°C (68°-77°F)
- do not freeze

INACTIVE INGREDIENTS

ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol

PRINCIPAL DISPLAY PANEL

Rugby

NDC 0536-1293-83

Regular Strength Antacid Liquid

Aluminum hydroxide, 400 mg Magnesium hydroxide, 400 mg Simethicone, 40 mg

Antacid and Antigas

Original Flavor

Relief of:

Heartburn

Acid Indigestion

Sour Stomach

Pressure & Bloating

12 FL OZ (355 mL)

Alcohol content 0.2% v/v



Regular Strength

Antacid

Aluminum Hydroxide, 400 mg Magnesium Hydroxide, 400 mg Simethicone, 40 mg

Antacid and Antigas Original Flavor

Relief of:

Heartburn **Acid Indigestion** Sour Stomach Pressure and Bloating

Alcohol Content 0.2% v/v

Drug Facts UNDER CAP IS BROKEN OR MISSING

DO NOT USE IF PRINTED SEAL

Active ingredients (in each 10 mL) Aluminum hydroxide (equiv. to dried gel, USP) 400 mg... Antacid Magnesium hydroxide 400 mg... .Antacid Simethicone 40 mg..

Uses relieves: ■ acid indigestion ■ heartburn ■ sour stomach upset stomach and gas associated with these symptoms

Warnings

Do not take more than 80 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

Ask a doctor before use if you have ■ kidney disease a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than two weeks

Keep out of reach of children.

Directions ■ shake well before use ■ mL = milliliter ■ adults and children 12 years and older: take 10 mL to 20 mL four times a day, or as directed by a doctor muchildren under 12 years: consult a doctor

Other information ■ each 10 mL contains: magnesium 170 mg, sodium 5 mg ■ store at controlled room temperature 20°C-25°C (68°F-77°F) ■ do not freeze

Inactive ingredients ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol.

Distributed by: RUGBY® LABORATORIES Livonia, MI 48152

Questions or comments? Call 1-800-645-2158 www.rugbylaboratories.com |

Rev. 05/21 R-30 Re-order No. 371027 MFR# 53040



UNVARNISHED AREA I FOR LOT & EXP (LINE DOES NOT PRINT) 1

ANTACID

aluminum hydroxide, magnesium hydroxide, simethicone liquid

Product Information HUMAN OTC DRUG Item Code (Source) NDC:0536-1293 **Product Type Route of Administration ORAL**

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 10 mL	
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	400 mg in 10 mL	
DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	40 mg in 10 mL	

Inactive Ingredients	
Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC10H)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics			
Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging				
#	# Item Code Package Description		Marketing Start Date	Marketing End Date
1	NDC:0536-1293- 83	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/02/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part331	06/02/2021	

Labeler - Rugby Laboratories (079246066)

Establishment					
Name	Address	ID/FEI	Business Operations		
Guardian Drug Company		119210276	MANUFACTURE(0536-1293)		

Revised: 6/2021 Rugby Laboratories