

**BERKLEY AND JENSEN IBUPROFEN CHILDRENS DYE FREE- ibuprofen suspension
BJWC**

BJWC Ibuprofen Oral Suspension USP Drug Facts

Active ingredient (in each 5 mL = 1 teaspoonful)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

**or as directed by a doctor

Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, artificial mixed berry flavor, glycerin, high fructose corn syrup, hypromellose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions or comments?

1-800-934-1204

Principal Display Panel

Compare to the active ingredient in Children's Motrin® Dye-Free

AGES 2 TO 11 YRS

children's

IBUPROFEN ORAL SUSPENSION

100mg PER 5mL

PAIN RELIEVER/FEVER REDUCER (NSAID)

LASTS UP TO 8 HOURS

BERRY FLAVOR

8 FL OZ (240mL)

GLUTEN FREE

DYE-FREE
NON-STAINING
ALCOHOL FREE



Compare to the active ingredient in Children's Motrin® Dye-Free***

Compare to the active ingredient in Children's Motrin® Dye-Free***

berkley jensen®

berkley jensen®

AGES 2 TO 11 YRS

**children's
IBUPROFEN**
ORAL SUSPENSION
100mg PER 5mL

AGES 2 TO 11 YRS

**children's
IBUPROFEN**
ORAL SUSPENSION
100mg PER 5mL

PAIN RELIEVER/FEVER REDUCER (NSAID)

PAIN RELIEVER/FEVER REDUCER (NSAID)



8 FL OZ (240mL) **gf** DYE-FREE NON-STAINING ALCOHOL FREE

8 FL OZ (240mL) **gf** DYE-FREE NON-STAINING ALCOHOL FREE



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts
Active ingredient (in each 5 mL = 1 teaspoonful)
Ibuprofen 100 mg (NSAID)
Purposes
Pain reliever/
fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily:
relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
 - asthma (wheezing)
 - skin redness/itching
 - blisters
 - facial swelling
 - shock
 - rash
- If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
has had stomach ulcers or bleeding problems
takes a blood thinning (anticoagulant) or steroid drug
takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
takes more or for a longer time than directed
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use
if the child has ever had an allergic reaction to any other pain reliever/fever reducer
right before or after heart surgery

Ask a doctor before use if
stomach bleeding warning applies to your child
child has a history of stomach problems, such as heartburn
child has problems or serious side effects from taking pain relievers or fever reducers
child has not been drinking fluids
child has lost a lot of fluid due to vomiting or diarrhea
child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
child has asthma
child is taking a diuretic

Ask a doctor or pharmacist before use if the child is
under a doctor's care for any serious condition
taking any other drug

Drug Facts (continued)

When using this product
give with food or milk if stomach upset occurs
the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
Stop use and ask a doctor if
child experiences any of the following signs of stomach bleeding
feels faint
vomits blood
has bloody or black stools
has stomach pain that does not get better
the child does not get any relief within first day (24 hours) of treatment
fever or pain gets worse or lasts more than 3 days
redness or swelling is present in the painful area
any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
this product does not contain directions or complete warnings for adult use
do not give more than directed
shake well before using
mL = milliliter; tsp = teaspoonful
find right dose on chart. If possible, use weight to dose; otherwise use age.
use only enclosed dosing cup. Do not use any other dosing device.
if needed, repeat dose every 6-8 hours
do not use more than 4 times a day
replace original bottle cap to maintain child resistance
wash dosage cup after each use

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

**or as directed by a doctor
Other information
each 5 mL (1 tsp) contains: sodium 2 mg
do not use if printed neckband is broken or missing
store at 20-25°C (68-77°F)
do not freeze
see bottom panel for lot number and expiration date

Inactive ingredients: anhydrous citric acid, artificial mixed berry flavor, glycerin, high fructose corn syrup, hypromellose, polyorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions or comments? 1-800-934-1204

***This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin® Dye-Free.



LOT NO.

EXP.

1 68534 B3 C7

BERKLEY AND JENSEN IBUPROFEN CHILDRENS DYE FREE
ibuprofen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68391-685
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-685-34	1 in 1 CARTON		
1		240 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074937	01/07/2010	

Labeler - BJWC (159082692)