WALGREENS SODIUM CHLORIDE OPHTHALMIC 5 PERCENT HYPERTONICITY EYE- sodium chloride ointment Walgreen Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Walgreens Sodium Chloride Ophthalmic Ointment 5 Percent Hypertonicity Eye Ointment

Walgreens Sodium Chloride Ophthalmic
Ointment 5% Hypertonicity Eye Ointment 3.5g
NDC 0363-7500-50
Drug Facts

Active ingredient

Sodium Chloride, 50 mg (5%)

Purpose

Hypertonicity Agent

Use

• For the temporary relief of corneal edema.

Warnings

- For use in the eyes only.
- Retain outer carton for full product drug facts.

Do not use

this product except under the advice and supervision of a doctor.

When using this product

- avoid contamination, do not touch tip of container to any surface.
- replace cap after use.
- this product may cause temporary burning and irritation on being instilled into the eye.

Stop use and ask a doctor if

- you experience eye pain, changes in vision, continued redness or irritation of the eye.
- condition worsens or persists for more than 72 hours.

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

• pull down lower lid of the affected eye(s) and apply a small amount (1/4 inch) to the inside of the eyelid and apply every 3-4 hours, or as directed by a doctor.

Other information

- Store at room temperature 15°- 30°C (59°- 86°F).
- Do not freeze.
- Keep tightly closed.
- See crimp of tube or box for lot number and expiration date.

Inactive ingredients

lanolin alcohol, mineral oil, water for injection and white petrolatum.

Questions or comments?

•1-800-925-4733

PRINCIPAL DISPLAY PANEL

NDC 0363-7500-50

Sodium Chloride

Ophthalmic

Ointment, 5%

Hypertonicity

Eye Ointment

NET WT 0.125 OZ (3.5 g)

Void Aqueous

Walgreens

DO NOT USE IF BOTTOM RIDGE OF TUBE IS EXPOSED, OR IF CAP OR TUBE IS DAMAGED

Drug Facts

Active ingredient

Purpose

Sodium chloride, (5%)

... Hypertonicity agent

Use

. For the temporary relief of corneal edema.

Warnings

- . For use in the eyes only.
- Retain outer carton for full product drug facts.

Do not use this product except under the advice and supervision of a doctor.

When using this product

- Avoid contamination, do not touch tip of container to any surface.
- . Replace cap after use.
- This product may cause temporary burning and irritation on being instilled into the eye.

Stop use and ask a doctor if

- You experience eye pain, changes in vision, continued redness or irritation of the eye.
- Condition worsens or persists for more than 72 hours.

Keep this and all drugs out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

 Pull down lower lid of the affected eye(s) and apply a small amount (1/4 inch) to the inside of the eyelid and apply every 3-4 hours, or as directed by a doctor.

Drug Facts (continued)

Other information

- Store at room temperature 15°-30°C (59°-86°F).
- . Do not freeze.
- · Keep tightly closed.
- See crimp of tube or box for lot number and expiration date.

Inactive ingredients

lanolin alcohol, mineral oil, purified water and white petrolatum.

Questions or comments?

1-800-925-4733



Wagreens Phermads t Recommended Wagreens Phermads t Survey "This product is not manufactured or distributed by Bausch & Lomb, owner of the registered trademark Muro 128° Hyperionicity Ophthalmic Ontment, 5%.

Walgreens

Compare to Muro 128® Hypertonicity Ophthalmic Ointment, 5% active ingredient#

NDC 0363-7500-50

Sodium Chloride Ophthalmic Ointment, 5% Hypertonicity Eye Ointment

- For temporary relief of corneal edema
- Sterile

NET WT 0.125 OZ (3.5 g)

ACTUAL SIZE



Void Aqueous

ORG0220-F Ini0120 R0120



WALGREENS SODIUM CHLORIDE OPHTHALMIC 5 PERCENT HYPERTONICITY EYE

sodium chloride ointment

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0363-7500

Route of Administration OPHTHALMIC

Active Ingredient/Active Moiety

LANOLIN ALCOHOLS (UNII: 884C3FA9HE)

Ingredient Name

Basis of Strength

SODIUM CHLORIDE (UNII: 451W47IQ8X) (CHLORIDE ION - UNII:Q32ZN48698)

SODIUM CHLORIDE

50 mg in 1 g

Inactive Ingredients

Ingredient Name
Strength

MINERAL OIL (UNII: T5L8T28FGP)

WATER (UNII: 059QF0K00R)

PETROLATUM (UNII: 4T6H12BN9U)

Item Code Package Description Marketing Start Date 1 NDC:0363-7500- 1 in 1 CARTON 04/01/2019 3.5 g in 1 TUBE; Type 0: Not a Combination Product

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part349	04/01/2019	

Labeler - Walgreen Company (008965063)

Revised: 3/2023 Walgreen Company