

REGULAR STRENGTH ACETAMINOPHEN- acetaminophen tablet
CARDINAL HEALTH

340R CARDINAL HEALTH 70000 0092 Acetaminophen Tablets

DRUG FACTS

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- the common cold
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

adults and children 12 years and over

- take 2 tablets (650 mg) every 4 to 6 hours while symptoms last
- do not take more than 10 tablets (3,250 mg) in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children 6 years to under 12 years

- take 1 tablet (325 mg) every 4 to 6 hours while symptoms last
- do not take more than 5 tablets (1,625 mg) in 24 hours
- do not use for more than 5 days unless directed by a doctor

children under 6 years

ask a doctor

OTHER INFORMATION

Other information

- SODIUM FREE
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain this ingredient

Questions or comments? Call 1-877-290-4008

Exp Date:
Lot No.:



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet) Purpose

Acetaminophen 325 mg Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - the common cold
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
 - hives
 - itching
 - blistering
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Drug Facts (continued)

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
 - pain gets worse or lasts more than 5 days in children under 12 years
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed

- adults and children 12 years and over
 - take 2 tablets (650 mg) every 4 to 6 hours while symptoms last
 - do not take more than 10 tablets (3,250 mg) in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

Drug Facts (continued)

children 6 years to under 12 years

- take 1 tablet (325 mg) every 4 to 6 hours while symptoms last
- do not take more than 5 tablets (1,625 mg) in 24 hours
- do not use for more than 5 days unless directed by a doctor

children under 6 years

ask a doctor

Other information

- SODIUM FREE**
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients povidone, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain this ingredient

Questions or comments? Call 1-877-290-4008

*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Regular Strength. 340R 0321

©2021 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners. DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43017 www.myleader.com 1-800-200-6313

100% Money Back Guarantee
Return to place of purchase if not satisfied.

LEADER²

NDC 70000-0092-1

Regular Strength Acetaminophen
Tablets, 325 mg | Pain Reliever / Fever Reducer

COMPARE TO
TYLENOL[®]
REGULAR STRENGTH
active ingredient*

100% Money Back Guarantee

Actual Size

100 TABLETS

100 TABLETS

Regular Strength
Tablets, 325 mg
Pain Reliever / Fever Reducer

LEADER²



RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION
Active ingredient (in each tablet) Acetaminophen 325 mg

Purpose Pain reliever/fever reducer
Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ the common cold ■ backache ■ minor pain of arthritis ■ toothache ■ muscular aches ■ premenstrual and menstrual cramps ■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if: ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours, which is the maximum daily amount ■ taken with other drugs containing acetaminophen ■ alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.
Do not use ■ with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed adults and children 12 years and over: ■ take 2 tablets (650 mg) every 4 to 6 hours while symptoms last. ■ do not take more than 10 tablets (3,250 mg) in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor children 6 years to under 12 years: ■ take 1 tablet (325 mg) every 4 to 6 hours while symptoms last. ■ do not take more than 5 tablets (1,625 mg) in 24 hours ■ do not use for more than 5 days unless directed by a doctor children under 6 years: ■ ask a doctor

Other information ■ TAMPER EVIDENCE: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING ■ SODIUM FREE

■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package
Questions or comments? Call 1-877-290-4008
 CN 5706767 REV. 4/21
 100% Money Back Guarantee
 Return to place of purchase if not satisfied.
 Lot No.:
 Exp. Date:

DIST. BY: CARL DUBOIN, PH. 45017 ©2017 Cardinal Health
 340R 0321

REGULAR STRENGTH ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0092
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL;340
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0092-1	1 in 1 CARTON	05/05/2021	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/05/2021	

Labeler - CARDINAL HEALTH (063997360)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(70000-0092)

Revised: 1/2024

CARDINAL HEALTH