

MOTION SICKNESS- dimenhydrinate tablet
Strategic Sourcing Services LLC

Foster and Thrive 44-198

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- alcohol, sedatives, and tranquilizers may increase drowsiness
- marked drowsiness may occur
- use caution when driving a motor vehicle or operating machinery
- avoid alcoholic beverages

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

adults and children 12 years and over	1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years	½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years	½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Other information

- **each tablet contains:** calcium 35 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°C-30°C (59°F-86°F)
- protect from moisture
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

Principal Display Panel

NDC 70677-1087-1

COMPARE TO DRAMAMINE®
ORIGINAL FORMULA ACTIVE INGREDIENT*

Foster & Thrive™

Motion Sickness

Dimenhydrinate 50 mg
ANTIEMETIC

Prevents:

• Nausea • Vomiting • Dizziness
for Children & Adults

ACTUAL SIZE

12 TABLETS

*This product is not manufactured or distributed by Medtech Products Inc.,
owner of the registered trademark Dramamine® Original Formula.
50844 ORG051819802 Rev. 03/23

Distributed by: McKesson Corp., via Strategic Sourcing
Services LLC., Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com

©2023 McKesson Corporation

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**



12 TABLETS

Prevents:
• Nausea • Vomiting • Dizziness
for Children & Adults

©2023 McKesson Corporation
Distributed by: McKesson Corp., via Strategic Sourcing
Services LLC, Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com
B-1243-198-02
OR6051819802

This product is not manufactured or distributed by Mediatech Products Inc.,
owner of the registered trademark Dramamine® Original Formula.
50844 OR6051819802 Rev. 03/23

Drug Facts

Active ingredient (in each tablet) Purpose
Dimenhydrinate 50 mg. Antiemetic

Uses for prevention and treatment of these symptoms associated with motion sickness:
nausea ■ vomiting ■ dizziness

Warnings
Do not use for children under 2 years of age unless directed by a doctor.
■ Ask a doctor before use if you have a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ difficulty in urination due to enlargement of the prostate gland
■ Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.
■ When using this product alcohol, sedatives, and tranquilizers may increase drowsiness ■ marked drowsiness may occur ■ use caution when driving a motor vehicle or operating machinery ■ avoid alcoholic beverages
■ If pregnant or breast-feeding, ask a health professional before use.
■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions
■ to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity
adults and children 1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
children 6 to under 12 years ½ to 1 tablet every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
children 2 to under 6 years ½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Other information
■ each tablet contains: calcium 35 mg
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BUSTERS IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture ■ see end flap for expiration date and lot number


Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?
Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BUSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

No Print/No Varnish Area
Lot & Exp. date



0 10939 95623 1

Foster and Thrive 44-198

MOTION SICKNESS

dimenhydrinate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1087
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M, 8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	44;198
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1087-1	2 in 1 CARTON	05/04/2023	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	05/04/2023	

Labeler - Strategic Sourcing Services LLC (116956644)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(70677-1087)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(70677-1087) , pack(70677-1087)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(70677-1087)