

## **NASAL DECONGESTANT- pseudoephedrine hcl tablet, film coated**

**CVS Pharmacy**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **CVS Pharmacy, Inc. Nasal Decongestant Drug Facts**

#### **Active ingredient (in each tablet)**

Pseudoephedrine HCl 30 mg

#### **Purpose**

Nasal decongestant

#### **Uses**

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

#### **Warnings**

##### **Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

##### **Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

##### **When using this product**

do not exceed recommended dosage

##### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

##### **If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 2 tablets every 4 to 6 hours</li><li>• do not take more than 8 tablets in 24 hours</li></ul>
children ages 6 to 11 years	<ul style="list-style-type: none"><li>• take 1 tablet every 4 to 6 hours</li><li>• do not take more than 4 tablets in 24 hours</li></ul>
children under 6 years	do not use this product in children under 6 years of age

**Other information**

- **each tablet contains:** calcium 20 mg
- store at 20°-25°C (68°-77°F)
- do not use if blister unit is broken or torn

**Inactive ingredients**

carnauba wax, dibasic calcium phosphate dihydrate, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, silicon dioxide, titanium dioxide

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

Compare to the active ingredient in Sudafed® Congestion

Non-Drowsy

Nasal Decongestant

PSEUDOEPHEDRINE HYDROCHLORIDE

Nasal decongestant

MAXIMUM STRENGTH

Relieves:

Nasal & sinus congestion

Sinus pressure

24 TABLETS 30 mg EACH

Actual Size

 **CVS**Health.

Non-Drowsy

# Nasal Decongestant

PSEUDOEPHEDRINE HYDROCHLORIDE/Nasal decongestant

 **CVS**Health™

Compare to the active ingredient  
in Sudafed® Congestion\*

NDC 59779-432-62

Non-Drowsy

# Nasal Decongestant

PSEUDOEPHEDRINE HYDROCHLORIDE

Nasal decongestant

**MAXIMUM STRENGTH**


Relieves:

- Nasal & sinus congestion
- Sinus pressure

**24 TABLETS 30 mg EACH**



Actual Size



43252 17 014

Important: Read all product information before using. Keep this box for important information.

OPEN OTHER END

**Drug Facts**

**Active ingredient (in each tablet) Purpose**  
Pseudoephedrine HCl 30 mg.....Nasal decongestant

**Uses**

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

**Warnings**

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Drug Facts (continued)**

**Directions**

adults and children 12 years and over	<ul style="list-style-type: none"> <li>take 2 tablets every 4 to 6 hours</li> <li>do not take more than 8 tablets in 24 hours</li> </ul>
children ages 6 to 11 years	<ul style="list-style-type: none"> <li>take 1 tablet every 4 to 6 hours</li> <li>do not take more than 4 tablets in 24 hours</li> </ul>
children under 6 years	do not use this product in children under 6 years of age

**Other information**

- each tablet contains: calcium 20 mg
- store at 20°-25°C (68°-77°F)
- do not use if blister unit is broken or torn

**Inactive ingredients**

carnauba wax, dibasic calcium phosphate dihydrate, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, silicon dioxide, titanium dioxide

**Questions or comments? 1-800-719-9260**

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Sudafed® Congestion.

**DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN**

CONVENIENT RECLOSING TAB

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2017 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS  
V-14017



#477074



**NASAL DECONGESTANT**

pseudoephedrine hcl tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:59779-432
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

**Inactive Ingredients**

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	

### Product Characteristics

<b>Color</b>	RED	<b>Score</b>	no score
<b>Shape</b>	ROUND (convex)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	L432
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-432-62	24 in 1 CARTON	07/15/1987	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:59779-432-67	48 in 1 CARTON	08/29/1997	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:59779-432-80	96 in 1 CARTON	07/28/1997	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/15/1987	

**Labeler** - CVS Pharmacy (062312574)