

CHILDRENS ROBITUSSIN COUGH AND COLD LONG-ACTING- chlorpheniramine maleate, dextrometrorphan hbr liquid
Haleon US Holdings LLC

Drug Facts

Active ingredients (in each 10 ml)

Chlorpheniramine maleate, USP 2 mg

Dextromethorphan HBr, USP 15 mg

Purposes

Antihistamine

Cough suppressant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- a cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- ml = milliliter
- do not take more than 4 doses in any 24-hour period

Age	Dose
Children under 6 years	do not use
children 6 to under 12 years	10 ml every 6 hours
adults and children 12 years and older	20 ml every 6 hours

Other information

- **each 10 ml contains:** sodium 6 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, artificial & natural flavors, FD&C red no. 40, glycerin, lactic acid, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

Call weekdays from 9 AM to 5 PM EST at **1-800-245-1040**

Additional Information

Packaged with Tamper-Evident bottle cap. Do Not Use if breakable ring is separated or missing.

Children's Robitussin liquid is specially formulated to provide **soothing action, control your child's cough** plus relieve other cold symptoms.

Should be 18 or older to purchase

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Use dosage cup included

Distributed by: GSK Consumer Healthcare,

Warren, NJ 07059

For most recent product information,

visit www.robitussin.com

Trademarks owned or licensed by GSK.

©2022 GSK or licensor

Made in Canada

PRINCIPAL DISPLAY PANEL

Children's

Robitussin

AGES 6 & OVER

Nighttime Cough

DM

Long-Acting

CHLORPHENIRAMINE MALEATE (Antihistamine)

DEXTROMETHORPHAN HBr (Cough Suppressant)

Relieves:

- **Cough Up to 8 Hours**
- **Runny Nose**

Alcohol-Free

fruit

punch

flavor

4 FL OZ

(118 ml)

PAA172103 Front Carton

Children's

Robitussin

AGES 6 & OVER

Nighttime
Cough



Long-Acting



CHLORPHENIRAMINE MALEATE
(Antihistamine)
DEXTROMETHORPHAN HBr (Cough Suppressant)

Relieves:

- ✓ Cough Up to **8 Hours**
- ✓ Runny Nose

Alcohol-Free



fruit
punch
flavor

4 FL OZ
(118 mL)

CHILDRENS ROBITUSSIN COUGH AND COLD LONG-ACTING

chlorpheniramine maleate, dextrometorphan hbr liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0031-8693
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
LACTIC ACID, UNSPECIFIED FORM (UNII: 33X04XA5AT)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	red (red liquid)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0031-8693-12	1 in 1 CARTON	01/01/2004	
1		118 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/01/2004	

Revised: 3/2024

Haleon US Holdings LLC