### IBUPROFEN- ibuprofen tablet Contract Pharmacy Services-PA

-----

### **Ibuprofen 200 mg Tablets**

#### **DRUG FACTS**

### **ACTIVE INGREDIENT (IN EACH TABLET)**

Ibuprofen 200 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

#### **PURPOSE**

Pain reliever/fever reducer

#### **USES**

- temporarily relieves minor aches and pains due to:
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

### WARNINGS

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### Ask a doctor before use if you are

- taking aspirin to prevent heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drugs

## When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

## If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222

#### **DIRECTIONS**

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

adults and children 12 years and older	take 1 tablet every 4 to 6 hours while symptoms persist	
	if pain or fever does not respond to 1 tablet, 2 tablets may be used do not exceed 6 tablets in 24 hours,	
	unless directed by a doctor	
children under 12	ask a doctor	
years		

#### OTHER INFORMATION

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid high humidity and excessive heat above 40°C (104°F)

#### **INACTIVE INGREDIENTS**

colloidal silicon dioxide, corn starch, croscarmellose sodium, hypromellose, iron oxides, microcrystalline cellulose, polyethylene glycol, polysorbate 80, stearic acid, titanium dioxide

### **QUESTIONS OR COMMENTS?**


Distributed by: Major Pharmaceuticals, 31778 Enterprise Drive, Livonia, MI 48150, U.S.A.

call toll free **734-743-6000** 

Repackaged by:

Contract Pharmacy Services-PA 125 Titus Ave Suite 200 Warrington, PA 18976 USA Original--08/2010--NJW

**Package Label** 



Blister of 30 Tablets

# **IBUPROFEN**

ibuprofen tablet

Dro	duct	Info	rmatio	1
ГІО			ııılatıvı	

Product Type HUMAN OTC DRUG Item Code (Source) NDC:67046-303(NDC:0904-7915)

Route of Administration ORAL

# **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPRO FEN (UNII: WK2XYI10 QM) (IBUPRO FEN - UNII: WK2XYI10 QM)	IBUPROFEN	200 mg

Product	Characteristics

Shape ROUND Size 10 mm	
Flavor Imprint Code 12	
Contains	

Packaging							
#	Item Code	Package Description	Marketing Start Date		M	Marketing End Date	
1 1	NDC:67046-303-30	30 in 1 BLISTER PACK					
Marketing Information							
M	arketing Category	Application Number or Monogra	ph Citation	Marketing Start	Date	Marketing End Date	
AN	DA AN	NDA072096		08/02/2010			

**Labeler -** Contract Pharmacy Services-PA (945429777)

Revised: 7/2010 Contract Pharmacy Services-PA