

FAMOTIDINE- famotidine tablet
BluePoint Laboratories

FAMOTIDINE tablet, film coated

Drug Facts

Active Ingredient in (Each Tablet)

Famotidine USP 10 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a Doctor Before Use if You Have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a Doctor or Pharmacist Before Use if You Are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop Use and ask a Doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If Pregnant or Breast-Feeding

ask a health professional before use.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away

Directions

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other Information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F)
- protect from moisture

Inactive Ingredients

carnauba wax, corn starch, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, red iron oxide, sodium starch glycolate, talc and titanium dioxide.

Questions or Comments?

call **1-855-274-4122**

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables

- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Do not use if carton is open or if printed foil seal under bottle cap is open or torn.

Manufactured by:

Aurobindo Pharma Limited

Unit-VII (SEZ), Mahabubnagar (Dt)-509302, India .

For BluePoint Laboratories

Rev: 02/2023

Package Label - Principal Display Panel

Aurobindo Pharma Limited. NDC: 68001-494-04

Famotidine Tablets USP 10mg 30s Label

NDC 68001-494-04
ORIGINAL STRENGTH
Famotidine
 Tablets USP

10 mg

Acid Reducer
Just One Tablet!
 Prevents & Relieves Heartburn
 Due to Acid Indigestion

30 Tablets

Do not use if printed foil seal on bottle is broken or missing.

Active Ingredient (in each tablet)
 Famotidine USP 10 mg.....Acid reducer

Purpose
 Famotidine USP 10 mg.....Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. **Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers.

Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with **lightheadness, sweating, or dizziness** ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadness. ■ frequent **chest pain** ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ kidney disease. **Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs. **Stop use and ask a doctor if** ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. **Directions** ■ adults and children 12 years and over: ■ to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew. ■ to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn ■ do not use more than 2 tablets in 24 hours ■ children under 12 years: ask a doctor. **Other information** ■ read the directions and warnings before use ■ keep the carton (68° to 77° F) ■ protect from moisture

Inactive ingredients See carton for complete list of inactive ingredients

Questions or comments? call 1-855-274-4122

Manufactured by:
Aurobindo Pharma Limited
 Unit-VII (SEZ), Mahabubnagar (Dt)-509302, India.
 For BluePoint Laboratories
 M.L.No.: 22/IN/AP/2009/FR Made in India

Rev: 02/2023

P14-32173 LM-5090

Aurobindo Pharma Limited. NDC: 68001-494-04

Famotidine Tablets USP 10mg 30s Carton

98805



NDC 68001-494-04

ORIGINAL STRENGTH

Famotidine
Tablets USP 10 mg
Acid Reducer
Just One Tablet!
Prevents & Relieves Heartburn
Due to Acid Indigestion

30 Tablets

Drug Facts (continued)**Directions**

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **15 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
 - children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F)
- protect from moisture

Inactive ingredients

carnauba wax, corn starch, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, red iron oxide, sodium starch glycolate, talc and titanium dioxide.

Questions or comments?

call 1-855-274-4122

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc. McNeil Consumer Healthcare Division, distributor of Original Strength Peppid® AC.

NDC 68001-494-04

**Compare to the Active Ingredient of
Original Strength Peppid® AC*

ORIGINAL STRENGTH

Famotidine
Tablets USP
10 mg
Acid Reducer
Just One Tablet!
Prevents & Relieves Heartburn
Due to Acid Indigestion

Actual Size

30 Tablets

Manufactured by: **Aurobindo Pharma Limited**
 Unit-VII (SEZ), Mahabubnagar (Dt)-509302, India.
 For BluePoint Laboratories

M.L.No.: 22/MN/AP/2009/F/R

Rev: 02/2023



N 3168001149404115

P1050886 LM-5089

Lot:

EXP.:

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Do not use if carton is open or if printed foil seal under bottle cap is open or torn.

Drug Facts

Active ingredient (in each tablet)	Purpose
Famotidine USP 10 mg.....	Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or

Drug Facts (continued)

black stools. These may be signs of a serious condition. See your doctor.

- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.

- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

FAMOTIDINE

famotidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68001-494
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)	
TALC (UNII: 7SEV7J4R1U)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND (biconvex)	Size	5mm
Flavor		Imprint Code	CC;58
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68001-494-04	1 in 1 CARTON	04/07/2021	
1		30 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:68001-494-06	1 in 1 CARTON	06/17/2021	
2		60 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206531	04/07/2021	

Labeler - BluePoint Laboratories (985523874)

Establishment

Name	Address	ID/FEI	Business Operations
Aurobindo Pharma Limited		650381903	manufacture(68001-494) , analysis(68001-494)