

**SUBVENTE: lamotrigine tablet**  
**REMEDYREPACK INC.**

**HIGHLIGHTS OF PRESCRIBING INFORMATION**  
These highlights do not include all the information needed to use SUBVENTE safely and effectively. See full prescribing information for SUBVENTE.

**SUBVENTE (lamotrigine) tablets USP, for oral use**

Initial U.S. Approval: 1994

**WARNING: SERIOUS SKIN RASHES**

- See full prescribing information for complete boxed warning.
- Cases of life-threatening serious rashes, including Stevens-Johnson syndrome and toxic epidermal necrolysis, and/or rash-related death have been caused by lamotrigine. The rate of serious rash is greater in pediatric patients than in adults. Additional factors that may increase the risk of rash include:
    - coadministration with valproate.
  - Exceeding recommended initial dose of SUBVENTE.
  - Exceeding recommended dose escalation for SUBVENTE. (5.1)
  - Rash-related deaths are also caused by SUBVENTE; however, it is not possible to predict which rashes will prove to be serious or life-threatening. Lamotrigine should be discontinued at the first sign of rash, unless the rash is clearly not drug related. (5.1)

**RECENT MAJOR CHANGES**

Warnings and Precautions: Lymphocytopenia 8/2019

Lymphohistiocytosis (5.2)

**INDICATIONS AND USAGE**

SUBVENTE is indicated for:

- **Adjunctive therapy in patients aged 2 years and older.**

- partial-onset seizures.
- primary generalized tonic-clonic seizures.
- generalized seizures of Lennox-Gastaut syndrome. (1.1)

**Warnings: monotherapy in patients aged 16 years and older.** Conversion to monotherapy in patients with partial-onset seizures who are receiving treatment with carbamazepine, phenytoin, phenobarbital, primidone, or valproate as the single AED. (1.1)

**Usage: Maintenance treatment of bipolar I disorder to delay the time to occurrence of mood episodes in patients treated for acute mood episodes with standard therapy. (1.2)**

**Limitations of Use:** Treatment of acute manic or mixed episodes is not recommended. Effectiveness of SUBVENTE in the acute treatment of mood episodes has not been established.

**DOSE AND ADMINISTRATION**

- Dosing is based on concomitant medication, indication, and patient age. (2.1, 2.2, 2.3, 2.4)

- To avoid an increased risk of rash, the recommended initial dose and subsequent dose escalations should not be exceeded. SUBVENTE starter kits are available for the first 5 weeks of treatment. (2.1, 6)

- Do not restart SUBVENTE in patients who discontinued due to rash unless the potential benefits clearly outweigh the risks. (2.1, 5.1)

- Adjustment to maintenance doses will be necessary in most patients starting or stopping estrogen-containing oral contraceptives. (2.1, 5.8)

- Discontinuation: Taper over a period of at least 2 weeks (approximately 50% dose reduction per week). (2.1, 5.9)

- **Follow-up:**
  - Adjunctive therapy—See Table 1 for patients older than 12 years and Tables 2 and 3 for patients aged 2 to 12 years. (2.3)
  - Conversion to monotherapy—See Table 4. (2.3)

**Usual dosing:** See Tables 5 and 6. (2.4)

**DOSE FORMS AND STRENGTHS**

- Tablets: 25 mg, 100 mg, 150 mg, and 200 mg; oral solution: 1.1, 1.1, 10

**CONTRAINDICATIONS**

Hypersensitivity to the drug or its ingredients. (Based on Warning, 4)

**WARNINGS AND PRECAUTIONS**

- Life-threatening serious rash and/or rash-related death. Discontinue at the first sign of rash, unless the rash is clearly not drug-related. (Based on Warning, 5.1)

- Hematophagocytic lymphohistiocytosis. Consider this diagnosis and evaluate patients immediately if they develop signs or symptoms of systemic inflammation. Discontinue SUBVENTE if an alternative etiology is not established. (5.2)

- Fatal life-threatening hypersensitivity reactions. Multorgan hypersensitivity reactions, also known as drug reaction with eosinophilia and systemic symptoms, may be fatal or life-threatening. Early signs may include rash, fever, and lymphadenopathy. These reactions may be associated with other organ involvement, such as hepatic, hepatic failure, blood dyscrasias, or acute myocardial infarction. SUBVENTE should be discontinued if alternate etiology for this reaction is not found. (5.3)

- Blood dyscrasias (e.g., neutropenia, thrombocytopenia, pancytopenia) may occur, either with or without an associated hypersensitivity syndrome. Monitor for signs of anemia, unexplained infections, or bleeding. (5.4)

- Suicidal behavior and ideation: Monitor for suicidal thoughts or behaviors. (5.5)

- Aseptic meningitis: Monitor for signs of meningitis. (5.6)

- Medication errors due to product name confusion: Strongly advise patients to visually inspect tablets to verify the recorded drug content. (5.7, 6, 17)

**ADVERSE REACTIONS**

**Follow-up:** Most common adverse reactions (incidence ≥ 10%) in adults were diarrhea, headache, dizziness, ataxia, nausea, vomiting, flu-like symptoms, rhinitis, pharyngitis, and rash. Additional adverse reactions (incidence ≥ 10%) reported in children included vomiting, infection, fever, accidental injury, diarrhea, abdominal pain, and tremor. (6.1)

**Usual dosing:** Most common adverse reactions (incidence ≥ 10%) in adults were nausea, insomnia, constipation, back pain, fatigue, rash, rhinitis, abdominal pain, and vertigo/dizziness. (6.1)

**To report SUSPECTED ADVERSE REACTIONS,** contact OWP Pharmaceuticals Inc. at 1-800-373-6729 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

**DRUG INTERACTIONS**

- Valproate increases lamotrigine concentrations more than 2-fold. (7, 12.3)

- Carbamazepine, phenytoin, phenobarbital, primidone, and rifampin decrease lamotrigine concentrations by approximately 60%. (7, 12.3)

- Estrogen-containing oral contraceptives decrease lamotrigine concentrations by approximately 50%. (7, 12.3)

- Protein inhibitors (bupropion) increase and azarone/bupropion decrease lamotrigine exposure by approximately 50% and 20%, respectively. (7, 12.3)

- Coadministration with organic cationic transporter 2 substrates with narrow therapeutic index is not recommended. (7, 12.3)

**USE IN SPECIFIC POPULATIONS**

- **Pregnancy:** Based on animal data may cause fetal harm. (8.1)

- **Hepatic impairment:** Dose adjustments required in patients with moderate and severe liver impairment. (2.1, 6)

- **Renal impairment:** Reduced maintenance doses may be effective for patients with significant renal impairment. (2.1, 8.7)

Revised: 5/2020

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

1.1 Epilepsy

1.2 Bipolar Disorder

**2 DOSAGE AND ADMINISTRATION**

2.1 General Dosing Considerations

2.2 Epilepsy-Adjunctive Therapy

2.3 Epilepsy-Conversion from Adjunctive Therapy to Monotherapy

2.4 Bipolar Disorder

**3 DOSAGE FORMS AND STRENGTHS**

3.1 Tablets

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

5.1 Serious Skin Rashes [see Boxed Warning]

5.2 Hematophagocytic Lymphohistiocytosis

5.3 Multorgan Hypersensitivity Reactions and Organ Failure

5.4 Blood Dyscrasias

5.5 Suicidal Behavior and Ideation

5.6 Aseptic Meningitis

5.7 Potential Medication Errors

5.8 Concomitant Use with Oral Contraceptives

5.9 Withdrawal Seizures

5.10 Status Epilepticus

5.11 Sudden Unexplained Death in Epilepsy (SUDEP)

5.12 Addition of SUBVENTE to a Multidrug Regimen that Includes Valproate

5.13 Binding in the Eye and Other Mucous-Membrane Tissues

5.14 Laboratory Tests

**6 ADVERSE REACTIONS**

6.1 Clinical Trial Experience

6.2 Other Adverse Reactions Observed in All Clinical Trials

6.3 Postmarketing Experience

**7 DRUG INTERACTIONS**

**8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.6 Hepatic Impairment

8.7 Renal Impairment

**10 OVERDOSAGE**

10.1 Human Overdose Experience

10.2 Management of Overdose

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14 CLINICAL STUDIES**

14.1 Epilepsy

14.2 Bipolar Disorder

**16 HOW SUPPLIED/STORAGE AND HANDLING**

\* Sections of subsection omitted from the full prescribing information are not listed.

**FULL PRESCRIBING INFORMATION**

**WARNING: SERIOUS SKIN RASHES**

SUBVENTE™ can cause serious rashes requiring hospitalization and discontinuation of treatment. The incidence of these rashes, which have included Stevens-Johnson syndrome, is approximately 0.3% to 0.8% in pediatric patients (aged 2 to 17 years) and 0.08% to 0.3% in adults receiving SUBVENTE. One rash-related death was reported in a prospectively followed cohort of 1,183 pediatric patients (aged 2 to 16 years) with epilepsy taking SUBVENTE as adjunctive therapy. In worldwide postmarketing experience, rare cases of toxic epidermal necrolysis and/or rash-related death have been reported in adult and pediatric patients, but their numbers are too few to permit a precise estimate of the rate.

Other than age, there are as yet no factors identified that are known to predict the risk of occurrence or the severity of rash caused by SUBVENTE. There are suggestions, yet to be proven, that the risk of rash may also be increased by (1) coadministration of SUBVENTE with valproate (includes valproic acid and divalproex sodium), (2) exceeding the recommended initial dose of SUBVENTE, or (3) exceeding the recommended dose escalation for SUBVENTE. However, cases have occurred in the absence of these factors.

Nearly all cases of life-threatening rashes caused by SUBVENTE have occurred within 2 to 8 weeks of treatment initiation. However, isolated cases have occurred after prolonged treatment (e.g., 6 months). Accordingly, duration of therapy cannot be relied upon as a means to predict the potential risk heralded by the first appearance of a rash.

Although benign rashes are also caused by SUBVENTE, it is not possible to predict reliably which rashes will prove to be serious or life-threatening. Accordingly, SUBVENTE should ordinarily be discontinued at the first sign of rash, unless the rash is clearly not drug related. Discontinuation of treatment may not prevent a rash from becoming life-threatening or permanently disabling or disfiguring [see Warnings and Precautions (5.1)].

**1 INDICATIONS AND USAGE**

1.1 Epilepsy

**Adjunctive Therapy**

SUBVENTE is indicated as adjunctive therapy for the following seizure types in patients aged 2 years and older:

- partial-onset seizures.
- primary generalized tonic-clonic (PGTC) seizures.
- generalized seizures of Lennox-Gastaut syndrome.

**Monotherapy**

SUBVENTE is indicated for conversion to monotherapy in adults (aged 16 years and older) with partial-onset seizures who are receiving treatment with carbamazepine, phenytoin, phenobarbital,

primidone, or valproate as the single antiepileptic drug (AED).

Safety and effectiveness of SUBVENTE have not been established (1) as initial monotherapy; (2) for conversion to monotherapy from AEDs other than carbamazepine, phenytoin, phenobarbital, primidone, or valproate; or (3) for simultaneous conversion to monotherapy from 2 or more concomitant AEDs.

## 1.2 Bipolar Disorder

SUBVENTE is indicated for the maintenance treatment of bipolar I disorder to delay the time to occurrence of mood episodes (depression, mania, hypomania, mixed episodes) in patients treated for acute mood episodes with standard therapy [see Clinical Studies (14.1)].

### Limitation of Use

Treatment of acute manic or mixed episodes is not recommended. Effectiveness of SUBVENTE in the acute treatment of mood episodes has not been established.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 General Dosing Considerations

#### Risk

There are suggestions, yet to be proven, that the risk of severe, potentially life-threatening rash may be increased by (1) coadministration of SUBVENTE with valproate, (2) exceeding the recommended initial dose of SUBVENTE, or (3) exceeding the recommended dose escalation for SUBVENTE. However, cases have occurred in the absence of these factors [see Black Box Warning]. Therefore, it is important that the dosing recommendations be followed closely.

The risk of non-serious rash may be increased when the recommended initial dose and/or the rate of dose escalation for SUBVENTE is exceeded and in patients with history of allergy or rash to other AEDs.

SUBVENTE Starter Kits provide SUBVENTE at doses consistent with the recommended titration schedule for the first 3 weeks of treatment, based upon concomitant medications for patients with epilepsy (older than 12 years) and bipolar I disorder (adults) and are intended to help reduce the potential for rash. The use of SUBVENTE Starter Kits is recommended for appropriate patients who are starting or restarting SUBVENTE [see How Supplied/Storage and Handling (16)].

It is recommended that SUBVENTE not be restarted in patients who discontinued due to rash associated with prior treatment with lamotrigine unless the potential benefits clearly outweigh the risks. If the decision is made to restart a patient who has discontinued SUBVENTE, the need to restart with the initial dosing recommendations should be assessed. The greater the interval of time since the previous dose, the greater consideration should be given to restarting with the initial dosing recommendation. If a patient has discontinued lamotrigine for a period of more than 3 half-lives, it is recommended that initial dosing recommendations and guidelines be followed. The half-life of lamotrigine is affected by other concomitant medications [see Clinical Pharmacology (12.3)].

#### SUBVENTE Added to Drugs Known to Inhibit Glucuronidation

Because lamotrigine is metabolized predominantly by glucuronic acid conjugation, drugs that are known to induce or inhibit glucuronidation may affect the apparent clearance of lamotrigine. Drugs that induce glucuronidation include carbamazepine, phenytoin, phenobarbital, primidone, rifampin, estrogen-containing oral contraceptives, and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir. Valproate inhibits glucuronidation. For dosing considerations for SUBVENTE in patients on estrogen-containing contraceptives and atazanavir/ritonavir, see below and Table 13. For dosing considerations for SUBVENTE in patients on other drugs known to induce or inhibit glucuronidation, see Tables 1, 2, 5-6, and 13.

#### Target Plasma Levels for Patients with Epilepsy or Bipolar Disorder

A therapeutic plasma concentration range has not been established for lamotrigine. Dosing of SUBVENTE should be based on therapeutic response [see Clinical Pharmacology (12.3)].

#### Women Taking Estrogen-Containing Oral Contraceptives

Starting SUBVENTE in Women Taking Estrogen-Containing Oral Contraceptives: Although estrogen-containing oral contraceptives have been shown to increase the clearance of lamotrigine [see Clinical Pharmacology (12.3)], no adjustments to the recommended dose-escalation guidelines for SUBVENTE should be necessary solely based on the use of estrogen-containing oral contraceptives. Therefore, dose escalation should follow the recommended guidelines for initiating adjunctive therapy with SUBVENTE based on the concomitant AED or other concomitant medications [see Tables 4, 5, and 7]. See below for adjustments to maintenance doses of SUBVENTE in women taking estrogen-containing oral contraceptives.

#### Adjustments to the Maintenance Dose of SUBVENTE in Women Taking Estrogen-Containing Oral Contraceptives

(1) Taking Estrogen-Containing Oral Contraceptives: In women not taking carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir that induce lamotrigine glucuronidation [see Drug Interactions (7), Clinical Pharmacology (12.3)], the maintenance dose of SUBVENTE will in most cases need to be increased by as much as 2-fold over the recommended target maintenance dose to maintain a consistent lamotrigine plasma level.

(2) Starting Estrogen-Containing Oral Contraceptives: In women taking a stable dose of SUBVENTE and not taking carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir that induce lamotrigine glucuronidation [see Drug Interactions (7), Clinical Pharmacology (12.3)], the maintenance dose will in most cases need to be increased by as much as 2-fold to maintain a consistent lamotrigine plasma level. The dose increases should begin at the same time that the oral contraceptive is introduced and continue, based on clinical response, no more rapidly than 50 to 100 mg/day every week. Dose increases should not exceed the recommended rate [see Tables 1 and (3)] unless lamotrigine plasma levels or clinical response support larger increases. Gradual transient increases in lamotrigine plasma levels may occur during the week of inactive hormonal preparation (pill-free week), and these increases will be greater if dose increases are made in the days before or during the week of inactive hormonal preparation. Increased lamotrigine plasma levels could result in additional adverse reactions, such as dizziness, nausea, and diplopia. If adverse reactions attributable to SUBVENTE consistently occur during the pill-free week, dose adjustments to the overall maintenance dose may be necessary. Dose adjustments limited to the pill-free week are not recommended. For women taking SUBVENTE in addition to carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir that induce lamotrigine glucuronidation [see Drug Interactions (7), Clinical Pharmacology (12.3)], no adjustment to the dose of SUBVENTE should be necessary.

(3) Stopping Estrogen-Containing Oral Contraceptives: In women not taking carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir that induce lamotrigine glucuronidation [see Drug Interactions (7), Clinical Pharmacology (12.3)], the maintenance dose of SUBVENTE will in most cases need to be decreased by as much as 50% in order to maintain a consistent lamotrigine plasma level. The decrease in dose of SUBVENTE should not exceed 25% of the total daily dose per week over a 2-week period, unless clinical response or lamotrigine plasma levels indicate otherwise [see Clinical Pharmacology (12.3)]. In women taking SUBVENTE in addition to carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir that induce lamotrigine glucuronidation [see Drug Interactions (7), Clinical Pharmacology (12.3)], no adjustment to the dose of SUBVENTE should be necessary.

#### Women and Other Hormonal Contraceptive Preparations or Hormone Replacement Therapy

The effect of other hormonal contraceptive preparations or hormone replacement therapy on the pharmacokinetics of lamotrigine has not been systematically evaluated. It has been reported that norgestrel, but not progesterone, increased the clearance of lamotrigine up to 2-fold, and the progestin-only pills had no effect on lamotrigine plasma levels. Therefore, adjustments to the dosage of SUBVENTE in the presence of progestogen alone will likely not be needed.

#### Patients Taking Atazanavir/Ritonavir

While atazanavir/ritonavir does reduce the lamotrigine plasma concentration, no adjustments to the recommended dose-escalation guidelines for SUBVENTE should be necessary solely based on the use of atazanavir/ritonavir. Dose escalation should follow the recommended guidelines for initiating adjunctive therapy with SUBVENTE based on concomitant AED or other concomitant medications (see Tables 1, 2, and 5). In patients already taking maintenance doses of SUBVENTE and not taking glucuronidation inducers, the dose of SUBVENTE may need to be increased if atazanavir/ritonavir is added, or decreased if atazanavir/ritonavir is discontinued [see Clinical Pharmacology (12.3)].

#### Patients with Hepatic Impairment

Experience in patients with hepatic impairment is limited. Based on a clinical pharmacology study in 24 subjects with mild, moderate, and severe liver impairment [see Use in Specific Populations (8.6), Clinical Pharmacology (12.3)], the following general recommendations can be made. No dosage adjustment is needed in patients with mild liver impairment. Initial, escalation, and maintenance doses should generally be reduced by approximately 25% in patients with moderate and severe liver impairment without ascites and 50% in patients with severe liver impairment with ascites. Escalation and maintenance doses may be adjusted according to clinical response.

#### Patients with Renal Impairment

Initial doses of SUBVENTE should be based on patients' concomitant medications (see Tables 1-3, and 5), reduced maintenance doses may be effective for patients with significant renal impairment [see Use in Specific Populations (8.7), Clinical Pharmacology (12.3)]. Few patients with severe renal impairment have been evaluated during chronic treatment with SUBVENTE. Because there is inadequate experience in this population, SUBVENTE should be used with caution in these patients.

#### Discontinuation Strategy

Epilepsy: For patients receiving SUBVENTE in combination with other AEDs, a re-evaluation of all AEDs in the regimen should be considered if a change in seizure control or an appearance or worsening of adverse reactions is observed. If a decision is made to discontinue therapy with SUBVENTE, a step-wise reduction of dose over at least 2 weeks (approximately 50% per week) is recommended unless safety concerns require a more rapid withdrawal [see Warnings and Precautions (5.9)].

Discontinuing carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir that induce lamotrigine glucuronidation should prolong the half-life of lamotrigine; discontinuing valproate should shorten the half-life of lamotrigine.

Bipolar Disorder: In the controlled clinical trials, there was no increase in the incidence, type, or severity of adverse reactions following abrupt termination of SUBVENTE. In the clinical development program in adults with bipolar disorder, 2 patients experienced seizures shortly after abrupt withdrawal of SUBVENTE. Discontinuation of SUBVENTE should involve a step-wise reduction of dose over at least 2 weeks (approximately 50% per week) unless safety concerns require a more rapid withdrawal [see Warnings and Precautions (5.9)].

### 2.2 Epilepsy-Adjunctive Therapy

This section provides specific dosing recommendations for patients older than 12 years and patients aged 2 to 12 years. Within each of these age-groups, specific dosing recommendations are provided depending upon concomitant AEDs or other concomitant medications (see Table 1 for patients older than 12 years and Table 2 for patients aged 2 to 12 years). A weight-based dosing guide for patients aged 2 to 12 years on concomitant valproate is provided in Table 3.

#### Patients Older than 12 Years

Recommended dosing guidelines are summarized in Table 1.

Table 1. Escalation Regimen for SUBVENTE in Patients Older than 12 Years with Epilepsy

	In Patients TAKING Valproate <sup>a</sup>	In Patients NOT TAKING Carbamazepine, Phenytoin, Phenobarbital, Primidone <sup>b</sup> or Valproate <sup>a</sup>	Patients TAKING Carbamazepine, Phenytoin, Phenobarbital, or Primidone <sup>b</sup> and NOT TAKING Valproate <sup>a</sup>
Weeks 1 and 2	25 mg every other day	25 mg every day	50 mg/day
Weeks 3 and 4	25 mg every other day	50 mg every day	100 mg/day (in 2 divided doses)
Week 5 onward to maintenance	Increase by 25 to 50 mg/day every 1 to 2 weeks	Increase by 50 mg/day every 1 to 2 weeks	Increase by 100 mg/day every 1 to 2 weeks
Usual maintenance dose	100 to 200 mg/day with valproate alone 100 to 400 mg/day with valproate and/or other drugs that induce glucuronidation <sup>c</sup> (in 1 or 2 divided doses)	225 to 375 mg/day (in 2 divided doses)	300 to 500 mg/day (in 2 divided doses)

<sup>a</sup> Valproate has been shown to inhibit glucuronidation and decrease the apparent clearance of lamotrigine [see Drug Interactions (7), Clinical Pharmacology (12.3)].

<sup>b</sup> Drugs that induce lamotrigine glucuronidation, such as carbamazepine, phenytoin, phenobarbital, primidone, rifampin, and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir. Dosing recommendations for oral contraceptives and the protease inhibitor atazanavir/ritonavir can be found in General Dosing Considerations for Dosage and Administration (2.1). Patients on rifampin and the protease inhibitor lopinavir/ritonavir should follow the same dosing/titration/maintenance regimen used with antiepileptic drugs that induce glucuronidation and increase clearance [see Dosage and Administration (2.1), Drug Interactions (7), and Clinical Pharmacology (12.3)].

#### Patients Aged 2 to 12 Years

Recommended dosing guidelines are summarized in Table 2.

Lower starting doses and slower dose escalations than those used in clinical trials are recommended because of the suggestion that the risk of rash may be decreased by lower starting doses and slower dose escalation. Therefore, maintenance doses will take longer to reach in clinical practice than in clinical trials. It may take several weeks to months to achieve an individualized maintenance dose. Maintenance doses in patients weighing less than 30 kg, regardless of age or concomitant AED, may need to be increased as much as 50%, based on clinical response.

The smallest available strength of SUBVENTE is 25 mg, and only whole tablets should be administered. If the calculated dose cannot be achieved using whole tablets, the dose should be rounded down to the nearest whole tablet [see How Supplied/Storage and Handling (16) and Medication Guide].

Table 2. Escalation Regimen for SUBVENTE in Patients Aged 2 to 12 Years with Epilepsy

	In Patients TAKING Valproate <sup>a</sup>	In Patients NOT TAKING Carbamazepine, Phenytoin, Phenobarbital, Primidone <sup>b</sup> or Valproate <sup>a</sup>	In Patients TAKING Carbamazepine, Phenytoin, Phenobarbital, or Primidone <sup>b</sup> and NOT TAKING Valproate <sup>a</sup>
Weeks 1 and 2	0.15 mg/kg/day in 1 or 2 divided doses, rounded	0.3 mg/kg/day in 1 or 2 divided doses, rounded down	0.6 mg/kg/day in 2 divided doses, rounded down to the

	down to the nearest whole tablet (see Table 3 for weight-based dosing guide)	to the nearest whole tablet	nearest whole tablet
Weeks 3 and 4	<b>0.3 mg/kg/day</b> in 1 or 2 divided doses, rounded down to the nearest whole tablet (see Table 3 for weight-based dosing guide)	<b>0.6 mg/kg/day</b> in 2 divided doses, rounded down to the nearest whole tablet	<b>1.2 mg/kg/day</b> in 2 divided doses, rounded down to the nearest whole tablet
Week 5 onward to maintenance	The dose should be increased every 1 to 2 weeks as follows: calculate 0.3 mg/kg/day, round this amount down to the nearest whole tablet, and add this amount to the previously administered daily dose.	The dose should be increased every 1 to 2 weeks as follows: calculate 0.6 mg/kg/day, round this amount down to the nearest whole tablet, and add this amount to the previously administered daily dose.	The dose should be increased every 1 to 2 weeks as follows: calculate 1.2 mg/kg/day, round this amount down to the nearest whole tablet, and add this amount to the previously administered daily dose.
Usual Maintenance Dose	<b>1 to 5 mg/kg/day</b> (maximum 200 mg/day in 1 or 2 divided doses) <b>to 3 mg/kg/day</b> with valproate alone	<b>4.5 to 7.5 mg/kg/day</b> (maximum 300 mg/day in 2 divided doses)	<b>5 to 15 mg/kg/day</b> (maximum 400 mg/day in 2 divided doses)
Maintenance dose in patients less than 30 kg	May need to be increased by as much as 20%, based on clinical response.	May need to be increased by as much as 20%, based on clinical response.	May need to be increased by as much as 20%, based on clinical response.

**Note:** Only whole tablets should be used for dosing.

\* Valproate has been shown to inhibit glucuronidation and decrease the apparent clearance of lamotrigine (see Drug Interactions (7), Clinical Pharmacology (12.3)).

† Drugs that induce lamotrigine glucuronidation and increase clearance, other than the specified antiepileptic drugs, include estrogen-containing oral contraceptives, rifampin, and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir. Dosing recommendations for oral contraceptives and the protease inhibitor atazanavir/ritonavir can be found in General Dosing Considerations (see Dosage and Administration (2.1)). Patients on rifampin and the protease inhibitor lopinavir/ritonavir should follow the same dosing titration/maintenance regimen used with antiepileptic drugs that induce glucuronidation and increase clearance (see Dosage and Administration (2.1), Drug Interactions (7), Clinical Pharmacology (12.3)).

**Table 3. The Initial Weight-Based Dosing Guide for Patients Aged 2 to 12 Years Taking Valproate (Weeks 1 to 4) with Epilepsy**

If the patient's weight is	Give this daily dose, using the most appropriate combination of lamotrigine 2 and 5-mg tablets		
	Greater than 14 kg	14 kg and less than 17 kg	17 kg and less than 25 kg
Greater than 14 kg	2 mg every other day	2 mg every day	2 mg every day
14 kg and less than 17 kg	2 mg every day	2 mg every day	4 mg every day
17 kg and less than 25 kg	4 mg every day	4 mg every day	8 mg every day
25 kg and less than 34 kg	4 mg every day	4 mg every day	8 mg every day
34 kg and less than 40 kg	5 mg every day	5 mg every day	10 mg every day

#### Usual Adjunctive Maintenance Dose for Epilepsy

The usual maintenance doses identified in Tables 1 and 2 are derived from dosing regimens employed in the placebo-controlled adjunctive trials in which the efficacy of SUBVENITE was established. In patients receiving multidrug regimens employing carbamazepine, phenytoin, phenobarbital, or primidone **adding valproate**, maintenance doses of adjunctive SUBVENITE as high as 700 mg/day have been used. In patients receiving **additive** adjunctive maintenance doses of adjunctive SUBVENITE as high as 200 mg/day have been used. The advantage of using doses above those recommended in Tables 1 to 4 has not been established in controlled trials.

#### 2.3 Epilepsy Conversion from Adjunctive Therapy to Monotherapy

The goal of the transition regimen is to attempt to maintain seizure control while mitigating the risk of serious rash associated with the rapid titration of SUBVENITE.

The recommended maintenance dose of SUBVENITE as monotherapy is 500 mg/day given in 2 divided doses.

To avoid an increased risk of rash, the recommended initial dose and subsequent dose escalation for SUBVENITE should not be exceeded (see Boxed Warning).

#### Conversion from Adjunctive Therapy with Carbamazepine, Phenytoin, Phenobarbital, or Primidone to Monotherapy with SUBVENITE

After achieving a dose of 500 mg/day of SUBVENITE using the guidelines in Table 1, the concomitant enzyme-inducing AED should be withdrawn by 20% decrements each week over a 4-week period. The regimen for the withdrawal of the concomitant AED is based on experience gained in the controlled monotherapy clinical trial.

#### Conversion from Adjunctive Therapy with Valproate to Monotherapy with SUBVENITE

The conversion regimen involves the 4 steps outlined in Table 4.

**Table 4. Conversion from Adjunctive Therapy with Valproate to Monotherapy with SUBVENITE in Patients Aged 16 Years and Older with Epilepsy**

	SUBVENITE	Valproate
Step 1	Achieve a dose of 200 mg/day according to guidelines in Table 1.	Maintain established stable dose.
Step 2	Maintain at 200 mg/day.	Decrease dose by decrements no greater than 500 mg/day/week to 500 mg/day and then maintain for 1 week.
Step 3	Increase to 300 mg/day and maintain for 1 week.	Simultaneously decrease to 250 mg/day and maintain for 1 week.
Step 4	Increase by 100 mg/day every week to achieve maintenance dose of 500 mg/day.	Discontinue.

#### Conversion from Adjunctive Therapy with Antiepileptic Drugs other than Carbamazepine, Phenytoin, Phenobarbital, Primidone, or Valproate to Monotherapy with SUBVENITE

No specific dosing guidelines can be provided for conversions to monotherapy with SUBVENITE with AEDs other than carbamazepine, phenytoin, phenobarbital, primidone, or valproate.

#### 2.4 Bipolar Disorder

The goal of maintenance treatment with SUBVENITE is to delay the time to occurrence of mood episodes (depression, mania, hypomania, mixed episodes) in patients treated for acute mood episodes with standard therapy (see Indications and Usage (1.2)).

Patients taking SUBVENITE for more than 16 weeks should be periodically reassessed to determine the need for maintenance treatment.

#### Adults

The target dose of SUBVENITE is 200 mg/day (100 mg/day in patients taking valproate, which decreases the apparent clearance of lamotrigine, and 400 mg/day in patients not taking valproate and taking either carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitor lopinavir/ritonavir that increase the apparent clearance of lamotrigine). In the clinical trials, doses up to 400 mg/day as monotherapy were evaluated; however, no additional benefit was seen at 400 mg/day compared with 200 mg/day (see Clinical Studies (14.2)). Accordingly, doses above 200 mg/day are not recommended.

Treatment with SUBVENITE is introduced, based on concurrent medication, according to the regimen outlined in Table 5. If other psychotropic medications are withdrawn following stabilization, the dose of SUBVENITE should be adjusted. In patients discontinuing valproate, the dose of SUBVENITE should be doubled over a 2-week period in equal weekly increments (see Table 6). In patients discontinuing carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir that induce lamotrigine glucuronidation, the dose of SUBVENITE should remain constant for the first week and then should be decreased by half over a 2-week period in equal weekly decrements (see Table 6). The dose of SUBVENITE may then be further adjusted to the target dose (200 mg) as clinically indicated.

If other drugs are subsequently initiated, the dose of SUBVENITE may need to be adjusted. In particular, the introduction of valproate requires reduction in the dose of SUBVENITE (see Drug Interactions (7), Clinical Pharmacology (12.3)).

To avoid an increased risk of rash, the recommended initial dose and subsequent dose escalation of SUBVENITE should not be exceeded (see Boxed Warning).

**Table 5. Escalation Regimens for SUBVENITE in Adults with Bipolar Disorder**

	In Patients TAKING Valproate <sup>a</sup> or Valproate <sup>a</sup>	In Patients NOT TAKING Carbamazepine, Phenytoin, Phenobarbital, Primidone, or Valproate <sup>a</sup>	In Patients TAKING Carbamazepine, Phenytoin, Phenobarbital, or Primidone and NOT TAKING Valproate <sup>a</sup>
Weeks 1 and 2	25 mg every other day	25 mg daily	50 mg daily
Weeks 1 and 2	50 mg daily	50 mg daily	100 mg daily, in divided doses
Week 5	50 mg daily	100 mg daily	200 mg daily, in divided doses
Week 6	100 mg daily	200 mg daily	300 mg daily, in divided doses
Week 7	100 mg daily	300 mg daily	400 mg daily, in divided doses

<sup>a</sup> Valproate has been shown to inhibit glucuronidation and decrease the apparent clearance of lamotrigine (see Drug Interactions (7), Clinical Pharmacology (12.3)).

† Drugs that induce lamotrigine glucuronidation and increase clearance, other than the specified antiepileptic drugs, include estrogen-containing oral contraceptives, rifampin, and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir. Dosing recommendations for oral contraceptives and the protease inhibitor atazanavir/ritonavir can be found in General Dosing Considerations (see Dosage and Administration (2.1)). Patients on rifampin and the protease inhibitor lopinavir/ritonavir should follow the same dosing titration/maintenance regimen used with antiepileptic drugs that induce glucuronidation and increase clearance (see Dosage and Administration (2.1), Drug Interactions (7), Clinical Pharmacology (12.3)).

**Table 6. Dose Adjustment to SUBVENITE in Adults with Bipolar Disorder Following Discontinuation of Psychotropic Medications**

	Discontinuation of Psychotropic Drugs (excluding Valproate, Carbamazepine, Phenytoin, Phenobarbital, or Primidone) <sup>a</sup>	Current Dose of Lamotrigine (mg/day)	
		100	400
Week 1	Maintain current dose of SUBVENITE	150	400
Week 2	Maintain current dose of SUBVENITE	200	400
Week 3 onward	Maintain current dose of SUBVENITE	200	200

<sup>a</sup> Valproate has been shown to inhibit glucuronidation and decrease the apparent clearance of lamotrigine (see Drug Interactions (7), Clinical Pharmacology (12.3)).

† Drugs that induce lamotrigine glucuronidation and increase clearance, other than the specified antiepileptic drugs, include estrogen-containing oral contraceptives, rifampin, and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir. Dosing recommendations for oral contraceptives and the protease inhibitor atazanavir/ritonavir can be found in General Dosing Considerations (see Dosage and Administration (2.1)). Patients on rifampin and the protease inhibitor lopinavir/ritonavir should follow the same dosing titration/maintenance regimen used with antiepileptic drugs that induce glucuronidation and increase clearance (see Dosage and Administration (2.1), Drug Interactions (7), Clinical Pharmacology (12.3)).

### 3. DOSAGE FORMS AND STRENGTHS

#### 3.1 Tablets

25 mg, White to off white, round shape, flat face beveled edge, uncoated tablets debossed with "2L" on one side and break line on other side.

100 mg, White to off white, round shape, flat face beveled edge, uncoated tablets debossed with "10LA" on one side and break line on other side.

150 mg, White to off white, round shape, flat face beveled edge, uncoated tablets debossed with "15LA" on one side and break line on other side.

200 mg, White to off white, round shape, flat face beveled edge, uncoated tablets debossed with "20LA" on one side and break line on other side.

#### 4 CONTRAINDICATIONS

SUBVENITE is contraindicated in patients who have demonstrated hypersensitivity (e.g., rash, angioedema, acute urticaria, extensive pruritus, mucocutaneous ulceration) to the drug or its ingredients (see Boxed Warning, Warnings and Precautions (5.1, 5.3)).

#### 5 WARNINGS AND PRECAUTIONS

##### Serious Skin Rashes (see Boxed Warning)

##### Pediatric Population

The incidence of serious rash associated with hospitalization and discontinuation of SUBVENITE in a prospectively followed cohort of pediatric patients (aged 2 to 17 years) is approximately 0.3% to 0.8%. One rash-related death was reported in a prospectively followed cohort of 1,263 pediatric patients (aged 2 to 16 years) with epilepsy taking SUBVENITE as adjunctive therapy. Additionally, there have been rare cases of toxic epidermal necrolysis with and without permanent sequelae and/or death in US and foreign postmarketing experience. There is evidence that the inclusion of valproate in a multidrug regimen increases the risk of serious, potentially life-threatening rash in pediatric patients. In pediatric patients who used valproate concomitantly for epilepsy, 1.2% (6 of 482) experienced a serious rash compared with 0.6% (6 of 952) patients not taking valproate.

##### Adult Population

Serious rash associated with hospitalization and discontinuation of SUBVENITE occurred in 0.3% (1 of 1,343) of adult patients who received SUBVENITE in premarketing clinical trials of epilepsy. In the bipolar and other mood disorders clinical trials, the rate of serious rash was 0.08% (1 of 1,233) of adult patients who received SUBVENITE as initial monotherapy and 0.13% (2 of 1,530) of adult patients who received SUBVENITE as adjunctive therapy. No fatalities occurred among these individuals. However, in worldwide postmarketing experience, rare cases of rash-related death have been reported, but their numbers are too few to permit a precise estimate of the rate.

Among the rashes leading to hospitalization were Stevens-Johnson syndrome, toxic epidermal necrolysis, angioedema, and those associated with multiorgan hypersensitivity (see Warnings and Precautions (5.3)).

There is evidence that the inclusion of valproate in a multidrug regimen increases the risk of serious, potentially life-threatening rash in adults. Specifically, of 584 patients administered SUBVENITE with

valproate in epilepsy clinical trials, 6 (1%) were hospitalized in association with rash. In contrast, 4 (0.16%) of 2,598 clinical trial patients and volunteers administered SUBVENTITE in the absence of valproate were hospitalized.

#### Patients with History of Allergy or Rash to Other Antiepileptic Drugs

The risk of serious rash may be increased when the recommended initial dose and/or the rate of dose escalation for SUBVENTITE is exceeded and in patients with a history of allergy or rash to other AEDs.

#### 5.2 Hemophagocytic Lymphohistiocytosis

Hemophagocytic lymphohistiocytosis (HLH) has occurred in pediatric and adult patients taking SUBVENTITE for various indications. HLH is a life-threatening syndrome of pathologic immune activation characterized by clinical signs and symptoms of extreme systemic inflammation. It is associated with high mortality rates if not recognized early and treated. Common findings include fever, hepatosplenomegaly, rash, lymphadenopathy, neurologic symptoms, cytopenias, high serum ferritin and liver function and coagulation abnormalities. In cases of HLH reported with SUBVENTITE, patients have presented with signs of systemic inflammation (fever, rash, hepatosplenomegaly, and organ system dysfunction) and blood dyscrasias. Symptoms have been reported to occur within 8 to 24 days following the initiation of treatment. Patients who develop early manifestations of pathologic immune activation should be evaluated immediately, and a diagnosis of HLH should be considered. SUBVENTITE should be discontinued if an alternative etiology for the signs or symptoms cannot be established.

#### 5.3 Multiorgan Hypersensitivity Reactions and Organ Failure

Multiorgan hypersensitivity reactions, also known as drug reaction with eosinophilia and systemic symptoms (DRESS), have occurred with SUBVENTITE. Some have been fatal or life threatening. DRESS typically, although not exclusively, presents with fever, rash, and/or lymphadenopathy in association with other organ system involvement, such as hepatitis, nephritis, hematologic abnormalities, myocarditis, or myositis, sometimes resembling an acute viral infection. Eosinophilia is often present. This disorder is variable in its expression, and other organ systems not noted here may be involved.

Fatality associated with acute multiorgan failure and various degrees of hepatic failure have been reported in 1 of 3,790 adult patients and 4 of 2,425 pediatric patients who received SUBVENTITE in epilepsy clinical trials. Rare fatalities from multiorgan failure have also been reported in postmarketing use.

Isolated liver failure without rash or involvement of other organs has also been reported with SUBVENTITE.

It is important to note that early manifestations of hypersensitivity (e.g., fever, lymphadenopathy) may be present even though a rash is not evident. If such signs or symptoms are present, the patient should be evaluated immediately. SUBVENTITE should be discontinued if an alternative etiology for the signs or symptoms cannot be established.

Prior to initiation of treatment with SUBVENTITE, the patient should be instructed that a rash or other signs or symptoms of hypersensitivity (e.g., fever, lymphadenopathy) may herald a serious medical event and that the patient should report any such occurrence to a healthcare provider immediately.

#### 5.4 Blood Dyscrasias

There have been reports of blood dyscrasias that may or may not be associated with multiorgan hypersensitivity (also known as DRESS) [see Warnings and Precautions (5.3)]. These have included neutropenia, leucopenia, anemia, thrombocytopenia, pancytopenia, and, rarely, aplastic anemia and pure red cell aplasia.

#### 5.5 Suicidal Behavior and Ideation

AEDs, including SUBVENTITE, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.

Pooled analyses of 199 placebo-controlled clinical trials (monotherapy and adjunctive therapy) of 11 different AEDs showed that patients randomized to 1 of the AEDs had approximately twice the risk (adjusted Relative Risk 1.8, 95% CI 1.2, 2.7) of suicidal thinking or behavior compared with patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence of suicidal behavior or ideation among 27,863 AED-treated patients was 0.43%, compared with 0.24% among 16,029 placebo-treated patients, representing an increase of approximately 1 case of suicidal thinking or behavior for every 530 patients treated. There were 4 suicides in drug-treated patients in the trials and none in placebo-treated patients, but the number of events is too small to allow any conclusion about drug effect on suicide.

The increased risk of suicidal thoughts or behavior with AEDs was observed as early as 1 week after starting treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behavior beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed.

The finding of an increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (6 to 100 years) in the clinical trials analyzed.

Table 7 shows absolute and relative risk by indication for all evaluated AEDs.

Table 7. Risk by Indication for Antiepileptic Drugs in the Pooled Analysis

Indication	Placebo Patients with Events per 1,000 Patients	Drug Patients with Events per 1,000 Patients	Relative Risk: Incidence of Events in Drug Patients/Incidence in Placebo Patients	Risk Difference: Additional Drug Patients with Events per 1,000 Patients
Epilepsy	1.0	1.8	1.8	0.8
Psychiatric	5.7	8.5	1.5	2.9
Other	1.0	1.9	1.9	0.9
Total	2.4	4.3	1.8	1.9

The relative risk for suicidal thoughts or behavior was higher in clinical trials for epilepsy than in clinical trials for psychiatric or other conditions, but the absolute risk differences were similar for the epilepsy and psychiatric indications.

Anyone considering prescribing SUBVENTITE or any other AED must balance the risks of suicidal thoughts or behavior with the risks of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior. Should suicidal thoughts and behavior emerge during treatment, the prescriber needs to consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

Patients, their caregivers, and families should be informed that AEDs increase the risk of suicidal thoughts and behavior and should be advised of the need to be alert for the emergence or worsening of the signs and symptoms of depression, any unusual changes in mood or behavior, the emergence of suicidal thoughts or suicidal behavior, or thoughts about self-harm. Behaviors of concern should be reported immediately to healthcare providers.

#### 5.6 Aseptic Meningitis

Therapy with SUBVENTITE increases the risk of developing aseptic meningitis. Because of the potential for serious outcomes of untreated meningitis due to other causes, patients should also be evaluated for other causes of meningitis and treated as appropriate.

Postmarketing cases of aseptic meningitis have been reported in pediatric and adult patients taking SUBVENTITE for various indications. Symptoms upon presentation have included headache, fever, nausea, vomiting, and neck rigidity. Rash, photophobia, myalgia, chills, altered consciousness, and somnolence were also noted in some cases. Symptoms have been reported to occur within 1 day to one and a half months following the initiation of treatment. In most cases, symptoms were reported to resolve after discontinuation of SUBVENTITE. Re-exposure resulted in a rapid return of symptoms (from within 30 minutes to 1 day following re-initiation of treatment) that were frequently more severe. Some of the patients treated with SUBVENTITE who developed aseptic meningitis had underlying diagnoses of systemic lupus erythematosus or other autoimmune diseases.

Cerebrospinal fluid (CSF) analyzed at the time of clinical presentation in reported cases was characterized by a mild to moderate pleocytosis, normal glucose levels, and mild to moderate increase in protein. CSF white blood cell count differentials showed a predominance of neutrophils in a majority of the cases, although a predominance of lymphocytes was reported in approximately one third of the cases. Some patients also had new onset of signs and symptoms of involvement of other organs (predominantly hepatic and renal involvement), which may suggest that in these cases the aseptic meningitis observed was part of a hypersensitivity reaction [see Warnings and Precautions (5.3)].

#### 5.7 Potential Medication Errors

Medication errors involving SUBVENTITE have occurred. In particular, the name SUBVENTITE can be confused with the names of other commonly used medications. Medication errors may also occur between the different formulations of SUBVENTITE. To reduce the potential of medication errors, write and say lamotrigine clearly. Depiction of the SUBVENTITE can be found in the Medication Guide that accompanies the product to highlight the distinctive markings, colors, and shapes that serve to identify the different presentations of the drug and thus may help reduce the risk of medication errors. To avoid medication error of using the wrong drug or formulation, patients should be strongly advised to visually inspect their tablets to verify that they are SUBVENTITE, as well as the correct formulation of SUBVENTITE, each time they fill their prescription.

#### 5.8 Concomitant Use with Oral Contraceptives

Some estrogen-containing oral contraceptives have been shown to decrease serum concentrations of lamotrigine [see Clinical Pharmacology (12.3)]. Dosage adjustments will be necessary in most patients who start or stop estrogen-containing oral contraceptives while taking SUBVENTITE [see Dosage and Administration (2.1)]. During the week of inactive hormone preparation (pill-free week) of oral contraceptive therapy, plasma lamotrigine levels are expected to rise, as much as doubling at the end of the week. Adverse reactions consistent with elevated levels of lamotrigine, such as dizziness, nausea, and diplopia, could occur.

#### 5.9 Withdrawal Seizures

As with other AEDs, SUBVENTITE should not be abruptly discontinued. In patients with epilepsy there is a possibility of increasing seizure frequency. In clinical trials in adults with bipolar disorder, 2 patients experienced seizures shortly after abrupt withdrawal of SUBVENTITE. Unless safety concerns require a more rapid withdrawal, the dose of SUBVENTITE should be tapered over a period of at least 2 weeks (approximately 50% reduction per week) [see Dosage and Administration (2.1)].

#### 5.10 Status Epilepticus

Valid estimates of the incidence of treatment-emergent status epilepticus among patients treated with SUBVENTITE are difficult to obtain because reporters participating in clinical trials did not all employ identical rules for identifying cases. At a minimum, 7 of 2,343 adult patients had episodes that could unequivocally be described as status epilepticus. In addition, a number of reports of variously defined episodes of seizure exacerbation (e.g., seizure clusters, seizure flurries) were made.

#### 5.11 Sudden Unexplained Death in Epilepsy (SUDEP)

During the premarketing development of SUBVENTITE, 20 sudden and unexplained deaths were recorded among a cohort of 4,700 patients with epilepsy (0.47 patients/year of exposure).

Some of these could represent seizure-related deaths in which the seizure was not observed, e.g., at night. This represents an incidence of 0.0035 deaths per patient-year. Although this rate exceeds that expected in a healthy population matched for age and sex, it is within the range of estimates for the incidence of sudden unexplained death in epilepsy (SUDEP) in patients not receiving SUBVENTITE (ranging from 0.0005 for the general population of patients with epilepsy, to 0.004 for a recently studied clinical trial population similar to that in the clinical development program for SUBVENTITE, to 0.005 for patients with refractory epilepsy). Consequently, whether these figures are reassuring or suggest concern depends on the comparability of the population reported upon with the cohort receiving SUBVENTITE and the accuracy of the estimates provided. Probably most reassuring is the similarity of estimated SUDEP rates in patients receiving SUBVENTITE and those receiving other AEDs, chemically unrelated to each other, that underwent clinical testing in similar populations. This evidence suggests, although it certainly does not prove, that the high SUDEP rates reflect population rates, not a drug effect.

#### 5.12 Addition of SUBVENTITE to a Multidrug Regimen that Includes Valproate

Because valproate reduces the clearance of SUBVENTITE, the dose of lamotrigine in the presence of valproate is less than half of that required in its absence [see Dosage and Administration (2.2, 2.3, 2.4), Drug Interactions (7)].

#### 5.13 Binding in the Eye and Other Melanin-Containing Tissues

Because lamotrigine binds to melanin, it could accumulate in melanin-rich tissues over time. This raises the possibility that lamotrigine may cause toxicity in these tissues after extended use. Although ophthalmological testing was performed in a controlled clinical trial, the testing was inadequate to exclude subtle effects or injury occurring after long-term exposure. Moreover, the capacity of available tests to detect potentially adverse consequences, if any, of lamotrigine's binding to melanin is unknown [see Clinical Pharmacology (12.2)].

Accordingly, although there are no specific recommendations for periodic ophthalmological monitoring, prescribers should be aware of the possibility of long-term ophthalmologic effects.

#### 5.14 Laboratory Tests

##### False-Positive Drug Test Results

Lamotrigine has been reported to interfere with the assay used in some rapid urine drug screens, which can result in false-positive readings, particularly for phenylethylamine (PCP). A more specific analytical method should be used to confirm a positive result.

##### Plasma Concentrations of Lamotrigine

The value of monitoring plasma concentrations of lamotrigine in patients treated with SUBVENITE has not been established. Because of the possible pharmacokinetic interactions between lamotrigine and other drugs, including AEDs (see Table 13), monitoring of the plasma levels of lamotrigine and concomitant drugs may be indicated, particularly during dosage adjustments. In general, clinical judgment should be exercised regarding monitoring of plasma levels of lamotrigine and other drugs and whether or not dosage adjustments are necessary.

**6 ADVERSE REACTIONS**

The following adverse reactions are described in more detail in the Warnings and Precautions section of the label:

- Serious skin rashes [see Warnings and Precautions (5.1)]
- Hemophagocytic Lymphohistiocytosis [see Warnings and Precautions (5.2)]
- Multisystem hypersensitivity reactions and organ failure [see Warnings and Precautions (5.3)]
- Blood dyscrasias [see Warnings and Precautions (5.4)]
- Suicidal behavior and ideation [see Warnings and Precautions (5.5)]
- Aseptic meningitis [see Warnings and Precautions (5.6)]
- Withdrawal seizures [see Warnings and Precautions (5.9)]
- Status epilepticus [see Warnings and Precautions (5.10)]
- Sudden unexpected death in epilepsy [see Warnings and Precautions (5.11)]

**6.1 Clinical Trial Experience**

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared with rates in the clinical trials of another drug and may not reflect the rates observed in practice.

**Epilepsy**

**Most Common Adverse Reactions in All Clinical Trials: Adjunctive Therapy in Adults with Epilepsy:** The most commonly observed (>5% for SUBVENITE and more common on drug than placebo) adverse reactions seen in association with SUBVENITE during adjunctive therapy in adults and not seen at an equivalent frequency among placebo-treated patients were: dizziness, ataxia, somnolence, headache, diplopia, blurred vision, nausea, vomiting, and rash. Dizziness, diplopia, ataxia, blurred vision, nausea, and vomiting were dose related. Dizziness, diplopia, ataxia, and blurred vision occurred more commonly in patients receiving carbamazepine with SUBVENITE than in patients receiving other AEDs with SUBVENITE. Clinical data suggest a higher incidence of rash, including serious rash, in patients receiving concomitant valproate than in patients not receiving valproate [see Warnings and Precautions (5.1)].

Approximately 11% of the 3,378 adult patients who received SUBVENITE as adjunctive therapy in premarketing clinical trials discontinued treatment because of an adverse reaction. The adverse reactions most commonly associated with discontinuation were rash (3.0%), dizziness (2.8%), and headache (2.5%).

In a dose-response trial in adults, the rate of discontinuation of SUBVENITE for dizziness, ataxia, diplopia, blurred vision, nausea, and vomiting was dose related.

**Monotherapy in Adults with Epilepsy:** The most commonly observed (>5% for SUBVENITE and more common on drug than placebo) adverse reactions seen in association with the use of SUBVENITE during the monotherapy phase of the controlled trial in adults not seen at an equivalent rate in the control group were vomiting, coordination abnormality, dyspepsia, nausea, dizziness, rhinitis, anxiety, insomnia, infection pain, weight decrease, chest pain, and dyspareunia. The most commonly observed (>5% for SUBVENITE and more common on drug than placebo) adverse reactions associated with the use of SUBVENITE during the conversion to monotherapy (add-on) period, not seen at an equivalent frequency among low-dose valproate-treated patients, were dizziness, headache, nausea, ataxia, coordination abnormality, vomiting, rash, somnolence, diplopia, ataxia, accidental injury, tremor, blurred vision, insomnia, myalgia, diarrhea, lymphadenopathy, gastritis, and sinusitis.

Approximately 10% of the 420 adult patients who received SUBVENITE as monotherapy in premarketing clinical trials discontinued treatment because of an adverse reaction. The adverse reactions most commonly associated with discontinuation were rash (4.5%), headache (3.1%), and ataxia (2.4%).

**Adjunctive Therapy in Pediatric Patients with Epilepsy:** The most commonly observed (>5% for SUBVENITE and more common on drug than placebo) adverse reactions seen in association with the use of SUBVENITE as adjunctive treatment in pediatric patients aged 2 to 16 years and not seen at an equivalent rate in the control group were infection, vomiting, rash, fever, somnolence, accidental injury, dizziness, diarrhea, abdominal pain, nausea, ataxia, tremor, ataxia, bronchitis, flu syndrome, and diplopia.

In 339 patients aged 2 to 16 years with partial-onset seizures or generalized seizures of Lennox-Gastaut syndrome, 4.2% of patients on lamotrigine and 2.9% of patients on placebo discontinued due to adverse reactions. The most commonly reported adverse reaction that led to discontinuation of lamotrigine was rash.

Approximately 11.5% of the 1,081 pediatric patients aged 2 to 16 years who received SUBVENITE as adjunctive therapy in premarketing clinical trials discontinued treatment because of an adverse reaction. The adverse reactions most commonly associated with discontinuation were rash (4.4%), reaction aggravated (1.7%), and ataxia (0.6%).

**Controlled Adjunctive Clinical Trials in Adults with Epilepsy:** Table 8 lists adverse reactions that occurred in adult patients with epilepsy treated with SUBVENITE in placebo-controlled trials. In these trials, either SUBVENITE or placebo was added to the patient's current AED therapy.

**Table 8. Adverse Reactions in Placebo-, Placebo-Controlled Adjunctive Trials in Adult Patients with Epilepsy<sup>a,b</sup>**

Body System/Adverse Reaction	Percent of Patients Receiving Adjunctive SUBVENITE (n = 711)	Percent of Patients Receiving Adjunctive Placebo (n = 419)
Body as a whole		
Headache	29	19
Flu syndrome	7	6
Fever	6	4
Abdominal pain	5	4
Neck pain	2	1
Reaction aggravated (asthma exacerbation)	2	1
Digestive		
Nausea	19	10
Vomiting	10	4
Diarrhea	6	4
Dyspepsia	5	2
Constipation	4	3
Anorexia	2	1
Musculoskeletal		
Arthralgia	2	0
Nervous		
Dizziness	38	13
Ataxia	22	6
Somnolence	7	14
Incoordination	6	2
Insomnia	6	2
Tremor	4	1
Depression	4	3
Anxiety	4	3
Lethargy	3	1
Coordination	3	2
Irritability	3	0
Speech disorder	3	0
Concentration disturbance	2	1
Respiratory		
Rhinitis	14	9
Pharyngitis	10	9
Cough increased	8	6
Skin and appendages		
Rash	10	5
Pruritus	3	2
Special senses		
Diplopia	26	7
Blurred vision	16	5
Vision abnormality	3	1
Urogenital		
Female patients only	(n = 365)	(n = 207)
Dysmenorrhea	7	6
Vaginitis	4	1
Amenorrhea	2	1

<sup>a</sup> Adverse reactions that occurred in at least 2% of patients treated with SUBVENITE and at a greater incidence than placebo.  
<sup>b</sup> Patients in these adjunctive trials were receiving 1% of the concomitant antiepileptic drugs carbamazepine, phenytoin, phenobarbital, or primidone in addition to SUBVENITE or placebo. Patients may have reported multiple adverse reactions during the trial or at discontinuation; thus, patients may be included in more than 1 category.

In a randomized, parallel trial comparing placebo with 300 and 500 mg/day of SUBVENITE, some of the more common drug-related adverse reactions were dose related (see Table 9).

**Table 9. Dose-Related Adverse Reactions from a Randomized, Placebo-Controlled Adjunctive Trial in Adults with Epilepsy<sup>a</sup>**

Adverse Reaction	Percent of Patients Experiencing Adverse Reactions		
	Placebo (n = 73)	SUBVENITE 300 mg (n = 71)	SUBVENITE 500 mg (n = 72)
Ataxia	10	10	10
Blurred vision	10	11	25 <sup>b</sup>
Diplopia	8	24 <sup>a</sup>	49 <sup>a</sup>
Dizziness	22	31	54 <sup>a</sup>
Nausea	11	18	25 <sup>a</sup>
Vomiting	11	11	18 <sup>a</sup>

<sup>a</sup> Significantly greater than placebo group (P<0.05).  
<sup>b</sup> Significantly greater than group receiving SUBVENITE 300 mg (P<0.05).

The overall adverse reaction profile for SUBVENITE was similar between females and males and was independent of age. Because the largest non-Caucasian racial subgroup was only 0% of patients exposed to SUBVENITE in placebo-controlled trials, there are insufficient data to support a statement regarding the distribution of adverse reaction reports by race. Generally, females receiving either SUBVENITE as adjunctive therapy or placebo were more likely to report adverse reactions than males. The only adverse reaction for which the reports on SUBVENITE were > 10% more frequent in females than males (without a corresponding difference by gender on placebo) was dizziness (difference = 16.5%). There was little difference between females and males in the rates of discontinuation of SUBVENITE for individual adverse reactions.

**Controlled Monotherapy Trial in Adults with Partial-Onset Seizures:** Table 10 lists adverse reactions that occurred in patients with epilepsy treated with monotherapy with SUBVENITE in a double-blind trial following discontinuation of either concomitant carbamazepine or phenytoin not seen at an equivalent frequency in the control group.

**Table 10. Adverse Reactions in a Controlled Monotherapy Trial in Adult Patients with Partial-Onset Seizures<sup>a,b</sup>**

Body System/Adverse Reaction	Percent of Patients Receiving SUBVENITE <sup>c</sup> as Monotherapy (n = 43)	Percent of Patients Receiving Low-Dose Valproate Monotherapy (n = 44)
Body as a whole		
Pain	5	0
Infection	5	2
Chest pain	5	2
Digestive		
Vomiting	9	0
Dyspepsia	7	2
Nausea	7	2
Metabolic and nutritional		
Weight decrease	5	2
Nervous		
Coordination abnormality	7	0
Dizziness	7	0
Anxiety	5	0
Insomnia	5	2
Respiratory		
Rhinitis	7	2
Urogenital (female patients only)	(n = 21)	(n = 28)
Dysmenorrhea	7	0

<sup>a</sup> Adverse reactions that occurred in at least 5% of patients treated with SUBVENITE and at a greater incidence than valproate-treated patients.  
<sup>b</sup> Patients in this trial were converted to SUBVENITE or valproate monotherapy from adjunctive therapy with carbamazepine or phenytoin. Patients may have reported multiple adverse reactions during the trial; thus, patients may be included in more than 1 category.  
<sup>c</sup> Up to 500 mg/day.  
<sup>d</sup> 1,000 mg/day.

Adverse reactions that occurred with a frequency of < 5% and > 2% of patients receiving SUBVENITE and numerically more frequent than placebo were:

Body as a Whole: Ataxia, fever.  
 Digestive: Anorexia, dry mouth, rectal hemorrhage, peptic ulcer.  
 Metabolic and Nutritional: Peripheral edema.

**Nervous System:** Amnesia, ataxia, depression, hyperreflexia, libido increase, decreased reflexes, increased reflexes, myasthenia, irritability, suicidal ideation.

**Respiratory:** Epistaxis, bronchitis, dyspnea.

**Skin and Appendages:** Contact dermatitis, dry skin, sweating.

**Special Senses:** Vision abnormality.

**Incidence in Controlled Adjuvant Trials in Pediatric Patients with Epilepsy:** Table 11 lists adverse reactions that occurred in 339 pediatric patients with partial-onset seizures or generalized seizures of Lennox-Gastaut syndrome who received SUBVENTE up to 15 mg/kg/day or a maximum of 750 mg/day.

**Table 11. Adverse Reactions in Pooled, Placebo-Controlled Adjuvant Trials in Pediatric Patients with Epilepsy\***

Body System/Adverse Reaction	Percent of Patients Receiving SUBVENTE (n = 160)	Percent of Patients Receiving Placebo (n = 171)
<b>Body as a whole</b>		
Infection	20	17
Fever	15	14
Accidental injury	14	12
Abdominal pain	10	5
Arthralgia	8	4
Flu syndrome	7	6
Pain	5	4
Facial edema	2	1
Photosensitivity	2	0
<b>Cardiovascular</b>		
Hemorrhage	2	1
<b>Digestive</b>		
Vomiting	20	16
Diarrhea	11	9
Nausea	10	2
Constipation	4	2
Dyspepsia	2	1
<b>Hemic and lymphatic</b>		
Lymphadenopathy	2	1
<b>Metabolic and nutritional</b>		
Edema	2	0
<b>Nervous system</b>		
Somnolence	17	15
Dizziness	14	4
Ataxia	11	3
Tremor	10	1
Emotional lability	4	2
Gait abnormality	4	2
Thinking abnormality	3	2
Convulsions	2	1
Nervousness	2	1
Vertigo	2	1
<b>Respiratory</b>		
Pharyngitis	14	11
Bronchitis	7	5
Increased cough	7	6
Sinusitis	2	1
Bronchospasm	2	1
<b>Skin</b>		
Rash	14	12
Eczema	2	1
Pruritus	2	1
<b>Special senses</b>		
Diplopia	5	1
Blurred vision	4	1
Vision abnormality	2	0
<b>Urogenital</b>		
Male and female patients		
Urinary tract infection	3	0

\* Adverse reactions that occurred in at least 2% of patients treated with SUBVENTE and at a greater incidence than placebo.

**Bipolar Disorder in Adults**

The most common adverse reactions seen in association with the use of SUBVENTE as monotherapy (100 to 400 mg/day) in adult patients (aged 18 to 82 years) with bipolar disorder in the 2 double-blind placebo-controlled trials of 18 months' duration are included in Table 12. Adverse reactions that occurred in at least 5% of patients and were numerically more frequent during the dose-escalation phase of SUBVENTE in these trials (when patients may have been receiving concomitant medications) compared with the monotherapy phase were: headache (27%), rash (11%), dizziness (10%), diarrhea (8%), dream abnormality (6%), and pruritus (6%).

During the monotherapy phase of the double-blind placebo-controlled trials of 18 months' duration, 13% of 227 patients who received SUBVENTE (100 to 400 mg/day), 16% of 190 patients who received placebo, and 23% of 166 patients who received lithium discontinued therapy because of an adverse reaction. The adverse reactions that most commonly led to discontinuation of SUBVENTE were rash (3%) and mania/hypomania/mixed mood adverse reactions (2%). Approximately 16% of 2,401 patients who received SUBVENTE (50 to 500 mg/day) for bipolar disorder in premarketing trials discontinued therapy because of an adverse reaction, most commonly due to rash (5%) and mania/hypomania/mixed mood adverse reactions (2%).

The overall adverse reaction profile for SUBVENTE was similar between females and males, between elderly and nonelderly patients, and among racial groups.

**Table 12. Adverse Reactions in 2 Placebo-Controlled Trials in Adult Patients with Bipolar Disorder\*\***

Body System/Adverse Reaction	Percent of Patients Receiving SUBVENTE (n = 227)	Percent of Patients Receiving Placebo (n = 190)
<b>General</b>		
Back pain	8	6
Fatigue	8	5
Abdominal pain	6	3
<b>Digestive</b>		
Nausea	14	11
Constipation	5	2
Vomiting	5	2
<b>Nervous System</b>		
Insomnia	10	6
Somnolence	9	7
Nervousness (dry mouth)	6	4
<b>Respiratory</b>		
Rhinitis	7	4
Exacerbation of cough	5	3
Pharyngitis	5	4
<b>Skin</b>		
Rash (nummular)*	7	5

\* Adverse reactions that occurred in at least 5% of patients treated with SUBVENTE and at a greater incidence than placebo.

\*\* Patients in these trials were converted to SUBVENTE (100 to 400 mg/day) or placebo monotherapy from add-on therapy with other psychotropic medications. Patients may have reported multiple adverse reactions during the trial; thus, patients may be included in more than 1 category.

† In the overall bipolar and other mood disorders clinical trials, the rate of serious rash was 0.08% (1 of 1,233) of adult patients who received SUBVENTE as initial monotherapy and 0.13% (2 of 1,538) of adult patients who received SUBVENTE as adjunctive therapy [see Warnings and Precautions (5.1)].

Other reactions that occurred in 5% or more patients but equally or more frequently in the placebo group included: dizziness, mania, headache, infection, influenza, pain, accidental injury, diarrhea, and dyspepsia.

Adverse reactions that occurred with a frequency of < 5% and > 1% of patients receiving SUBVENTE and numerically more frequent than placebo were:

**General:** Fever, neck pain.

**Cardiovascular:** Migraine.

**Digestive:** Flatulence.

**Metabolic and Nutritional:** Weight gain, edema.

**Musculoskeletal:** Arthralgia, myalgia.

**Nervous System:** Amnesia, depression, agitation, emotional lability, dyspraxia, abnormal thoughts, dream abnormality, hyposthesia.

**Respiratory:** Sinusitis.

**Urogenital:** Urinary frequency.

**Adverse Reactions following Abrupt Discontinuation:** In the 2 controlled clinical trials, there was no increase in the incidence, severity, or type of adverse reactions in patients with bipolar disorder after abruptly terminating therapy with SUBVENTE. In the clinical development program in adults with bipolar disorder, 2 patients experienced seizures shortly after abrupt withdrawal of SUBVENTE [see Warnings and Precautions (5.3)].

**Mania/Hypomania/Mixed Episodes:** During the double-blind placebo-controlled clinical trials in bipolar I disorder in which adults were converted to monotherapy with SUBVENTE (100 to 400 mg/day) from other psychotropic medications and followed for up to 18 months, the rates of mania or hypomanic or mixed mood episodes reported as adverse reactions were 5% for patients treated with SUBVENTE (n = 227), 4% for patients treated with lithium (n = 166), and 7% for patients treated with placebo (n = 190). In all bipolar controlled trials combined, adverse reactions of mania (including hypomania and mixed mood episodes) were reported in 5% of patients treated with SUBVENTE (n = 956), 3% of patients treated with lithium (n = 200), and 4% of patients treated with placebo (n = 803).

**6.2 Other Adverse Reactions Observed in All Clinical Trials**

SUBVENTE has been administered to 6,694 individuals for whom complete adverse reaction data was captured during all clinical trials, only some of which were placebo controlled. During these trials, all adverse reactions were recorded by the clinical investigators using terminology of their own choosing.

To provide a meaningful estimate of the proportion of individuals having adverse reactions, similar types of adverse reactions were grouped into a smaller number of standardized categories using modified COSTART dictionary terminology. The frequencies presented represent the proportion of the 6,694 individuals exposed to SUBVENTE who experienced an event of the type cited on at least 1 occasion while receiving SUBVENTE. All reported adverse reactions are included except those already listed in the previous tables or elsewhere in the labeling; those too general to be informative, and those not reasonably associated with the use of the drug.

Adverse reactions are further classified within body system categories and enumerated in order of decreasing frequency using the following definitions: frequent adverse reactions are defined as those occurring in at least 1/100 patients; infrequent adverse reactions are those occurring in 1/100 to 1/1,000 patients; rare adverse reactions are those occurring in fewer than 1/1,000 patients.

**Body as a Whole**

**Infrequent:** Allergic reaction, chills, malaise.

**Cardiovascular System**

**Infrequent:** Flushing, hot flashes, hypertension, palpitations, postural hypotension, syncope, tachycardia, vasodilation.

**Dermatological**

**Infrequent:** Acne, alopecia, hirsutism, maculopapular rash, skin discoloration, urticaria. **Rare:** Angioedema, erythema, exfoliative dermatitis, fungal dermatitis, herpes zoster, leukoderma, multiforme erythema, petechial rash, pustular rash, Stevens-Johnson syndrome, vesiculobullous rash.

**Digestive System**

**Infrequent:** Dysphagia, eructation, gastritis, gingivitis, increased appetite, increased salivation, liver function tests abnormal, mouth ulceration.

**Rare:** Gastrointestinal hemorrhage, glossitis, gum hemorrhage, gum hyperplasia, hematemesis, hemorrhagic colitis, hepatitis, melena, stomach ulcer, stomatitis, tongue edema.

**Endocrine System**

**Rare:** Goiter, hypothyroidism.

**Hematologic and Lymphatic System**

**Infrequent:** Ecchymosis, leukopenia.

**Rare:** Anemia, eosinophilia, fibrin decrease, fibrinogen decrease, iron deficiency anemia, leukocytosis, lymphocytosis, macrocytic anemia, pernicia, thrombocytopenia.

**Metabolic and Nutritional Disorders**

**Infrequent:** Aspartate transaminase increased.

**Rare:** Alcohol intolerance, alkaline phosphatase increase, alanine transaminase increase, bilirubinemia, general edema, gamma glutamyl transpeptidase increase, hyperglycemia.

**Musculoskeletal System**

**Infrequent:** Arthritis, leg cramps, myasthenia, twitching.

**Rare:** Bursitis, muscle atrophy, pathological fracture, tendinous contracture.

**Nervous System**

**Frequent:** Confusion, paresthesia.

**Infrequent:** Anhedonia, apathy, aphasia, central nervous system depression, depersonalization, dysarthria, dyskinesia, euphoria, hallucinations, hostility, hyperkinesia, hypertonia, libido decreased, memory decrease, mind racing, movement disorder, myoclonus, panic attack, paranoid reaction, personality disorder, psychosis, sleep disorder, stupor, suicidal ideation.

**Rare:** Chorea/athetosis, delirium/delusions, dysphoria, dystonia, extrapyramidal syndrome, faintness, grand mal convulsions, hemiplegia, hyperalgesia, hyperesthesia, hypokinesia, hypotonia, manic depression reaction, muscle spasm, neuralgia, neuritis, paralysis, peripheral neuritis.

**Respiratory System**

**Infrequent:** Yawn

**Rare:** Hiccup, hyperventilation

**Special Senses**

**Frequent:** Amblyopia

**Infrequent:** Abnormality of accommodation, conjunctivitis, dry eyes, ear pain, photophobia, taste perversion, tinnitus.

**Rare:** Deafness, lacrimation disorder, oscillopsia, parosmia, ptosis, strabismus, taste loss, uveitis, visual field defect.

**Urinary System**

**Infrequent:** Abnormal ejaculation, hematuria, impotence, menorrhagia, polyuria, urinary incontinence.

**Rare:** Acute kidney failure, arnigostris, breast abscess, breast neoplasm, creatinine increase, cystitis, dysuria, epididymitis, female lactation, kidney failure, kidney pain, nocturia, urinary retention, urinary urgency.

**6.3 Postmarketing Experience**

The following adverse reactions have been identified during postapproval use of SUBVENITE. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

**Blood and Lymphatic**

Agranulocytosis, hemolytic anemia, lymphadenopathy not associated with hypersensitivity disorder.

**Gastrointestinal**

Eosinophilic esophagitis.

**Hepatobiliary Tract and Pancreas**

Pancreatitis.

**Immunologic**

Lupus-like reaction, vasculitis.

**Lower Respiratory**

Aphasia.

**Musculoskeletal**

Rhabdomyolysis has been observed in patients experiencing hypersensitivity reactions.

**Nervous System**

Aggression, exacerbation of Parkinsonian symptoms in patients with pre-existing Parkinson's disease, tics.

**Neutropenic**

Progressive immunosuppression.

**7 DRUG INTERACTIONS**

Significant drug interactions with SUBVENITE are summarized in this section.

Uridine 5'-diphospho-glucuronyl transferase (UGT) have been identified as the enzymes responsible for metabolism of lamotrigine. Drugs that induce or inhibit glucuronidation may, therefore, affect the apparent clearance of lamotrigine. Strong or moderate inducers of the cytochrome P450 3A4 (CYP3A4) enzyme, which is also known to induce UGT, may also enhance the metabolism of lamotrigine.

Those drugs that have been demonstrated to have a clinically significant impact on lamotrigine metabolism are outlined in Table 13. Specific dosing guidance for these drugs is provided in the Dosage and Administration section [see Dosage and Administration (2.1)].

Additional details of these drug interaction studies are provided in the Clinical Pharmacology section [see Clinical Pharmacology (12.3)].

**Table 13. Established and Other Potentially Significant Drug Interactions**

Concomitant Drug	Effect on Concentration of SUBVENITE or Concomitant Drug	Clinical Comment
Estrogen-containing oral contraceptive preparations containing 30 mg ethinylestradiol and 0.02 mg levonorgestrel	↓ lamotrigine ↓ levonorgestrel	Decreased lamotrigine concentration approximately 50%. Decrease in levonorgestrel component by 19%.
Carbamazepine and carbamazepine epoxide	↓ lamotrigine ↓ carbamazepine epoxide	Addition of carbamazepine decreases lamotrigine concentration approximately 40%. May increase carbamazepine epoxide levels.
Opavivir/ritonavir	↓ lamotrigine	Decreased lamotrigine concentration approximately 50%.
Atazanavir/ritonavir	↓ lamotrigine	Decreased lamotrigine AUC approximately 32%.
Phenobarbital/phenytoin	↓ lamotrigine	Decreased lamotrigine concentration approximately 40%.
Phenytoin	↓ lamotrigine	Decreased lamotrigine concentration approximately 40%.
Rifampin	↓ lamotrigine	Decreased lamotrigine AUC approximately 40%.
Valproate	↑ lamotrigine ? valproate	Increased lamotrigine concentrations slightly more than 2-fold. There are conflicting study results regarding effect of lamotrigine on valproate concentrations: 1) a mean 25% decrease in valproate concentrations in healthy volunteers, 2) no change in valproate concentration in controlled clinical trials in patients with epilepsy.
↑: Decreased (induces lamotrigine glucuronidation) ↓: Increased (inhibits lamotrigine glucuronidation) ?: Conflicting data.		

**Effect of SUBVENITE on Organic Cationic Transporter 2 Substrates**

Lamotrigine is an inhibitor of renal tubular secretion via organic cationic transporter 2 (OCT2) protein [see Clinical Pharmacology (12.3)]. This may result in increased plasma levels of certain drugs that are substantially excreted via this route. Coadministration of SUBVENITE with OCT2 substrates with a narrow therapeutic index (e.g., dofetilide) is not recommended.

**8 USE IN SPECIFIC POPULATIONS**

**8.1 Pregnancy**

As with other AEDs, physiological changes during pregnancy may affect lamotrigine concentrations and/or therapeutic effect. There have been reports of decreased lamotrigine concentrations during pregnancy and restoration of pre-pregnancy concentration after delivery. Dosage adjustments may be necessary to maintain clinical response.

**Pregnancy Category C**

There are no adequate and well-controlled studies in pregnant women. In animal studies, lamotrigine was developmentally toxic at doses lower than those administered clinically. SUBVENITE should be used during pregnancy only if the potential benefit justifies the potential risks to the fetus. When lamotrigine was administered to pregnant mice, rats, or rabbits during the period of organogenesis (oral doses of up to 125, 25, and 30 mg/kg, respectively), reduced fetal body weight and increased incidences of fetal skeletal variations were seen in mice and rats at doses that were also maternally toxic. The no-effect doses for embryofetal developmental toxicity in mice, rats, and rabbits (75, 6.25, and 30 mg/kg, respectively) are similar to (mice and rabbits) or less than (rats) the human dose of 400 mg/day on a body surface area (mg/m<sup>2</sup>) basis.

In a study in which pregnant rats were administered lamotrigine (oral doses of 5 or 25 mg/kg) during the period of organogenesis and offspring were evaluated postnatally, behavioral abnormalities were observed in exposed offspring at both doses. The lowest effect dose for developmental neurotoxicity in rats is less than the human dose of 400 mg/day on a mg/m<sup>2</sup> basis. Maternal toxicity was observed at the higher dose tested.

When pregnant rats were administered lamotrigine (oral doses of 5, 10, or 20 mg/kg) during the latter part of gestation, increased offspring mortality (including stillbirths) was seen at all doses. The lowest effect dose for perinatal developmental toxicity in rats is less than the human dose of 400 mg/day on a mg/m<sup>2</sup> basis. Maternal toxicity was observed at the 2 highest doses tested.

Lamotrigine decreases fetal folate concentrations in rat, an effect known to be associated with adverse pregnancy outcomes in animals and humans.

**Pregnancy Registry**

To provide information regarding the effects of in utero exposure to SUBVENITE, physicians are advised to recommend that pregnant patients taking SUBVENITE enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry. This can be done by calling the toll-free number 1-888-233-2334 and must be done by patients themselves. Information on the registry can also be found at the website <http://www.aedpregnancyregistry.org>.

**8.2 Lactation**

**Risk Summary**

Lamotrigine is present in milk from lactating women taking SUBVENITE (see Data). Neonates and young infants are at risk for high serum levels because maternal serum and milk levels can rise to high levels postpartum if lamotrigine dosage has been increased during pregnancy but is not reduced after delivery to the pre-pregnancy dosage. Glucuronidation is required for drug clearance. Glucuronidation capacity is immature in the infant and this may also contribute to the level of lamotrigine exposure. Events including rash, apnea, drowsiness, poor sucking, and poor weight gain (requiring hospitalization in some cases) have been reported in infants who have been milk-fed by mothers using lamotrigine; whether or not these events caused by lamotrigine is unknown. No data are available on the effects of the drug on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for SUBVENITE and any potential adverse effects on the breastfed infant from SUBVENITE or from the underlying maternal condition.

**Clinical Consideration**

Human milk-fed infants should be closely monitored for adverse events resulting from lamotrigine. Measurement of infant serum levels should be performed to rule out toxicity if concerns arise. Human milk-feeding should be discontinued in infants with lamotrigine toxicity.

**Data**

Data from multiple small studies indicate that lamotrigine plasma levels in nursing infants have been reported to be as high as 50% of maternal plasma concentrations.

**8.4 Pediatric Use**

**Indications**

SUBVENITE is indicated as adjunctive therapy in patients aged 2 years and older for partial-onset seizures, the generalized seizures of Lennox-Gastaut syndrome, and PGTC seizures.

Safety and efficacy of SUBVENITE used as adjunctive treatment for partial-onset seizures were not demonstrated in a small, randomized, double-blind, placebo-controlled withdrawal trial in very young pediatric patients (aged 1 to 24 months). SUBVENITE was associated with an increased risk for infections adverse reactions (SUBVENITE 37%, Placebo 5%), and respiratory adverse reaction (SUBVENITE 20%, Placebo 5%). Infections adverse reactions included bronchitis, bronchitis, ear infection, eye infection, otitis externa, pharyngitis, urinary tract infection, and viral infection. Respiratory adverse reactions included nasal congestion, cough, and apnea.

**Bipolar Disorder**

Safety and efficacy of SUBVENITE for the maintenance treatment of bipolar disorder were not established in a double-blind, randomized withdrawal, placebo-controlled trial that evaluated 301 pediatric patients aged 10 to 17 years with a current manic/hypomanic, depressed, or mixed mood episode as defined by DSM-IV-TR. In the randomized phase of the trial, adverse reactions that occurred in at least 5% of patients taking placebo (n = 86) were influenza (SUBVENITE 8%, placebo 2%), oropharyngeal pain (SUBVENITE 8%, placebo 2%), vomiting (SUBVENITE 6%, placebo 2%), upper abdominal pain (SUBVENITE 5%, placebo 1%), and suicidal ideation (SUBVENITE 5%, placebo 0%).

**Juvenile Animal Data**

In a juvenile animal study in which lamotrigine (oral doses of 0, 5, 15, or 30 mg/kg) was administered to young rats from postnatal day 7 to 62, decreased viability and growth were seen at the highest dose tested and long-term neurobehavioral abnormalities (decreased locomotor activity, increased reactivity, and learning deficits in animals tested as adults) were observed at the 2 highest doses. The no-effect dose for adverse developmental effects in juvenile animals is less than the human dose of 400 mg/day on a mg/m<sup>2</sup> basis.

**8.5 Geriatric Use**

Clinical trials of SUBVENITE for epilepsy and bipolar disorder did not include sufficient numbers of patients aged 65 years and older to determine whether they respond differently from younger patients or exhibit a different safety profile than that of younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

**8.6 Hepatic Impairment**

Experience in patients with hepatic impairment is limited. Based on a clinical pharmacology study in 24

subjects with mild, moderate, and severe liver impairment [see Clinical Pharmacology (12.3)], the following general recommendations can be made. No dosage adjustment is needed in patients with mild liver impairment. Initial, escalation, and maintenance doses should generally be reduced by approximately 25% in patients with moderate and severe liver impairment without ascites and 50% in patients with severe liver impairment with ascites. Escalation and maintenance doses may be adjusted according to clinical response [see Dosage and Administration (2.1)].

### 8.7 Renal Impairment

Lamotrigine is metabolized mainly by glucuronic acid conjugation, with the majority of the metabolites being recovered in the urine. In a small study comparing a single dose of lamotrigine in subjects with varying degrees of renal impairment with healthy volunteers, the plasma half-life of lamotrigine was approximately twice as long in the subjects with chronic renal failure [see Clinical Pharmacology (12.3)].

Initial doses of SUBVENTE should be based on patients' AED regimens; reduced maintenance doses may be effective for patients with significant renal impairment. Few patients with severe renal impairment have been evaluated during chronic treatment with lamotrigine. Because there is inadequate experience in this population, SUBVENTE should be used with caution in these patients [see Dosage and Administration (2.1)].

## 10 OVERDOSAGE

### 10.1 Human Overdose Experience


Overdoses involving quantities up to 15 g have been reported for SUBVENTE, some of which have been fatal. Overdose has resulted in ataxia, myoclonus, seizures (including tonic-clonic seizures), decreased level of consciousness, coma, and intraventricular conduction delay.

### 10.2 Management of Overdose

There are no specific antidotes for lamotrigine. Following a suspected overdose, hospitalization of the patient is advised. General supportive care is indicated, including frequent monitoring of vital signs and close observation of the patient. If indicated, emesis should be induced, and precautions should be taken to protect the airway. It should be kept in mind that immediate-release lamotrigine is rapidly absorbed [see Clinical Pharmacology (12.2)]. It is uncertain whether hemodialysis is an effective means of removing lamotrigine from the blood. In 6 renal failure patients, about 20% of the amount of lamotrigine in the body was removed by hemodialysis during a 4-hour session. A Poison Control Center should be contacted for information on the management of overdose of SUBVENTE.

## 11 DESCRIPTION

SUBVENTE is an AED of the phenyltriazine class, is chemically unrelated to existing AEDs. Lamotrigine's chemical name is 3,5-diamino-6-(2,3-dichlorophenyl)-pyrimidine; its molecular formula is  $C_{10}H_{10}N_4Cl_2$  and its molecular weight is 256.09. Lamotrigine, USP is a white to pale cream-colored powder and has a pK<sub>a</sub> of 5.7. Lamotrigine, USP is very slightly soluble in water (0.17 mg/mL at 25°C) and slightly soluble in 0.1 M HCl (4.1 mg/mL at 25°C). The structural formula is:



SUBVENTE (lamotrigine) tablets, USP are supplied for oral administration as 25-mg (white to off white), 100-mg (white to off white), 150-mg (white to off white), and 200-mg (white to off white) tablets. Each tablet contains the labeled amount of lamotrigine, USP and the following inactive ingredients: lactose monohydrate; magnesium stearate; microcrystalline cellulose; polyvidone; and sodium starch glycolate.

### Meets USP Dissolution Test 3

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

The precise mechanism(s) by which lamotrigine exerts its anticonvulsant action are unknown. In animal models designed to detect anticonvulsant activity, lamotrigine was effective in preventing seizure spread in the maximum electroshock (MES) and pentylenetetrazol (scMet) tests, and prevented seizures in the visually and electrically evoked after-discharge (EAD) tests for antiepileptic activity. Lamotrigine also displayed inhibitory properties in the kindling model in rats both during kindling development and in the fully kindled state. The relevance of these models to human epilepsy, however, is not known.

One proposed mechanism of action of lamotrigine, the relevance of which remains to be established in humans, involves an effect on sodium channels. *In vitro* pharmacological studies suggest that lamotrigine inhibits voltage-sensitive sodium channels, thereby stabilizing neuronal membranes and consequently modulating presynaptic transmitter release of excitatory amino acids (e.g., glutamate and aspartate).

#### Effect of Lamotrigine on N-Methyl-D-Aspartate Receptor-Mediated Activity

Lamotrigine did not inhibit N-methyl-D-aspartate (NMDA)-induced depolarizations in rat cortical slices or NMDA-induced cyclic GMP formation in immature rat cerebellum, nor did lamotrigine displace compounds that are either competitive or noncompetitive ligands at this glutamate receptor complex (CNQX, CGS, TCHP). The IC<sub>50</sub> for lamotrigine effects on NMDA-induced currents (in the presence of 3 μM of glycine) in cultured hippocampal neurons exceeded 100 μM.

The mechanisms by which lamotrigine exerts its therapeutic action in bipolar disorder have not been established.

### 12.2 Pharmacodynamics

#### Folate Metabolism

*In vitro*, lamotrigine inhibited dihydrofolate reductase, the enzyme that catalyzes the reduction of dihydrofolate to tetrahydrofolate. Inhibition of this enzyme may interfere with the biosynthesis of nucleic acids and protein. When oral daily doses of lamotrigine were given to pregnant rats during organogenesis, fetal, placental, and maternal folate concentrations were reduced. Significantly reduced concentrations of folate are associated with teratogenesis [see Use in Specific Populations (8.1)]. Folate concentrations were also reduced in male rats given repeated oral doses of lamotrigine. Reduced concentrations were partially returned to normal when supplemented with folic acid.

#### Accumulation in Kidneys

Lamotrigine accumulated in the kidney of the male rat, causing chronic progressive nephrosis, necrosis, and mineralization. These findings are attributed to α<sub>2</sub>-microglobulin, a species- and sex-specific protein that has not been detected in humans or other animal species.

#### Melanin Binding

Lamotrigine binds to melanin-containing tissues, e.g., in the eye and pigmented skin. It has been found in the uveal tract up to 52 weeks after a single dose in rodents.

#### Cardiovascular

In dogs, lamotrigine is extensively metabolized to a 2-N-methyl metabolite. This metabolite causes dose-dependent prolongation of the PR interval, widening of the QRS complex, and, at higher doses, complete AV conduction block. Similar cardiovascular effects are not anticipated in humans because only trace amounts of the 2-N-methyl metabolite (<0.6% of lamotrigine dose) have been found in human urine [see Clinical Pharmacology (12.2)]. However, it is conceivable that plasma concentrations of this metabolite could be increased in patients with a reduced capacity to glucuronidate lamotrigine (e.g., in patients with liver disease, patients taking concomitant medications that inhibit glucuronidation).

### 12.3 Pharmacokinetics

The pharmacokinetics of lamotrigine have been studied in subjects with epilepsy, healthy young and elderly volunteers, and volunteers with chronic renal failure. Lamotrigine pharmacokinetic parameters for adult and pediatric subjects and healthy normal volunteers are summarized in Tables 14 and 16.

Table 14. Mean Pharmacokinetic Parameters\* in Healthy Volunteers and Adult Subjects with Epilepsy

Adult Study Population	Number of Subjects	T <sub>max</sub> <sup>b</sup> , Time of Maximum Plasma Concentration (h)	t <sub>1/2</sub> <sup>c</sup> , Elimination Half-life (h)	CL/F: Apparent Plasma Clearance (mL/min/kg)
Healthy volunteers taking no other medications: Single-dose SUBVENTE	179	2.2 (0.25 to 12.0)	32.8 (14.0 to 103.0)	0.44 (0.12 to 1.10)
	36	1.7 (0.5 to 4.0)	25.4 (11.6 to 61.6)	0.50 (0.24 to 1.15)
Healthy volunteers taking valproate: Single-dose SUBVENTE	6	1.8 (1.0 to 4.0)	48.3 (31.5 to 88.6)	0.30 (0.14 to 0.42)
	18	1.5 (0.5 to 3.5)	70.3 (41.9 to 113.5)	0.18 (0.12 to 0.33)
Subjects with epilepsy taking valproate only: Single-dose SUBVENTE	4	4.8 (1.8 to 8.4)	58.8 (30.5 to 88.8)	0.28 (0.16 to 0.40)
Subjects with epilepsy taking carbamazepine, phenytoin, phenobarbital, or primidone plus valproate: Single-dose SUBVENTE	25	3.8 (1.0 to 10.0)	27.2 (11.2 to 51.6)	0.53 (0.27 to 1.04)
Subjects with epilepsy taking carbamazepine, phenytoin, phenobarbital, or primidone <sup>d</sup> : Single-dose SUBVENTE	24	2.3 (0.5 to 5.0)	14.4 (6.4 to 30.4)	1.10 (0.51 to 2.22)
	17	2.0 (0.75 to 5.93)	12.6 (7.5 to 23.1)	1.21 (0.66 to 1.82)

\*The majority of parameter means determined in each study had coefficients of variation between 20% and 40% for half-life and CL/F and between 30% and 70% for t<sub>max</sub>. The overall mean values were calculated from individual study means that were weighted based on the number of volunteers/subjects in each study. The numbers in parentheses below each parameter mean represent the range of individual volunteer/subject values across studies.

<sup>b</sup>Carbamazepine, phenytoin, phenobarbital, and primidone have been shown to increase the apparent clearance of lamotrigine. Enzyme-inducing oral contraceptives and other drugs, such as rifampin and protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir, that induce lamotrigine glucuronidation have also been shown to increase the apparent clearance of lamotrigine [see Drug Interactions (7)].

### Absorption

Lamotrigine is rapidly and completely absorbed after oral administration with negligible first-pass metabolism (absolute bioavailability is 98%). The bioavailability is not affected by food. Peak plasma concentrations occur anywhere from 1.8 to 4.8 hours following drug administration.

#### Dose Proportionality

In healthy volunteers not receiving any other medications and given single doses, the plasma concentrations of lamotrigine increased in direct proportion to the dose administered over the range of 50 to 400 mg. In 2 small studies (n=7 and 8) of patients with epilepsy who were maintained on other AEDs, there also was a linear relationship between dose and lamotrigine plasma concentrations at steady state following doses of 50 to 350 mg twice daily.

#### Distribution

Estimates of the mean apparent volume of distribution (VdF) of lamotrigine following oral administration ranged from 0.9 to 1.3 L/kg. VdF is independent of dose and is similar following single and multiple doses in both patients with epilepsy and in healthy volunteers.

#### Protein Binding

Data from *in vitro* studies indicate that lamotrigine is approximately 55% bound to human plasma proteins at plasma lamotrigine concentrations from 1 to 10 mg/mL (10 mg/mL is 4 to 6 times the trough plasma concentration observed in the controlled efficacy trials). Because lamotrigine is not highly bound to plasma proteins, clinically significant interactions with other drugs through competition for protein binding sites are unlikely. The binding of lamotrigine to plasma protein did not change in the presence of therapeutic concentrations of phenytoin, phenobarbital, or valproate. Lamotrigine did not displace other AEDs (carbamazepine, phenytoin, phenobarbital) from protein-binding sites.

#### Metabolism

Lamotrigine is metabolized predominantly by glucuronic acid conjugation; the major metabolite is an inactive 2-N-glucuronide conjugate. After oral administration of 240 mg of [<sup>14</sup>C]-lamotrigine (15 μCi) to 6 healthy volunteers, 94% was recovered in the urine and 2% was recovered in the feces. The radioactivity in the urine consisted of unchanged lamotrigine (10%), the 2-N-glucuronide (70%), 5-N-glucuronide (10%), 2-N-methyl metabolite (0.14%), and other unidentified minor metabolites (4%).

#### Enzyme Induction

The effects of lamotrigine on the induction of specific families of mixed-function oxidase isozymes have not been systematically evaluated.

Following multiple administration (150 mg twice daily) to normal volunteers taking no other medication, lamotrigine induced its own metabolism, resulting in a 25% decrease in t<sub>1/2</sub> and a 37%



increase in CL<sub>T</sub> at steady state compared with values obtained in the same volunteers following a single dose. Evidence gathered from other sources suggests that self-induction by lamotrigine may not occur when lamotrigine is given as adjunctive therapy in patients receiving enzyme-inducing drugs such as carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir that induce lamotrigine glucuronidation [see Drug Interactions (7)].

#### Elimination

The elimination half-life and apparent clearance of SUBVENTE following oral administration of lamotrigine to adult subjects with epilepsy and healthy volunteers is summarized in Table 14. Half-life and apparent oral clearance vary depending on concomitant AEDs.

#### Drug Interactions

The apparent clearance of lamotrigine is affected by the coadministration of certain medication [see Warnings and Precautions (5.8, 5.12), Drug Interactions (7)].

The net effects of drug interaction with lamotrigine are summarized in Tables 13 and 15, followed by details of the drug interaction studies below.

Table 15. Summary of Drug Interactions with Lamotrigine

Drug	Drug Plasma Concentration with Adjunctive Lamotrigine	Lamotrigine Plasma Concentration with Adjunctive Drugs
Oral contraceptives (e.g., ethinyl diethyl levonorgestrel) <sup>a</sup>	-- <sup>d</sup>	↓
Aripiprazole	Not assessed	-- <sup>e</sup>
Atazanavir/ritonavir	-- <sup>f</sup>	↓
Bupropion	Not assessed	↓
Carbamazepine	--	↓
Carbamazepine epoxide <sup>g</sup>	?	↓
Felbamate	Not assessed	--
Gabapentin	Not assessed	--
Lacosamide	Not assessed	--
Levetiracetam	--	--
Lithium	--	Not assessed
Lopinavir/ritonavir	-- <sup>h</sup>	↓
Olanzapine	--	-- <sup>h</sup>
Oxcarbazepine	--	--
10-Menthylhydroxy oxcarbazepine metabolite <sup>i</sup>	--	--
Perampanel	Not assessed	--
Phenobarbital/primidone	--	↓
Phenytoin	--	↓
Pregabalin	--	↓
Rifampin	Not assessed	↓
Risperidone	--	Not assessed
β-Hydroxyrisperidone <sup>j</sup>	--	↓
Rufinamide	--	↓
Valproate	↓	↑
Valproate + phenytoin and/or lamotrigine	Not assessed	--
Zonisamide	Not assessed	--

<sup>a</sup> From adjunctive clinical trial and volunteer trials.

<sup>b</sup> Net effects were estimated by comparing the mean clearance values obtained in adjunctive clinical trials and volunteer trials.

<sup>c</sup> The effect of other hormonal contraceptive preparations or hormone replacement therapy on the pharmacokinetics of lamotrigine has not been systematically evaluated in clinical trials, although the effect may be similar to that seen with the ethinyl diethyl levonorgestrel combination.

<sup>d</sup> Slight decrease in levetiracetam.

<sup>e</sup> Slight decrease, not expected to be clinically meaningful.

<sup>f</sup> Compared with historical controls.

<sup>g</sup> Not administered, but an active metabolite of carbamazepine.

<sup>h</sup> Not administered, but an active metabolite of oxcarbazepine.

<sup>i</sup> Not administered, but an active metabolite of oxcarbazepine.

<sup>j</sup> Slight increase, not expected to be clinically meaningful.

-- = No significant effect.

? = Conflicting data.

#### Estrogen-Containing Oral Contraceptives

In 16 female volunteers, an oral contraceptive preparation containing 30 mcg ethinylestradiol and 150 mcg levonorgestrel increased the apparent clearance of lamotrigine (300 mg/day) by approximately 2-fold with mean decreases in AUC of 52% and in C<sub>max</sub> of 30%. In this study, trough serum lamotrigine concentrations gradually increased and were approximately 2-fold higher on average at the end of the week of the inactive hormone preparation compared with trough lamotrigine concentrations at the end of the active hormone cycle.

Gradual transient increases in lamotrigine plasma levels (approximately 2-fold increase) occurred during the week of inactive hormone preparation (pill-free weeks) for women not also taking a drug that increased the clearance of lamotrigine (carbamazepine, phenytoin, phenobarbital, primidone, or other drugs such as rifampin and the protease inhibitors lopinavir/ritonavir and atazanavir/ritonavir that induce lamotrigine glucuronidation) [see Drug Interactions (7)]. The increase in lamotrigine plasma levels will be greater if the dose of SUBVENTE is increased the few days before or during the pill-free week. Increases in lamotrigine plasma levels could result in dose-dependent adverse reactions.

In the same study, coadministration of lamotrigine (300 mg/day) in 16 female volunteers did not affect the pharmacokinetics of the ethinylestradiol component of the oral contraceptive preparation. There were mean decreases in the AUC and C<sub>max</sub> of the levonorgestrel component of 19% and 12%, respectively. Measurement of serum progesterone indicated that there was no hormonal evidence of ovulation in any of the 16 volunteers, although measurement of serum FSH, LH, and estradiol indicated that there was some loss of suppression of the hypothalamic-pituitary-ovarian axis.

The effects of doses of lamotrigine other than 300 mg/day have not been systematically evaluated in controlled clinical trials.

The clinical significance of the observed hormonal changes on ovulatory activity is unknown. However, the possibility of decreased contraceptive efficacy in some patients cannot be excluded. Therefore, patients should be instructed to promptly report changes in their menstrual pattern (e.g., break-through bleeding).

Dosage adjustments may be necessary for women receiving estrogen-containing oral contraceptive preparation [see Dosage and Administration (2.1)].

#### Other Hormonal Contraceptives or Hormone Replacement Therapy

The effect of other hormonal contraceptive preparations or hormone replacement therapy on the pharmacokinetics of lamotrigine has not been systematically evaluated. It has been reported that ethinylestradiol, not progestogens, increased the clearance of lamotrigine up to 2-fold, and the progestin-only pills had no effect on lamotrigine plasma levels. Therefore, adjustments to the dosage of SUBVENTE in the presence of progestin alone will likely not be needed.

#### Aripiprazole

In 18 patients with bipolar disorder on a stable regimen of 100 to 400 mg/day of lamotrigine, the lamotrigine AUC and C<sub>max</sub> were reduced by approximately 10% in patients who received aripiprazole 10 to 30 mg/day for 7 days, followed by 30 mg/day for an additional 7 days. This reduction in lamotrigine exposure is not considered clinically meaningful.

#### Atazanavir/Ritonavir

In a study in healthy volunteers, daily doses of atazanavir/ritonavir (300 mg/100 mg) reduced the plasma AUC and C<sub>max</sub> of lamotrigine (single 100-mg dose) by an average of 32% and 6%, respectively, and shortened the elimination half-lives by 27%. In the presence of atazanavir/ritonavir (300 mg/100 mg), the metabolite-to-lamotrigine ratio was increased from 0.45 to 0.71 consistent with induction of glucuronidation. The pharmacokinetics of atazanavir/ritonavir were similar in the presence of concomitant lamotrigine to the historical data of the pharmacokinetics in the absence of lamotrigine.

#### Bupropion

The pharmacokinetics of a 100-mg single dose of lamotrigine in healthy volunteers (n = 12) were not changed by coadministration of bupropion sustained-release formulation (150 mg twice daily) starting 11 days before lamotrigine.

#### Carbamazepine

Lamotrigine has no appreciable effect on steady-state carbamazepine plasma concentration. Limited clinical data suggest there is a higher incidence of dizziness, diplopia, ataxia, and blurred vision in patients receiving carbamazepine with lamotrigine than in patients receiving other AEDs with lamotrigine [see Adverse Reactions (6.1)]. The mechanism of this interaction is unclear. The effect of lamotrigine on plasma concentrations of carbamazepine-epoxide is unclear. In a small subset of patients (n = 7) studied in a placebo-controlled trial, lamotrigine had no effect on carbamazepine-epoxide plasma concentrations, but in a small, uncontrolled study (n = 9), carbamazepine-epoxide levels increased. The addition of carbamazepine decreases lamotrigine steady-state concentrations by approximately 40%.

#### Felbamate

In a trial in 21 healthy volunteers, coadministration of felbamate (1200 mg twice daily) with lamotrigine (100 mg twice daily for 10 days) appeared to have no clinically relevant effects on the pharmacokinetics of lamotrigine.

#### Folate Inhibitors

Lamotrigine is a weak inhibitor of dihydrofolate reductase. Prescribers should be aware of this action when prescribing other medications that inhibit folate metabolism.

#### Lacosamide

Plasma concentrations of lamotrigine were not affected by concomitant lacosamide (200, 400, or 600 mg/day) in placebo-controlled clinical trials in patients with partial-onset seizures.

#### Gabapentin

Based on a retrospective analysis of plasma levels in 34 subjects who received lamotrigine both with and without gabapentin, gabapentin does not appear to change the apparent clearance of lamotrigine.

#### Levetiracetam

Potential drug interaction between levetiracetam and lamotrigine were assessed by evaluating serum concentrations of both agents during placebo-controlled clinical trials. These data indicate that lamotrigine does not influence the pharmacokinetics of levetiracetam and that levetiracetam does not influence the pharmacokinetics of lamotrigine.

#### Lithium

The pharmacokinetics of lithium were not altered in healthy subjects (n = 20) by coadministration of lamotrigine (100 mg/day) for 6 days.

#### Lopinavir/Ritonavir

The addition of lopinavir (400 mg twice daily)/ritonavir (100 mg twice daily) decreased the AUC, C<sub>max</sub>, and elimination half-life of lamotrigine by approximately 50% to 55.4% in 18 healthy subjects. The pharmacokinetics of lopinavir/ritonavir were similar with concomitant lamotrigine, compared with that in historical controls.

#### Olanzapine

The AUC and C<sub>max</sub> of olanzapine were similar following the addition of olanzapine (15 mg once daily) to lamotrigine (200 mg once daily) in healthy male volunteers (n = 16) compared with the AUC and C<sub>max</sub> in healthy male volunteers receiving olanzapine alone (n = 16).

In the same trial, the AUC and C<sub>max</sub> of lamotrigine were reduced on average by 24% and 20%, respectively, following the addition of olanzapine to lamotrigine in healthy male volunteers compared with those receiving lamotrigine alone. This reduction in lamotrigine plasma concentrations is not expected to be clinically meaningful.

#### Oxcarbazepine

The AUC and C<sub>max</sub> of oxcarbazepine and its active 10-menthylhydroxy oxcarbazepine metabolite were not significantly different following the addition of oxcarbazepine (600 mg twice daily) to lamotrigine (200 mg once daily) in healthy male volunteers (n = 13) compared with healthy male volunteers receiving oxcarbazepine alone (n = 13).

In the same trial, the AUC and C<sub>max</sub> of lamotrigine were similar following the addition of oxcarbazepine (600 mg twice daily) to lamotrigine in healthy male volunteers compared with those receiving lamotrigine alone. Limited clinical data suggest a higher incidence of headache, dizziness, nausea, and somnolence with coadministration of lamotrigine and oxcarbazepine compared with lamotrigine alone or oxcarbazepine alone.

#### Perampanel

In a pooled analysis of data from 3 placebo-controlled clinical trials investigating adjunctive perampanel in patients with partial-onset and primary generalized tonic-clonic seizures, the highest perampanel dose evaluated (12 mg/day) increased lamotrigine clearance by <10%. An effect of this magnitude is not considered to be clinically relevant.

#### Phenobarbital/Primidone

The addition of phenobarbital or primidone decreases lamotrigine steady-state concentrations by approximately 40%.

#### Phenytoin

Lamotrigine has no appreciable effect on steady-state phenytoin plasma concentrations in patients with

epilepsy. The addition of phenytoin decreases lamotrigine steady-state concentration by approximately 40%.

#### Pregabalin

Steady-state trough plasma concentrations of lamotrigine were not affected by concomitant pregabalin (200 mg 3 times daily) administration. There are no pharmacokinetic interactions between lamotrigine and pregabalin.

#### Rifampin

In 10 male volunteers, rifampin (600 mg/day for 5 days) significantly increased the apparent clearance of a single 25-mg dose of lamotrigine by approximately 2-fold (AUC decreased by approximately 40%).

#### Risperidone

In a 14 healthy volunteers study, multiple oral doses of lamotrigine 400 mg daily had no clinically significant effect on the single-dose pharmacokinetics of risperidone 2 mg and its active metabolite 9-OH risperidone. Following the coadministration of risperidone 2 mg with lamotrigine, 12 of the 14 volunteers reported somnolence compared with 1 out of 20 when risperidone was given alone, and none when lamotrigine was administered alone.

#### Topiramate

Topiramate resulted in no change in plasma concentration of lamotrigine. Administration of lamotrigine resulted in a 15% increase in topiramate concentration.

#### Valproate

When lamotrigine was administered to healthy volunteers (n = 18) receiving valproate, the trough steady-state valproate plasma concentrations decreased by an average of 25% over a 3-week period, and then stabilized. However, adding lamotrigine to the existing therapy did not cause a change in valproate plasma concentration in either adult or pediatric patients in controlled clinical trials. The addition of valproate increased lamotrigine steady-state concentrations in normal volunteers by slightly more than 2-fold. In 1 trial, maximal inhibition of lamotrigine clearance was reached at valproate doses between 250 and 500 mg/day and did not increase as the valproate dose was further increased.

#### Zonisamide

In a study in 18 patients with epilepsy, coadministration of zonisamide (200 to 400 mg/day) with lamotrigine (150 to 500 mg/day for 35 days) had no significant effect on the pharmacokinetics of lamotrigine.

#### Known Inhibitors or Inhibitors of Glucuronidation

Drugs other than those listed above have not been systematically evaluated in combination with lamotrigine. Since lamotrigine is metabolized predominantly by glucuronic acid conjugation, drugs that are known to induce or inhibit glucuronidation may affect the apparent clearance of lamotrigine and doses of lamotrigine may require adjustment based on clinical response.

#### Other

In vitro assessment of the inhibitory effect of lamotrigine at OCT2 demonstrate that lamotrigine, but not the N(2)-glucuronide metabolite, is an inhibitor of OCT2 at a potentially clinically relevant concentration, with IC<sub>50</sub> values of 53.8 μM [see Drug Interactions (7)].

Results of in vitro experiments suggest that clearance of lamotrigine is unlikely to be reduced by concomitant administration of antiepileptic, clonazepam, clozapine, fluoxetine, haloperidol, lorazepam, phenelzine, sertraline, or zolpidem.

Results of in vitro experiments suggest that lamotrigine does not reduce the clearance of drugs eliminated predominantly by CYP2D6.

#### Specific Population

**Patients with Renal Impairment:** Twelve volunteers with chronic renal failure (mean creatinine clearance = 11 mL/min; range: 6 to 23) and another 6 individuals undergoing hemodialysis were each given a single 100-mg dose of lamotrigine. The mean plasma half-lives determined in the study were 42.9 hours (chronic renal failure), 13.0 hours (during hemodialysis), and 57.4 hours (between hemodialysis) compared with 26.2 hours in healthy volunteers. On average, approximately 20% (range: 5.6 to 35.1) of the amount of lamotrigine present in the body was eliminated by hemodialysis during a 4-hour session [see Dosage and Administration (2.1)].

**Patients with Hepatic Impairment:** The pharmacokinetics of lamotrigine following a single 100-mg dose of lamotrigine were evaluated in 24 subjects with mild, moderate, and severe hepatic impairment (Child-Pugh classification system) and compared with 12 subjects without hepatic impairment. The subjects with severe hepatic impairment were without ascites (n = 2) or with ascites (n = 5). The mean apparent clearance of lamotrigine in subjects with mild (n = 12), moderate (n = 5), severe without ascites (n = 2), and severe with ascites (n = 3) liver impairment were 0.30, 0.07, 0.24, 0.15, 0.21 ± 0.04, and 0.15 ± 0.09 mL/min/kg, respectively, as compared with 0.37 ± 0.1 mL/min/kg in the healthy controls. Mean half-lives of lamotrigine in subjects with mild, moderate, severe without ascites, and severe with ascites hepatic impairment were 46 ± 20, 72 ± 44, 67 ± 11, and 109 ± 48 hours, respectively, as compared with 33 ± 7 hours in healthy controls [see Dosage and Administration (2.1)].

**Pediatric Patients:** The pharmacokinetics of lamotrigine following a single 2-mg/kg dose were evaluated in 2 studies in pediatric subjects (n = 29 for subjects aged 10 months to 5.9 years and n = 26 for subjects aged 5 to 11 years). Forty-three subjects received concomitant therapy with other AEDs and 12 subjects received lamotrigine as monotherapy. Lamotrigine pharmacokinetic parameters for pediatric patients are summarized in Table 16.

Population pharmacokinetic analysis involving subjects aged 2 to 18 years demonstrated that lamotrigine clearance was influenced predominantly by total body weight and concurrent AED therapy. The oral clearance of lamotrigine was higher, on a body weight basis, in pediatric patients than in adults. Weight-normalized lamotrigine clearance was higher in those subjects weighing <30 kg compared with those weighing ≥30 kg. Accordingly, patients weighing <30 kg may need an increase of as much as 50% in maintenance doses, based on clinical response, as compared with subjects weighing more than 30 kg being administered the same AEDs [see Dosage and Administration (2.2)]. These analyses also revealed that, after accounting for body weight, lamotrigine clearance was not significantly influenced by age. Thus, the same weight-adjusted doses should be administered to children irrespective of differences in age. Concomitant AEDs which influence lamotrigine clearance in adults were found to have similar effects in children.

Table 16. Mean Pharmacokinetic Parameters in Pediatric Subjects with Epilepsy

Pediatric Study Population	Number of Subjects	T <sub>max</sub> (h)	t <sub>1/2</sub> (h)	CL/F (mL·min/kg)
<b>Age 10 months to 5.3 years</b> Subjects taking carbamazepine, phenytoin, phenobarbital, or primidone <sup>a</sup>	10	3.0 (1.0 to 5.9)	7.7 (5.7 to 11.4)	3.62 (2.44 to 5.28)
Subjects taking antiepileptic drugs with no known effect on the apparent clearance of lamotrigine. Subjects taking valproate only	7	5.2 (2.9 to 6.1)	19.0 (12.9 to 27.1)	1.2 (0.75 to 2.42)
<b>Age 5 to 11 years</b> Subjects taking carbamazepine, phenytoin, phenobarbital, or primidone <sup>a</sup>	8	7	16.0 (10 to 3.0) <sup>b</sup>	2.54 (1.35 to 5.58)
Subjects taking carbamazepine, phenytoin, phenobarbital, or primidone <sup>a</sup> plus valproate	8	3.3 (1.0 to 6.4)	19.1 (7.0 to 31.2)	0.89 (0.39 to 1.93)
<b>Age 13 to 18 years</b> Subjects taking carbamazepine, phenytoin, phenobarbital, or primidone <sup>a</sup>	3	4.5 (1.0 to 6.0)	65.8 (60.7 to 73.7)	0.24 (0.21 to 0.26)
Subjects taking valproate only	4	— <sup>c</sup>	— <sup>c</sup>	1.3
		— <sup>c</sup>	— <sup>c</sup>	0.3

<sup>a</sup> Carbamazepine, phenytoin, phenobarbital, and primidone have been shown to increase the apparent clearance of lamotrigine. Enzyme-containing oral contraceptives, ethanol, and the protein inhibitor leprosin (bromelain) also have been shown to increase the apparent clearance of lamotrigine [see Drug Interactions (7)].

<sup>b</sup> Two subjects were included in the calculation for mean T<sub>max</sub>.

<sup>c</sup> Parameter not estimated.

**Geriatric Patients:** The pharmacokinetics of lamotrigine following a single 150-mg dose of lamotrigine were evaluated in 12 elderly volunteers between the ages of 65 and 76 years (mean creatinine clearance = 61 mL/min; range: 33 to 108 mL/min). The mean half-life of lamotrigine in these subjects was 31.2 hours (range: 24.5 to 43.4 hours), and the mean clearance was 0.40 mL/min/kg (range: 0.20 to 0.48 mL/min/kg).

**Male and Female Patients:** The clearance of lamotrigine is not affected by gender. However, during dose escalation of lamotrigine in 1 clinical trial in patients with epilepsy on a stable dose of valproate (n = 77), mean trough lamotrigine concentrations unadjusted for weight were 24% to 40% higher (0.3 to 1.7 mg/mL) in females than in males.

**Racial or Ethnic Groups:** The apparent oral clearance of lamotrigine was 25% lower in non-Caucasian than Caucasian.

### 13 NONCLINICAL TOXICOLOGY

#### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No evidence of carcinogenicity was seen in mouse or rat following oral administration of lamotrigine for up to 2 years at doses up to 30 mg/kg/day and 10 to 15 mg/kg/day in mouse and rat, respectively. The highest doses tested are less than the human dose of 400 mg/day on a body surface area (mg/m<sup>2</sup>) basis.

Lamotrigine was negative in in vitro gene mutation (Ames and mouse lymphoma 4) assays and in clastogenicity (in vitro human lymphocyte and in vivo rat bone marrow) assays.

No evidence of impaired fertility was detected in rats given oral doses of lamotrigine up to 20 mg/kg/day. The highest dose tested is less than the human dose of 400 mg/day on a mg/m<sup>2</sup> basis.

### 14 CLINICAL STUDIES

#### 14.1 Epilepsy

##### Monotherapy with SUBVENITE in Adults with Partial-Onset Seizures Already Receiving Treatment with Carbamazepine, Phenytoin, Phenobarbital, or Primidone as the Single Antiepileptic Drug

The effectiveness of monotherapy with SUBVENITE was established in a multicenter double-blind clinical trial enrolling 156 adult outpatients with partial-onset seizures. The patients experienced at least 4 simple partial-onset, complex partial-onset, and/or secondarily generalized seizures during each of 2 consecutive 4-week periods while receiving carbamazepine or phenytoin monotherapy during baseline. SUBVENITE (target dose of 500 mg/day) or valproate (1,000 mg/day) was added to either carbamazepine or phenytoin monotherapy over a 4-week period. Patients were then converted to monotherapy with SUBVENITE or valproate during the next 4 weeks, then continued on monotherapy for an additional 12-week period.

Trial endpoints were completion of all weeks of trial treatment or meeting an escape criterion. Criteria for escape relative to baseline were: (1) doubling of average monthly seizure count, (2) doubling of highest consecutive 2-day seizure frequency, (3) emergence of a new seizure type (defined as a seizure that did not occur during the 8-week baseline) that is more severe than seizure types that occur during study treatment, or (4) clinically significant prolongation of generalized tonic-clonic seizures. The primary efficacy variable was the proportion of patients in each treatment group who met escape criteria.

The percentages of patients who met escape criteria were 42% (32/76) in the group receiving SUBVENITE and 69% (52/80) in the valproate group. The difference in the percentage of patients meeting escape criteria was statistically significant (P = 0.0012) in favor of SUBVENITE. No differences in efficacy-based metrics, sex, or race were detected.

Patients in the control group were intentionally treated with a relatively low dose of valproate; as such, the sole objective of this trial was to demonstrate the effectiveness and safety of monotherapy with SUBVENITE, and cannot be interpreted to imply the superiority of SUBVENITE to an adequate dose of valproate.

##### Adjunctive Therapy with Lamotrigine in Adults with Partial-Onset Seizures

The effectiveness of SUBVENITE as adjunctive therapy (added to other AEDs) was initially established in 3 pivotal, multicenter, placebo-controlled, double-blind clinical trials in 355 adults with refractory partial-onset seizures. The patients had a history of at least 4 partial-onset seizures per month in spite of receiving 1 or more AEDs at therapeutic concentrations and in 2 of the trials were observed on their established AED regimen during baselines that varied between 8 to 12 weeks. In the third trial, patients were not observed in a prospective baseline. In patients continuing to have at least 4 seizures per month during the baseline, SUBVENITE or placebo was then added to the existing therapy. In all 3 trials, change from baseline in seizure frequency was the primary measure of effectiveness. The results given below are for all partial-onset seizures in the intent-to-treat population (all patients who received at least 1 dose of treatment) in each trial, unless otherwise indicated. The median seizure frequency at baseline was 3 per week while the mean at baseline was 6.6 per week for all patients enrolled in efficacy trials.

One trial (n = 216) was a double-blind, placebo-controlled, parallel trial consisting of a 24-week treatment period. Patients could not be on more than 2 other anticonvulsants and valproate was not allowed. Patients were randomized to receive placebo, a target dose of 300 mg/day of SUBVENITE, or a target dose of 500 mg/day of SUBVENITE. The median reductions in the frequency of all partial-onset seizures relative to baseline were 8% in patients receiving placebo, 20% in patients receiving 300 mg/day of SUBVENITE, and 36% in patients receiving 500 mg/day of SUBVENITE. The seizure frequency reduction was statistically significant in the 500-mg/day group compared with the placebo group, but not in the 300-mg/day group.

A second trial (n = 95) was a double-blind, placebo-controlled, randomized, crossover trial consisting of two 14-week treatment periods (the last 2 weeks of which consisted of dose tapering) separated by a 4-week washout period. Patients could not be on more than 2 other anticonvulsants and valproate was not allowed. The target dose of SUBVENITE was 400 mg/day. When the first 12 weeks of the treatment

periods were analyzed, the median change in seizure frequency was a 25% reduction on SUBVENTE compared with placebo ( $P < 0.001$ ).

The third trial ( $n = 41$ ) was a double-blind, placebo-controlled, crossover trial consisting of two 12-week treatment periods separated by a 4-week washout period. Patients could not be on more than 2 other anticonvulsants. Thirteen patients were on concomitant valproate; these patients received 150 mg/day of SUBVENTE. The 28 other patients had a target dose of 300 mg/day of SUBVENTE. The median change in seizure frequency was a 26% reduction on SUBVENTE compared with placebo ( $P < 0.01$ ).

No differences in efficacy based on age, sex, or race, as measured by change in seizure frequency, were detected.

#### Adjunctive Therapy with SUBVENTE in Pediatric Patients with Partial-Onset Seizures

The effectiveness of SUBVENTE as adjunctive therapy in pediatric patients with partial-onset seizures was established in a multicenter, double-blind, placebo-controlled trial in 189 patients aged 2 to 16 years ( $n = 90$  on SUBVENTE,  $n = 101$  on placebo). Following an 8-week baseline phase, patients were randomized to 18 weeks of treatment with SUBVENTE or placebo added to their current AED regimen of up to 2 drugs. Patients were dosed based on body weight and valproate use. Target doses were designed to approximate 5 mg/kg/day for patients taking valproate (maximum dose: 200 mg/day) and 15 mg/kg/day for the patients not taking valproate (maximum dose: 750 mg/day). The primary efficacy endpoint was percentage change from baseline in all partial-onset seizures. For the intent-to-treat population, the median reduction of all partial-onset seizures was 36% in patients treated with SUBVENTE and 7% on placebo, a difference that was statistically significant ( $P < 0.01$ ).

#### Adjunctive Therapy with SUBVENTE in Pediatric and Adult Patients with Lennox-Gastaut Syndrome

The effectiveness of SUBVENTE as adjunctive therapy in patients with Lennox-Gastaut syndrome was established in a multicenter, double-blind, placebo-controlled trial in 169 patients aged 3 to 25 years ( $n = 79$  on SUBVENTE,  $n = 90$  on placebo). Following a 4-week, single-blind, placebo phase, patients were randomized to 16 weeks of treatment with SUBVENTE or placebo added to their current AED regimen of up to 3 drugs. Patients were dosed on a fixed-dose regimen based on body weight and valproate use. Target doses were designed to approximate 5 mg/kg/day for patients taking valproate (maximum dose: 200 mg/day) and 15 mg/kg/day for patients not taking valproate (maximum dose: 400 mg/day). The primary efficacy endpoint was percentage change from baseline in major motor seizures (aetia, tonic, major myoclonic, and tonic-clonic seizures). For the intent-to-treat population, the median reduction of major motor seizures was 12% in patients treated with SUBVENTE and 9% on placebo, a difference that was statistically significant ( $P < 0.05$ ). Drop attacks were significantly reduced by SUBVENTE (34%) compared with placebo (9%), as were tonic-clonic seizures (36% reduction versus 10% increase for SUBVENTE and placebo, respectively).

#### Adjunctive Therapy with SUBVENTE in Pediatric and Adult Patients with Primary Generalized Tonic-Clonic Seizures

The effectiveness of SUBVENTE as adjunctive therapy in patients with PGTC seizures was established in a multicenter, double-blind, placebo-controlled trial in 117 pediatric and adult patients aged 2 years and older ( $n = 58$  on SUBVENTE,  $n = 59$  on placebo). Patients with at least 3 PGTC seizures during an 8-week baseline phase were randomized to 19 to 24 weeks of treatment with SUBVENTE or placebo added to their current AED regimen of up to 2 drugs. Patients were dosed on a fixed-dose regimen, with target doses ranging from 3 to 12 mg/kg/day for pediatric patients and from 200 to 400 mg/day for adult patients based on concomitant AEDs.

The primary efficacy endpoint was percentage change from baseline in PGTC seizures. For the intent-to-treat population, the median percent reduction in PGTC seizures was 66% in patients treated with SUBVENTE and 34% on placebo, a difference that was statistically significant ( $P = 0.006$ ).

## 14.2 Bipolar Disorder

### Adults

The effectiveness of SUBVENTE in the maintenance treatment of bipolar I disorder was established in 2 multicenter, double-blind, placebo-controlled trials in adult patients (aged 18 to 82 years) who met DSM-IV criteria for bipolar I disorder. Trial 1 enrolled patients with a current or recent (within 60 days) depressive episode as defined by DSM-IV and Trial 2 included patients with a current or recent (within 60 days) episode of mania or hypomania as defined by DSM-IV. Both trials included a cohort of patients (30% of 404 subjects in Trial 1 and 28% of 171 patients in Trial 2) with rapid cycling bipolar disorder (4 to 6 episodes per year).

In both trials, patients were titrated to a target dose of 200 mg of SUBVENTE as add-on therapy or as monotherapy with gradual withdrawal of any psychotropic medications during an 8- to 16-week open-label period. Overall 81% of 1,305 patients participating in the open-label period were receiving 1 or more other psychotropic medications, including benzodiazepines, selective serotonin reuptake inhibitors (SSRIs), atypical antipsychotics (including olanzapine), valproate, or lithium, during titration of SUBVENTE. Patients with a CGI-severity score of 3 or less maintained for at least 4 continuous weeks, including at least the final week on monotherapy with SUBVENTE, were randomized to a placebo-controlled, double-blind treatment period for up to 18 months. The primary endpoint was TIME (time to intervention for a mood episode or one that was emerging, time to discontinuation for either an adverse event that was judged to be related to bipolar disorder, or for lack of efficacy). The mood episode could be depression, mania, hypomania, or a mixed episode.

In Trial 1, patients received double-blind monotherapy with SUBVENTE 50 mg/day ( $n = 50$ ), SUBVENTE 200 mg/day ( $n = 124$ ), SUBVENTE 400 mg/day ( $n = 47$ ), or placebo ( $n = 123$ ). SUBVENTE 200- and 400-mg/day treatment groups combined was superior to placebo in delaying the time to occurrence of a mood episode (Figure 1). Separate analyses of the 200- and 400-mg/day dose groups revealed no added benefit from the higher dose.

In Trial 2, patients received double-blind monotherapy with SUBVENTE (100 to 400 mg/day,  $n = 59$ ) or placebo ( $n = 70$ ). SUBVENTE was superior to placebo in delaying time to occurrence of a mood episode (Figure 2). The median dose of SUBVENTE was about 211 mg/day.

Although these trials were not designed to separately evaluate time to the occurrence of depression or mania, a combined analysis for the 2 trials revealed a statistically significant benefit for SUBVENTE over placebo in delaying the time to occurrence of both depression and mania, although the finding was more robust for depression.

Figure 1: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Mood Episode (Trial 1)





Figure 2: Kaplan-Meier Estimation of Cumulative Proportion of Patients with Mood Episode (Trial 2)



## 16 HOW SUPPLIED/STORAGE AND HANDLING

SUBVENTE (lamotrigine) tablets, USP 200 mg

White to off white, round shape, flat face beveled edge, uncoated tablets debossed with "20LA" on one side and break line on other side.

NDC: 70518-2645-00 100 in 1 BOX

NDC: 70518-2645-01 1 in 1 POUCH

### Storage

Store at 20° to 25° C (68° to 77° F); excursions permitted to 15° to 30° C (59° to 86° F) [See USP Controlled Room Temperature].

### Repackaging and Distribution By:

Remedy Repack, Inc.  
625 Kolter Dr. Suite #4 Indian, PA 1-724-465-6762

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

### Rash

Prior to initiation of treatment with SUBVENTE, inform patients that a rash or other signs or symptoms of hypersensitivity (e.g., fever, lymphadenopathy) may herald a serious medical event and instruct them to report any such occurrence to their healthcare providers immediately.

### Hemophagocytic Lymphohistiocytosis

Prior to initiation of treatment with SUBVENTE, inform patients that excessive immune activation may occur with SUBVENTE and that they should report signs or symptoms such as fever, rash, or lymphadenopathy to a healthcare provider immediately.

### Multifocal Hypersensitivity Reaction, Blood Dyscrasias, and Organ Failure

Inform patients that multifocal hypersensitivity reactions and acute multiorgan failure may occur with SUBVENTE. Isolated organ failure or isolated blood dyscrasias without evidence of multifocal hypersensitivity may also occur. Instruct patients to contact their healthcare providers immediately if they experience any signs or symptoms of these conditions [see Warnings and Precautions (3.3, 5.4)].

### Suicidal Thinking and Behavior

Inform patients, their caregivers, and families that AEDs, including SUBVENTE, may increase the risk of suicidal thoughts and behavior. Instruct them to be alert for the emergence or worsening of symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts or behavior or thoughts about self-harm. Instruct them to immediately report behaviors of concern to their healthcare providers.

### Worsening of Seizures

Instruct patients to notify their healthcare providers if worsening of seizure control occurs.

### Central Nervous System Adverse Effects

Inform patients that SUBVENTE may cause dizziness, somnolence, and other symptoms and signs of central nervous system depression. Accordingly, instruct them neither to drive a car nor to operate other complex machinery until they have gained sufficient experience on SUBVENTE to gauge whether or not it adversely affects their mental and/or motor performance.

### Pregnancy and Nursing

Instruct patients to notify their healthcare providers if they become pregnant or intend to become pregnant during therapy and if they intend to breastfeed or are breastfeeding an infant.

Encourage patients to enroll in the NAEAD Pregnancy Registry if they become pregnant. This registry is collecting information about the safety of antiepileptic drugs during pregnancy. To enroll, patients can call the toll-free number 1-888-233-2344 [see Use in Specific Populations (6.1)].

Inform patients who intend to breastfeed that SUBVENTE is present in breast milk and advise them to monitor their child for potential adverse effects of this drug. Discuss the benefits and risks of continuing breastfeeding.

### Oral Contraceptive Use

Instruct women to notify their healthcare providers if they plan to start or stop use of oral

contraceptives or other female hormonal preparations. Starting estrogen-containing oral contraceptives may significantly decrease lamotrigine plasma levels and stopping estrogen-containing oral contraceptives (including the pill-free week) may significantly increase lamotrigine plasma levels (see Warnings and Precautions (5.1), Clinical Pharmacology (12.3)). Also instruct women to promptly notify their healthcare providers if they experience adverse reactions or changes in menstrual pattern (e.g., break-through bleeding) while receiving SUBVENTE in combination with these medications.

**Discontinuing SUBVENTE**

Instruct patients to notify their healthcare providers if they stop taking SUBVENTE for any reason and not to resume SUBVENTE without consulting their healthcare providers.

**Asseptic Meningitis**

Inform patients that SUBVENTE may cause aseptic meningitis. Instruct them to notify their healthcare providers immediately if they develop signs and symptoms of meningitis such as headache, fever, nausea, vomiting, stiff neck, rash, abnormal sensitivity to light, myalgia, chills, confusion, or drowsiness while taking SUBVENTE.

**Parenteral Medication Errors**

To avoid a medication error of using the wrong drug or formulation, strongly advise patients to visually inspect their tablets to verify that they are lamotrigine, as well as the correct formulation of SUBVENTE, each time they fill their prescription (see Dosage Forms and Strengths (3.1, 3.2, 3.3), How Supplied/Storage and Handling (16)). Refer the patient to the Medication Guide that provides depictions of the SUBVENTE tablets.

**Repackaged and Distributed By:**

Remedy Repack, Inc.  
625 Kolter Dr. Suite #4 Indiana, PA 1-724-465-8762

**Repackaged and Distributed By:**

Remedy Repack, Inc.  
625 Kolter Dr. Suite #4 Indiana, PA 1-724-465-8762

**DRUG:**

NDC: 70518-2645-0



NDC: 70518-2645-1


PACKAGING: 1 in 1 POUCH

OUTER PACKAGING: 100 in 1 BOX

ACTIVE INGREDIENT(S):

INACTIVE INGREDIENT(S):

<b>Subventite</b>		NDC #: 70518-2645-00	
Lamotrigine		Expires:	
200 mg Tablet		LOT #: 69102-0320-01	
QTY: 100		Source NDC: 69102-0320-01	
		MFG: CMAP, West Chicago, IL 60185	
RX ONLY		MFG: CMAP, West Chicago, IL 60185	
Keep this and all medication out of the reach of children			
Directions For Use: See Package Insert			
Store at 20-25°C (68-77°F); excursions permitted to 15-30°C (59-86°F) (see USP Controlled Room Temperature). See USP Controlled Room Temperature. Repackaged by: RemedyRepack Inc., Indiana, PA 15109, 724-465-8762			

<b>Subventite</b>		NDC #: 70518-2645-01	
Lamotrigine		Expires:	
200 mg Tablet		LOT #: 69102-0320-01	
QTY: 1		Source NDC: 69102-0320-01	
		MFG: CMAP, West Chicago, IL 60185	
RX ONLY		MFG: CMAP, West Chicago, IL 60185	
Keep this and all medication out of the reach of children			
Directions For Use: See Package Insert			
Store at 20-25°C (68-77°F); excursions permitted to 15-30°C (59-86°F) (see USP Controlled Room Temperature). See USP Controlled Room Temperature. Repackaged by: RemedyRepack Inc., Indiana, PA 15109, 724-465-8762			

SUBVENTE			
Lamotrigine tablet			
<b>Product Information</b>			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC 70518-2645 (NDC 69102-0320)
Route of Administration	ORAL		
<b>Active Ingredient/Active Moiety</b>			
	Ingredient Name	Rank of Strength	Strength
LAMOTRIGINE (UNE 13827698X) LAMOTRIGINE	UN13827698X	LAMOTRIGINE	200 mg
<b>Product Characteristics</b>			
Color	white (tablet or off-white)	Score	2 pieces
Shape	ROUND (Round, flat face beveled-edge)	Size	12mm
Imprint		Imprint Code	264A
Contains			
<b>Packaging</b>			
#	Item Code	Package Description	Marketing Start Date
1	NDC 70518-2645-0	100 in 1 BOX	03/24/2020
1	NDC 70518-2645-1	1 in 1 POUCH Type 0: Not a Combination Product	
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078947	03/24/2020	
<b>Labeler - REMEDY REPACK INC. (025572556)</b>			