

**ALL DAY RELIEF- naproxen sodium tablet, film coated**  
**Kroger Company**

-----  
**Kroger Co. All Day Relief Drug Facts**

**Active ingredient (in each caplet)**

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - you have difficulty swallowing
  - it feels like the pill is stuck in your throat
  - redness or swelling is present in the painful area
  - any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"><li>• take 1 caplet every 8 to 12 hours while symptoms last</li><li>• for the first dose you may take 2 caplets within the first hour</li><li>• do not exceed 2 caplets in any 8- to 12-hour period</li><li>• do not exceed 3 caplets in a 24-hour period</li></ul>
Children under 12 years	<ul style="list-style-type: none"><li>• ask a doctor</li></ul>

**Other information**

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?**

1-800-632-6900

**Principal Display Panel**

COMPARE TO active ingredient of ALEVE® CAPLETS

See side panel

SEE NEW WARNINGS INFORMATION

Strength to Last 12 Hours

All Day Relief

Naproxen Sodium Tablets, USP 220 mg

Pain Reliever - Fever Reducer (NSAID)

Actual Size

\*\*Capsule-Shaped Tablets

EXP.  
: 36878 45 C12

LOT NO.



KEEP CARTON FOR REFERENCE



QUALITY GUARANTEED

COMPARE TO active ingredient of  
**ALEVE® CAPLETS** †See side panel

NDC 30142-368-78

SEE NEW WARNINGS INFORMATION

Strength to Last 12 Hours

# All Day Relief

Naproxen Sodium Tablets, USP 220 mg

Pain Reliever-Fever Reducer (NSAID)



Actual Size

100 CAPLETS\*\* \*\*Capsule-Shaped Tablets

Our  
Pharmacists  
Recommend

### Drug Facts (continued)

Questions or comments?  
1-800-632-6900

†Aleve® is a registered trademark of Bayer HealthCare LLC, Pittsburgh, PA 15205. Bayer HealthCare LLC is not affiliated with The Kroger Co. or this product.

GLUTEN FREE

Distributed by The Kroger Co., Cincinnati, Ohio 45202

QUALITY GUARANTEED

If you are not completely satisfied with this product, return it for a refund or replacement.

Comments or questions? [www.kroger.com](http://www.kroger.com)



#### Drug Facts (continued)

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

#### Directions

■ do not take more than directed  
■ the smallest effective dose should be used  
■ drink a full glass of water with each dose

Adults and children 12 years and older  
■ take 1 caplet every 8 to 12 hours while symptoms last  
■ for the first dose you may take 2 caplets within the first hour  
■ do not exceed 2 caplets in any 8- to 12-hour period  
■ do not exceed 3 caplets in a 24-hour period

Children under 12 years

■ ask a doctor

#### Other information

■ each caplet contains: sodium 21 mg  
■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients** FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, etc, titanium dioxide

**DO NOT USE IF PRINTED FOIL UNDER  
CAP IS BROKEN OR MISSING**

### **Drug Facts**

#### **Active ingredient (in each caplet) Purposes**

Naproxen sodium 220 mg Pain reliever/  
(naproxen 200 mg) (NSAID)\* .....fever reducer  
\*nonsteroidal anti-inflammatory drug

#### **Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

#### **Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
- are age 60 or older
  - have had stomach ulcers or bleeding problems
  - take a blood thinning (anticoagulant) or steroid drug
  - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
  - have 3 or more alcoholic drinks every day while using this product
  - take more or for a longer time than directed ▶



### **Drug Facts (continued)**

#### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

#### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug



#### **When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear ▶





## ALL DAY RELIEF

naproxen sodium tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30 142-368
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

### Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONES (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	BLUE (Light Blue)	Score	no score
Shape	CAPSULE (oval)	Size	12mm
Flavor		Imprint Code	L368
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30 142-368-71	1 in 1 CARTON		
1		50 in 1 BOTTLE		

2	NDC:30 142-368-62	1 in 1 CARTON		
2		24 in 1 BOTTLE		
3	NDC:30 142-368-78	1 in 1 CARTON		
3		100 in 1 BOTTLE		
4	NDC:30 142-368-90	500 in 1 BOTTLE		
5	NDC:30 142-368-76	1 in 1 CARTON		
5		120 in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	01/16/1997	

**Labeler** - Kroger Company (006999528)

Revised: 12/2012

Kroger Company